



Shirley A. Jones

ECG Success

Exercises in ECG Interpretation



ECG Success

Exercises in ECG Interpretation



ECG Success

Exercises in ECG Interpretation

Shirley A. Jones, MS Ed, MHA, EMT-D

Emergency Medical Services Educator

Arrhythmia Instructor

Riverview Hospital

Noblesville, Indiana

Basic Life Support Instructor

American Heart Association

Advanced Cardiac Life Support Instructor

American Heart Association



F. A. DAVIS COMPANY • Philadelphia

F. A. Davis Company
1915 Arch Street
Philadelphia, PA 19103
www.fadavis.com

Copyright © 2008 by F. A. Davis Company

Copyright © 2008 by F. A. Davis Company. All rights reserved. This product is protected by copyright. No part of it may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise, without written permission from the publisher.

Printed in the United States of America

Last digit indicates print number: 10 9 8 7 6 5 4 3 2 1

Acquisitions Editor: Lisa B. Deitch
Project Editor: Ilysa Richman, Padraic J. Maroney
Director of Content Development: Darlene D. Pedersen
Art and Design Manager: Carolyn O'Brien

As new scientific information becomes available through basic and clinical research, recommended treatments and drug therapies undergo changes. The author(s) and publisher have done everything possible to make this book accurate, up to date, and in accord with accepted standards at the time of publication. The author(s), editors, and publisher are not responsible for errors or omissions or for consequences from application of the book, and make no warranty, expressed or implied, in regard to the contents of the book. Any practice described in this book should be applied by the reader in accordance with professional standards of care used in regard to the unique circumstances that may apply in each situation. The reader is advised always to check product information (package inserts) for changes and new information regarding dose and contraindications before administering any drug. Caution is especially urged when using new or infrequently ordered drugs.

Library of Congress Cataloging-in-Publication Data

Jones, Shirley A.

ECG success : exercises in ECG interpretation / Shirley A. Jones.

p. ; cm.

ISBN-13: 978-0-8036-1577-9

ISBN-10: 0-8036-1577-9

1. Electrocardiography—Interpretation—Problems, exercises, etc. I. Title.

[DNLM: 1. Electrocardiography—methods—Problems and Exercises. 2. Arrhythmia—diagnosis—Problems and Exercises. WG 18.2 J76e 2007]

RC683.5.E5J575 2007

616.1'2075470076—dc22

2007017019

Authorization to photocopy items for internal or personal use, or the internal or personal use of specific clients, is granted by F. A. Davis Company for users registered with the Copyright Clearance Center (CCC) Transactional Reporting Service, provided that the fee of \$.10 per copy is paid directly to CCC, 222 Rosewood Drive, Danvers, MA 01923. For those organizations that have been granted a photocopy license by CCC, a separate system of payment has been arranged. The fee code for users of the Transactional Reporting Service is: 8036-1577-07 0 + \$.10.



Dedication

To the memory of my father George Francis Jones who was and still is my hero in life. And, to my sister Virginia Kelleher, MD, for all her love and support. Also to my best buddies who have always given me unconditional love, Zachary, Chelsea, Little Zachary, Darby, Spirit, and Francis.

Shirley A. Jones



Preface

No one is born knowing how to read ECG strips. We learn to do many things in a lifetime, and nearly all of them get better with practice. If you're planning to use this book, ECG isn't completely new to you—you have a good idea of what's involved in generating and interpreting a tracing.

ECG Success covers all the information you will need—anatomy and physiology, practice, and case scenarios, and relevant emergency care—to help you feel competent and in control, whether the situation involves an emergency or just a nonthreatening ECG. This book has staying power. You will find its content useful across a spectrum of situations, from classroom study through clinical experience and later in actual practice.

Pattern recognition lies at the heart of ECG interpretation. This skill develops with experience, gained through repetition and variety. You need to see the same patterns over and over again, but you also need to see as great a diversity as possible. In *ECG Success* you'll find more than 550 ECG tracings.

The book is organized into four units. First, two introductory chapters review the background information you need for working with ECG. Chapter One discusses heart anatomy and physiology, including biomechanics and electrophysiology. Chapter Two gives you the basics of ECG: limb and chest leads, electrode placement, cable connections, components of a tracing, rhythm strip analysis, and more.

In Unit Two, seven chapters explain and illustrate the different types of rhythm, some dangerous, others merely troublesome, and a few even normal. Each of these chapters gives you a group of nine practice strips to analyze, with the answers given at the end of the chapter. All types of arrhythmias are discussed and illustrated: sinus, atrial, junctional, and ventricular; atrioventricular and bundle branch blocks; artifact; and

artificial pacemaker rhythm. The section ends with a chapter on myocardial infarction and the 12-lead ECG.

The chapter practice strips will warm you up for Unit Three, the working core of the book. You'll find four test chapters with a total of 300 strips and the answers given at the end of each chapter so you can check your work. In case you're hungry for more, the two chapters in Unit Four comprise eleven real-life case studies, followed by multiple-choice questions and illustrated by more ECG strips. Four appendices round out the book: Healthcare Provider Guidelines for Cardiopulmonary Resuscitation, Advanced Cardiac Life Support Protocols, Emergency Medications, and Emergency Medical Skills.

As you page through this book you'll find some special features to guide you. In Units One and Two, frequent Clinical Tips provide valuable information on how an arrhythmia can affect the patient. Hints on rhythm interpretation appear throughout the first practice strip chapter.

I couldn't have written this book without building up a track record of my own ECG successes. The secret: I had a good instructor who was patient and explained everything in detail. She kept emphasizing that we had to follow every step when analyzing a rhythm; shortcuts are dangerous because you can miss critical details on the rhythm strip. Then we had to practice, practice, practice. That repetition, combined with careful attention to every step, was the real key to my success.

Take your time now, and use *ECG Success* to improve your skills. Once you run into a genuine emergency you will have only minutes, or less, to interpret the ECG correctly and ensure the right treatment for the patient.

Shirley A. Jones



Consultants

Dawn McKay, RN, MSN, CCRN

Critical Care Instructor
Assistant Professor of Nursing
Liberty University
Lynchburg, Virginia
Basic Life Support Instructor,
American Heart Association
Advanced Cardiac Life Support Instructor, American
Heart Association
Trauma Nursing Core Course Instructor, Emergency
Nurses Association
Fundamental Critical Care Support Instructor, Society
of Critical Care Medicine

Carmen J. Petrin, MS, APRN, BC

Nurse Practitioner
New England Heart Institute at
Catholic Medical Center
Manchester, New Hampshire
Former Critical Care Educator
Catholic Medical Center
Manchester, New Hampshire
Basic Life Support Instructor,
American Heart Association
Advanced Cardiac Life Support Instructor,
American Heart Association
Pediatric Advanced Life Support Instructor,
American Heart Association
National Advanced Cardiac Life Support
Faculty, American Heart Association
Pediatric Advanced Life Support Training
Center Faculty, American Heart Association
Basic Life Support Training Center
Faculty, American Heart Association

Shirley A. Jones has worked in the field of emergency medical services for more than 30 years. She received her Master of Science in Education and her Master of Health Administration degrees from Indiana University. She has been awarded five first-place honors in tri-state and state-wide advanced life support competitions, served on the faculty of national conferences, and won honors from the Medical Writers Association for two textbooks. She is an accomplished writer and educator in the fields of electrocardiology and pharmacological and mechanical therapy. She welcomes the comments, criticisms, and ideas of readers for the improvement of future editions.



Reviewers

Lori Baker, RN

Surgical Nurse Educator
Lions Gate Hospital
North Vancouver, Canada

Carole Berube, MA, MSN, RN

Professor Emerita in Nursing
Bristol Community College
Fall River, Massachusetts

Daryl Boucher, MSN, RN

Flight Nurse/Nursing and Allied
Health Faculty
Aroostook Medical Center/Northern
Maine Community College
Presque Isle, Maine

Carmen Carpenter, RN, MS, CMA

Department Chair, Allied Health
Sciences and Medical Assisting
South University
West Palm Beach, Florida

**Barbara Chamberlain,
MSN, APRN, BC**

Critical Care Clinical Nurse
Specialist
Kennedy Health System
Turnersville, New Jersey

Pam Chambers, MPH, PA-C

Associate Professor, Physician
Assistant Program
Des Moines University
Des Moines, Iowa

Julie Chew, RN, MS, PhD

Clinical Educator
Sacred Heart Medical Center
Eugene, Oregon

Jeff Chianfagna, BS, MA, PA

Clinical Physician Assistant and
Academic Instructor/Faculty
Pace Lenox Hill Hospital Physician
Assistant Program
New York, New York

Nancy Edge, RN, BSN

Clinical Educator, Cardiac Surgery
Vancouver General Hospital
Vancouver, Canada

Linda Latham, RN, BSN, MA

Lead Instructor
Forsyth Technical Community
College
Winston-Salem, North Carolina

Cindy Light, RN, MSN, CEN

Nursing Instructor
Baker University School of Nursing
Topeka, Kansas

Dawn McKay, RN, MSN, CCRN

Assistant Professor
Liberty University
Lynchburg, Virginia

Susan Moore, PhD, RN

Professor of Nursing
New Hampshire Community
Technical College
Manchester, New Hampshire

Deborah Opacic, EdD, PA-C

Professor, Clinical Educator
Duquesne University
Pittsburgh, Pennsylvania

Patricia Richards, RN, MSN, CCRN

Faculty, Nursing Department
Central Maine Community College
Auburn, Maine

Catherine Richmond, RN, BSN, MSN

Professor, Nursing
SUNY Alfred State College
Alfred, New York

Robert Spears, MPAS, PA-C

Assistant Professor
The University of Findlay
Findlay, Ohio

Walt Stoy, PhD

Director of Education
Center for Emergency Medicine
Pittsburgh, Pennsylvania

Debbie Sullivan, PhD, PA-C

Assistant Professor/PA Program
Faculty/Interim Director
Midwestern University
Glendale, Arizona

Rita Tomasewski, MSN, ARCNP

Cardiovascular Clinical Nurse
Specialist
St. Francis Health Center
Topeka, Kansas

Marilyn Turner, RN, CMA

Medical Assisting Program Director
Ogeechee Technical College
Statesboro, Georgia



Contents

UNIT ONE

Heart Structure and Electrical Activity

<i>Chapter 1:</i>	ANATOMY AND PHYSIOLOGY OF THE HEART 2
	Overview 2
	<i>Anatomy of the Heart 2</i>
	Layers of the Heart 2
	Heart Valves 2
	Heart Chambers and Great Vessels 3
	Coronary Vessel Circulation 3
	<i>Anatomy of the Cardiovascular System 4</i>
	Blood Vessel Structures 4
	Arterial Circulation 4
	Venous Circulation 5
	<i>Physiology of the Heart 5</i>
	Mechanical Physiology 5
	Electrophysiology 6
<i>Chapter 2:</i>	THE ELECTROCARDIOGRAM 9
	Overview 9
	Limb Leads 10
	Standard Chest Leads 10
	Cable Connections 11
	Electrode Placement Using a Three-wire Cable 11
	Electrode Placement Using a Five-wire Cable 12
	Modified Chest Leads 12
	The Right-sided 12-Lead ECG 12
	The 15-Lead ECG 13
	Recording of the ECG 13
	Components of an ECG Tracing 14
	Methods for Calculating Heart Rate 14
	Rhythm Strip Analysis 16
	Instructions for Analyzing ECG Practice and Test Strips 16

UNIT TWO

Rhythms and Their Analysis

<i>Chapter 3:</i>	SINOATRIAL NODE ARRHYTHMIAS 20
	Normal Sinus Rhythm (NSR) 20
	Sinus Bradycardia 21
	Sinus Tachycardia 21
	Sinus Arrhythmia 22
	Sinus Pause (Sinus Arrest) 22
	Sinoatrial (SA) Block 23
	ECG Practice Strips 23
	Answers to Chapter 3 ECG Practice Strips 26
<i>Chapter 4:</i>	ATRIAL ARRHYTHMIAS 28
	Wandering Atrial Pacemaker (WAP) 28
	Multifocal Atrial Tachycardia (MAT) 29
	Premature Atrial Contraction (PAC) 29
	Atrial Tachycardia 30
	Supraventricular Tachycardia (SVT) 30
	Paroxysmal Supraventricular Tachycardia (PSVT) 31
	Atrial Flutter 31
	Atrial Fibrillation 32
	Wolff-Parkinson-White (WPW) Syndrome 32
	ECG Practice Strips 33
	Answers to Chapter 4 ECG Practice Strips 36
<i>Chapter 5:</i>	JUNCTIONAL ARRHYTHMIAS 37
	Junctional Rhythm 37
	Accelerated Junctional Rhythm 38
	Junctional Tachycardia 38
	Junctional Escape Beat 38

Premature Junctional Contraction (PJC)	39	Artificial Pacemaker Rhythm	64
ECG Practice Strips	39	Single-chamber Pacemaker Rhythm—Atrial	65
Answers to Chapter 5 ECG Practice Strips	42	Single-chamber Pacemaker Rhythm—Ventricular	65
Chapter 6: VENTRICULAR ARRHYTHMIAS 44		Dual-chamber Pacemaker Rhythm—Atrial and Ventricular	65
Idioventricular Rhythm	44	Pacemaker Malfunctions	65
Accelerated Idioventricular Rhythm	45	Failure to Capture	66
Premature Ventricular Contraction (PVC)	45	Failure to Sense	66
Premature Ventricular Contraction: Uniform	46	Oversensing	66
Premature Ventricular Contraction: Multiform	46	ECG Practice Strips	67
Premature Ventricular Contraction: Ventricular Bigeminy	46	Answers to Chapter 8 ECG Practice Strips	70
Premature Ventricular Contraction: Ventricular Trigeminy	46	Chapter 9: ARTIFACT 71	
Premature Ventricular Contraction: Ventricular Quadrigeminy	47	Loose Electrical Connection	71
Premature Ventricular Contraction: Couplets	47	Variation with Respiration	71
Premature Ventricular Contraction: R on T Phenomenon	47	60-cycle Interference	72
Premature Contraction: Interpolated PVC	48	Muscle Artifact	72
Ventricular Tachycardia (VT): Monomorphic	48	ECG Practice Strips	72
Ventricular Tachycardia (VT): Polymorphic	49	Answers to Chapter 9 ECG Practice Strips	75
Torsade de Pointes	49	Chapter 10: THE 12-LEAD ECG AND ACUTE MYOCARDIAL INFARCTION 77	
Ventricular Fibrillation (VF)	50	Troubleshooting ECG Problems	77
Pulseless Electrical Activity (PEA)	50	R Wave Progression	78
Asystole	51	Electrical Axis Deviation	78
ECG Practice Strips	51	Ischemia, Injury, and Infarction	78
Answers to Chapter 6 ECG Practice Strips	54	Progression of an Acute Myocardial Infarction	79
Chapter 7: ATRIOVENTRICULAR AND BUNDLE BRANCH BLOCKS 56		ST Segment Elevation and Depression	79
First-degree AV Block	56	The Normal 12-lead ECG	80
Second-degree AV Block: Type I (Mobitz I or Wenckebach)	57	Anterior Myocardial Infarction	80
Second-degree AV Block: Type II (Mobitz II)	57	Inferior Myocardial Infarction	80
Third-degree AV Block	58	Lateral Myocardial Infarction	81
Bundle Branch Block (BBB)	58	Septal Myocardial Infarction	81
ECG Practice Strips	59	Posterior Myocardial Infarction	82
Answers to Chapter 7 ECG Practice Strips	62	Left Bundle Branch Block	82
Chapter 8: ARTIFICIAL CARDIAC PACEMAKERS 63		Right Bundle Branch Block	82
Artificial Pacemaker	63	UNIT THREE ECG Practice Tests	
Pacemaker Modes	64	Chapter 11: ECG PRACTICE TEST ONE 86	
Understanding Pacemaker Codes	64	Test Strip Section One	86
		Answers to Practice Test One	115
		Chapter 12: ECG PRACTICE TEST TWO 122	
		Test Strip Section Two	122
		Answers to Practice Test Two	147
		Chapter 13: ECG PRACTICE TEST THREE 154	
		Test Strip Section Three	154
		Answers to Practice Test Three	179

<i>Chapter 14:</i>	ECG PRACTICE TEST FOUR	186
	Test Strip Section Four	186
	Answers to Practice Test Four	211

UNIT FOUR Case Studies

<i>Chapter 15:</i>	ECG CASE STUDIES ONE	220
	Case Study One	220
	Case Study Two	222
	Case Study Three	223
	Case Study Four	226
	Case Study Five	228
	Answers to Case Study One	229
	Answers to Case Study Two	229
	Answers to Case Study Three	230
	Answers to Case Study Four	231
	Answers to Case Study Five	231

<i>Chapter 16:</i>	ECG CASE STUDIES TWO	232
	Case Study One	232
	Case Study Two	233
	Case Study Three	235
	Case Study Four	237
	Case Study Five	239
	Case Study Six	241
	Answers to Case Study One	242
	Answers to Case Study Two	243
	Answers to Case Study Three	243
	Answers to Case Study Four	244
	Answers to Case Study Five	244
	Answers to Case Study Six	244

APPENDICES

<i>Appendix A:</i>	HEALTHCARE PROVIDER GUIDELINES FOR CARDIOPULMONARY RESUSCITATION (CPR)	247
	CPR Healthcare Provider Skill Performance	247
	CPR: Adult	247
	CPR: Child	248
	CPR: Infant	249
	OBSTRUCTED AIRWAY:	
	Conscious Adult or Child	249

OBSTRUCTED AIRWAY:	
Conscious Infant	250
OBSTRUCTED AIRWAY:	
Unconscious Adult	250
OBSTRUCTED AIRWAY:	
Unconscious Child	251
OBSTRUCTED AIRWAY:	
Unconscious Infant	251
CPR and Obstructed Airway Positions	252

<i>Appendix B:</i>	ADVANCED CARDIAC LIFE SUPPORT PROTOCOLS	253
	Ventricular Fibrillation (VF) or Pulseless Ventricular Tachycardia (VT)	253
	Pulseless Electrical Activity (PEA)	254
	Asystole	254
	Acute Coronary Syndrome (ACS)	255
	Bradycardia	255
	Tachycardia—Unstable	256
	Narrow-complex Tachycardia—Stable Regular Rhythm	257
	Narrow-complex Tachycardia—Stable Irregular Rhythm	257
	Wide-complex Tachycardia—Stable Regular Rhythm	258
	Wide-complex Tachycardia—Stable Irregular Rhythm	258

<i>Appendix C:</i>	EMERGENCY MEDICATIONS	259
--------------------	------------------------------	------------

<i>Appendix D:</i>	EMERGENCY MEDICAL SKILLS	265
	Defibrillation	265
	Cardioversion (Synchronized)	266
	Transcutaneous Pacing	267
	Carotid Sinus Massage (Vagal Maneuver)	267
	<i>Abbreviations Used in Text</i>	269
	<i>Selected References</i>	271
	<i>Index</i>	273

Heart Structure and Electrical Activity



Anatomy and Physiology of the Heart

OVERVIEW

Cardiovascular disease is a common cause of medical problems. Patients may present with symptoms ranging from chest pain to sudden collapse. You should already be familiar with the most important causes of cardiovascular emergencies: angina, congestive heart failure, acute myocardial infarction, pulmonary edema, cardiogenic shock, arrhythmias, hypertensive emergency, and cardiac arrest. This chapter reviews the anatomy, physiology, and electrical conduction system of the heart.

ANATOMY OF THE HEART

The heart, located in the mediastinum, is the central structure of the cardiovascular system. It is protected by the bony structures of the sternum anteriorly, the spinal column posteriorly, and the rib cage (FIG. 1-1).




Figure 1.1 ■ Location of the heart.

This fist-sized muscular organ is roughly conical. The base of the cone is at the top of the heart and the apex (the pointed part) is at the bottom. The heart is rotated slightly counterclockwise, with the apex tipped anteriorly so that the back surface of the heart actually lies over the diaphragm.



Clinical Tip:

The cone-shaped heart has its tip (apex) just above the diaphragm to the left of the midline. This is why we may think of the heart as being on the left side—the strongest beat can be heard or felt there.

LAYERS OF THE HEART

The heart is composed of several different layers of tissue (FIG. 1-2). Surrounding the heart itself is a protective sac called the pericardium. This double-walled sac has an inner, serous (visceral) layer and an outer, fibrous (parietal) layer. Between these layers is the pericardial cavity, which contains a small amount of lubricating fluid to prevent friction during heart contraction. The layers of the heart wall itself include the epicardium, or outermost layer; the myocardium, the thick middle layer of cardiac muscle; and the endocardium, the smooth layer of connective tissue that lines the inside of the heart.

Myocardial tissue is a special type of contractile tissue found only in the heart. Although it is similar in appearance to skeletal muscle tissue, myocardial tissue has some unique structural and electrical properties. These properties are described more fully in the discussion of electrophysiology.

HEART VALVES

To prevent the backflow of blood during cardiac contraction, the atria and ventricles are separated from




Figure 1.2 ■ Layers of the heart.

each other by two sets of valves composed of endocardial and connective tissue (FIG. 1-3). The fibrous connective tissue prevents enlargement of valve openings and anchors valve flaps.

The first set of heart valves, the atrioventricular (AV) valves, is located between each atrium and ventri-




Figure 1.3 ■ Valves of the heart. The atria have been removed in this superior view.

cle. The tricuspid valve, which is between the right atrium and ventricle, derives its name from its construction of three feathery leaflets, or cusps. The left AV valve, which has only two cusps, is called the mitral valve. The AV valves open to allow ventricular filling when the intra-atrial pressure exceeds the intraventricular pressure during atrial contraction. The onset of ventricular contraction creates pressure to close the AV valves.

The other set of valves, called semilunar valves, functions by similar pressure changes and prevents the flow of blood back into the ventricles after contraction. The two semilunar valves are the pulmonic valve, located in the outflow tract from the right ventricle to the pulmonary artery, and the aortic valve, between the left ventricle and the aorta.

HEART CHAMBERS AND GREAT VESSELS

The heart is a hollow muscle with an internal skeleton of connective tissue that creates four separate chambers (FIG. 1-4). The superior chambers of the heart are the right and left atria. These chambers primarily collect blood as it enters the heart and help fill the lower chambers.

The more thickly muscled lower chambers of the heart are called ventricles. These are the primary pumping chambers, the left having a thicker myocardial layer than the right. Vertical walls, composed of connective and muscle tissue, separate the two atria and the two ventricles. These walls are called the interatrial septum and the interventricular septum, respectively.

The pulmonary artery, the aorta, the superior and inferior vena cava, and the pulmonary veins are the largest blood vessels in the heart and are often referred to collectively as the great vessels (FIG. 1-4).

CORONARY VESSEL CIRCULATION

The coronary arteries and veins provide the blood supply to the heart muscle and the electrical conduction




Figure 1.4 ■ Heart chambers and great vessels.




Figure 1.5 ■ Coronary vessels. (A) Coronary vessels in anterior view. (B) Coronary vessels in posterior view.

system. The coronary arteries, left and right, are the first to branch off the aorta, just above the leaflets of the aortic valve (Fig. 1-5). The left coronary artery has two major branches. The anterior descending branch runs along the anterior surface of the heart, while the circumflex branch courses in the groove between the left atrium and ventricle to the posterior surface of the heart. These two branches supply arterial blood to the left ventricle, interventricular septum, part of the right ventricle, and certain electrical conduction structures in those areas.

The right coronary artery arises from the aorta and courses along the right AV groove to the posterior surface of the heart. This artery supplies the right atrium, right ventricle, and part of the left ventricle, in addition to certain electrical conduction structures. The heart's arterial anatomy provides the left ventricle with a dual blood supply, from both coronary arteries.

The coronary veins correspond in distribution to the coronary arteries. They drain venous blood into the right atrium. The largest of these veins, the coronary sinus, provides venous drainage of the left ventricle.



Clinical Tip:

A protective feature of the coronary vessels is that many interconnections, or anastomoses, exist between the arterioles of the coronary arteries, allowing for development of collateral circulation, if needed.

ANATOMY OF THE CARDIOVASCULAR SYSTEM

The cardiovascular system is a closed system consisting of blood vessels and the heart. Arteries and veins are

connected by smaller structures in which electrolytes are exchanged across cell membranes.

BLOOD VESSEL STRUCTURES

The systemic, or peripheral, circulation is composed of a circuit of blood vessels that transport substances needed for cellular metabolism to body systems and remove the waste products of metabolism from those same tissues. With the exception of the interconnecting capillaries, the anatomy of all blood vessel walls is a similar three-layer design (Fig. 1-6). The tunica intima is the smooth single-cell layer that lines the inside of all blood vessel walls. The middle layer of elastic fibers and muscle, the tunica media, gives strength and recoil. Contraction or relaxation of this muscle layer varies the diameter of the blood vessel lumen, the cavity through which blood flows. Finally, the tunica externa, a tough outer layer of fibrous tissue, protects the blood vessel from damage.

ARTERIAL CIRCULATION

Arteries carry blood away from the heart and, with the exception of the pulmonary artery, transport oxygenated blood. Arteries carry blood under high pressure and, therefore, are equipped with a much thicker medial layer than other blood vessels. Major arteries of the body to recognize include the aorta and the subclavian, internal and external carotid, axillary, brachial, radial, common iliac, and femoral arteries (Fig. 1-7).




Figure 1.6 ■ Blood vessel structures.




Figure 1.7 ■ Arterial circulation.

VENOUS CIRCULATION

With the exception of the pulmonary veins, veins carry blood that is low in oxygen and high in carbon dioxide content. Veins carry blood under much lower pressure than arteries. Movement of this low-pressure blood, often against gravity, is aided by the “milking” action of large muscles that surround veins, particularly in the legs, and a series of intermittent one-way valves in the veins to prevent backflow of blood between heart contractions. Major veins of the body include the superior and inferior vena cava and the internal and external jugular, subclavian, axillary, iliac, and femoral veins (FIG. 1-8).

PHYSIOLOGY OF THE HEART

Normal blood flow through the heart begins at the right atrium, which receives systemic venous blood from the superior and inferior venae cavae (see Fig. 1-4). Blood passes from the right atrium, across the tricuspid valve, to the right ventricle. It is then pumped across the pulmonary valve into the pulmonary artery.

Outside the heart, the two branches (left and right) of the pulmonary artery distribute blood to the lungs for gas exchange in the pulmonary capillaries. Oxygenated blood returns to the heart’s left atrium through




Figure 1.8 ■ Venous circulation.

four pulmonary veins. After passing across the mitral valve, blood enters the left ventricle, where it is pumped across the aortic valve and then enters the coronary arteries and the peripheral circulation through the aorta.

MECHANICAL PHYSIOLOGY

The complete cycle of mechanical pumping of blood through the heart and pulmonary circulation is referred to as the cardiac cycle. In this cycle, the right and left atria contract just before the beginning of right and left ventricular contraction. Much of the blood flow from the atria to the ventricles occurs by gravity, but atrial contraction is necessary to fill the ventricles to maximum capacity. The right and left ventricles contract simultaneously while the atria relax. The resulting pressure closes the AV valves, opens the aortic and pulmonary valves, and propels blood into the pulmonary and systemic circulation.

The contraction phase of the cardiac cycle is called systole, which generally refers to ventricular contraction versus atrial contraction. Diastole is the relaxation

phase of the cardiac cycle, when the ventricles are filling (FIG. 1-9). This phase lasts much longer than systole.



Clinical Tip:

Increases in heart rate reduce the length of diastole more significantly than that of systole. The duration of the diastolic phase is important, as this is when about 70% of the coronary artery flow occurs and complete filling of the ventricles takes place.

The amount of blood ejected from either ventricle with a single contraction is called the stroke volume. Stroke volume is determined and affected by three factors: preload, afterload, and cardiac contractility.

Preload can be thought of as the pressure under which the ventricle fills. This pressure is influenced by the amount of venous blood return. A feature of myocardial muscle is that the more it is stretched (up to a limit), the greater its force of contraction. Therefore, stroke volume can be increased considerably by increasing the blood volume that fills the ventricles and thus increasing the amount of myocardial muscle fiber stretch. This concept is known as Starling's law of the heart. Afterload, or the resistance against which the ventricles contract, also influences stroke volume. Afterload is determined by systemic arterial resistance. Cardiac contractility is the third major determinant of stroke volume. It is the intrinsic state of the heart muscle's force of contraction, also called the heart's contractile, or inotropic, state.

Stroke volume (SV) times heart rate (HR) determines the heart's cardiac output (CO), the amount of blood pumped through the circulatory system per minute:

$$CO = SV \times HR$$

An increase in stroke volume alone can improve the cardiac output. However, heart rate also has great impact. Rate increases in the healthy heart can improve the cardiac output up to threefold.

ELECTROPHYSIOLOGY

Automaticity

Arrhythmia interpretation is based on an understanding of the normal anatomy and physiology of electrical conduction in the heart. Myocardial fibers possess




Figure 1.9 ■ Systolic and diastolic phases in the heart.

highly specialized electrical properties, in addition to the mechanical property of contractility. Automaticity is the ability to generate an electrical impulse inde-

pendently of stimulation by the nervous system or any other source. This property is unique to certain cardiac cells called pacemaker cells.

Excitability and Conductivity

Two other electrical properties, excitability and conductivity, are shared by all myocardial cells. Excitability is the ability of cells to respond to electrical stimulation. Conductivity is the ability to pass or propagate an electrical impulse from cell to cell through the heart. These three properties are constantly involved in the electrical conduction system of the heart.

The heart's electrical conduction system is a network of structures that allows electrical impulses to spread through the heart much faster than if they had to spread through muscle cells alone. The structures of the conduction system, in sequence of normal electrical conduction, are shown in Figure 1-10 and listed below:

Sinoatrial (SA) node. This node is the dominant pacemaker of the heart, located in the upper portion of the right atrium. Intrinsic rate is 60–100 bpm.

Internodal pathways. These cells direct electrical impulses between the SA and AV nodes and spread them across the atrial muscle.

Atrioventricular (AV) node. This node is part of an area called AV junctional tissue, which includes some surrounding tissue plus the connected bundle of His. Although AV junctional tissue contains pacemaker cells, none are thought to exist in the AV node itself. The AV node slows conduction, creating a slight delay




Figure 1.10 ■ Conduction system of the heart.

before electrical impulses are carried to the ventricles. Intrinsic rate is 40–60 bpm.

Bundle of His. Located at the top of the interventricular septum, this bundle of fibers extends directly from the AV node and connects the atria and ventricles electrically.

Bundle branches. The bundle of His splits into two conduction paths called the right and left bundle branches. These bundles carry electrical impulses at high speed to the tissue of the interventricular septum, and to each ventricle simultaneously.

Purkinje system. The bundle branches terminate with this network of fibers, which spread electrical impulses rapidly throughout the ventricular walls. Intrinsic rate is 20–40 bpm.

The creation of electrical impulses and the spread of impulses through the electrical conduction system occur through a process called depolarization. Chemical pumps in the cell walls alter the precise concentration of electrolytes maintained inside and outside the cell. During depolarization, the electrical charge of a cell is changed by the electrolyte concentration shift on either side of the cell membrane. This change in electrical charge stimulates the muscle fiber to contract. A resting, or “polarized,” cell is normally more electrically negative on the inside of the cell wall than on the outside (FIG. 1-11).

Electrical stimulation, however, changes the permeability of the cell wall and allows positively charged ions, particularly sodium (Na^+), to move into the cell. The rush of sodium, along with the slower influx of calcium (Ca^{++}), causes the inside of the cell to change from negative to positive. The cell is then said to be depolarized. The response of the muscle to this electrical charge is contraction. Because of conductivity, this process of depolarization moves rapidly from cell to cell in the conduction pathway and throughout the muscle cells of the heart (FIG. 1-12).

After depolarization, myocardial cells must return to their resting state of internal negativity for further depolarization to occur. The proper distribution of electrolytes is re-established by the cell wall chemical pumps, which pump sodium (Na^+) out of the cell and return potassium (K^+) into the cell. This process of re-establishing the internal negative charge of the cell is called repolarization.




Figure 1.11 ■ The depolarization process. (A) A single cell has depolarized. (B) A wave propagates from cell to cell (C) until all are depolarized. (D) Repolarization that restores each cell's normal polarity.




Figure 1.12 ■ Progression of depolarization through the heart.

The Electrocardiogram

OVERVIEW

In Chapter 1 you learned that movement of electrolytes across the membranes of myocardial cells (depolarization and repolarization) produces a flow of electrical current and creates an electrical field. Depolarization and repolarization can be seen on the electrocardiogram (ECG) (FIG. 2-1).



Clinical Tip:

It is important to keep in mind that the ECG shows only electrical activity; it tells us nothing about how well the heart is working mechanically.

The body acts as a giant conductor of electrical current. Electrical activity that originates in the heart can be detected on the body's surface through an electrocardiogram (ECG). Electrodes are applied to the skin to measure voltage changes in the cells between the electrodes. These voltage changes are amplified and visually displayed on an oscilloscope and graph paper. This chapter focuses on the ECG and its analysis and interpretation.

The following summarizes a basic ECG:

- An ECG is a series of waves and deflections recording the heart's electrical activity from a certain "view."
- Many views, each called a lead, monitor voltage changes between electrodes placed in different positions on the body.
- Leads I, II, and III are bipolar leads, which consist of two electrodes of opposite polarity (positive and negative). The third (ground) electrode minimizes electrical activity from other sources.
- Leads aVR, aVL, and aVF are unipolar leads and consist of a single positive electrode and a reference

point (with zero electrical potential) that lies in the center of the heart's electrical field.

- Leads V_1 through V_6 are unipolar leads and consist of a single positive electrode with a negative reference point, found at the electrical center of the heart.
- Voltage changes are amplified and visually displayed on an oscilloscope and graph paper.
- An ECG tracing looks different in each lead because the recorded angle of electrical activity changes with each lead.




Figure 2.1 ■ Correlation of depolarization and repolarization with the ECG.

- Several different angles allow a more accurate perspective than a single one would.
- The ECG machine can be adjusted to make any skin electrode positive or negative. The polarity depends on which lead the machine is recording.
- A cable attached to the patient is divided into several different-colored wires: three, four, or five for monitoring purposes, or ten for a 12-lead ECG.
- Incorrect placement of electrodes may turn a normal ECG tracing into an abnormal one.



Clinical Tip:

Patients should be treated according to their symptoms, not merely their ECG tracing.



Clinical Tip:

To obtain a 12-lead ECG, four wires are attached to each limb and six wires are attached at different locations on the chest. The total of ten wires provides 12 views (12 leads).

LIMB LEADS

Electrodes are placed on the right arm (RA), left arm (LA), right leg (RL), and left leg (LL). With only four electrodes, six leads are viewed (FIG. 2-2). These leads include the standard leads—I, II, and III—and the augmented leads—aVR, aVL, and aVF.




Figure 2.2 ■ Standard limb lead electrode placement.




Figure 2.3 ■ Standard leads: I, II, and III.

Table 2.1 ■ ELEMENTS OF STANDARD LEADS

Lead	Positive Electrode	Negative Electrode	View of Heart
I	LA	RA	Lateral
II	LL	RA	Inferior
III	LL	LA	Inferior

Standard Leads

Leads I, II, and III make up the standard leads. If electrodes are placed on the right arm, left arm, and left leg, three leads are formed (FIG. 2-3). If you draw an imaginary line between each of these electrodes, an axis is formed between each pair of leads. The axes of these three leads form an equilateral triangle with the heart in the center (Einthoven's triangle). TABLE 2-1 shows the composition of the standard leads.

Augmented Leads

Leads aVR, aVL, and aVF make up the augmented leads (FIG. 2-4). Each letter of an augmented lead refers to a specific term: a = augmented, V = voltage, R = right arm, L = left arm, F = foot (the left foot). TABLE 2-2 shows the composition of the augmented leads.

STANDARD CHEST LEADS

The chest leads are identified as V_1 , V_2 , V_3 , V_4 , V_5 , and V_6 (FIG. 2-5). Each electrode placed in a "V" position is positive. TABLE 2-3 shows the composition of the chest leads.




Figure 2.4 ■ Augmented leads: aVR, aVL, and aVF.

Table 2.2 ■ ELEMENTS OF AUGMENTED LEADS

Lead	Positive Electrode	View of Heart
aVR	RA	None
aVL	LA	Lateral
aVF	LL	Inferior

Table 2.3 ■ ELEMENTS OF CHEST LEADS

Lead	Positive Electrode Placement	View of Heart
V ₁	Fourth intercostal space to right of sternum	Septum
V ₂	Fourth intercostal space to left of sternum	Septum
V ₃	Directly between V ₂ and V ₄	Anterior
V ₄	Fifth intercostal space at left mid-clavicular line	Anterior
V ₅	Level with V ₄ at left anterior axillary line	Lateral
V ₆	Level with V ₅ at left midaxillary line	Lateral




Figure 2.5 ■ Chest leads: V₁, V₂, V₃, V₄, V₅, and V₆.

Table 2.4 ■ MONITORING CABLE CONNECTIONS

U.S.	Connect to	Europe
White	Right arm	Red
Black	Left arm	Yellow
Red	Left leg	Green
Green	Right leg	Black
Brown	Chest	White

CABLE CONNECTIONS

Before you attach the cable to the patient, you need to know whether the cable is an American or European cable. Improper placement of the connections can give incorrect ECG recordings. The colors of the wires differ as shown in TABLE 2-4.

ELECTRODE PLACEMENT USING A THREE-WIRE CABLE

A three-wire patient cable is used to monitor the standard leads, I, II, and III (FIG. 2-6).




Figure 2.6 ■ Three-wire patient cable.



Clinical Tip:

Lead II is commonly called a monitoring lead. It provides information on heart rate, regularity, conduction time, and ectopic beats. The presence or location of an acute myocardial infarction (MI) should be further diagnosed with a 12-lead ECG.

ELECTRODE PLACEMENT USING A FIVE-WIRE CABLE

A five-wire patient cable is used to monitor the standard leads, I, II, and III; augmented leads, aVR, aVL, aVF; and one chest lead, usually V₁ (FIG. 2-7).




Figure 2.7 ■ Five-wire patient cable.



Clinical Tip:

Five-wire telemetry units are commonly used to monitor leads I, II, III, aVR, aVL, aVF, and V₁ in critical care settings.

MODIFIED CHEST LEADS

Modified chest leads (MCL) are useful in detecting bundle branch blocks and premature beats. Lead MCL₁ simulates chest lead V₁ and views the ventricular septum (FIG. 2-8). Lead MCL₆ simulates chest lead V₆ and views the lateral wall of the left ventricle (FIG. 2-9).




Figure 2.8 ■ Lead MCL₁ electrode placement.




Figure 2.9 ■ Lead MCL₆ electrode placement.



Clinical Tip:

Write on the rhythm strip which simulated lead was used.

THE RIGHT-SIDED 12-LEAD ECG

Other chest leads that are not part of a standard 12-lead ECG may be used to view specific surfaces of the heart. In a right-sided 12-lead ECG, the limb leads are placed




Figure 2.10 ■ The right-sided chest lead placement.

as usual, but the chest leads are a mirror image of the standard chest lead placement (FIG. 2-10). The ECG machine cannot recognize that the leads have been reversed. It will still print “V₁–V₆” next to the tracing. Be sure to cross this out and write the new lead positions on the ECG paper. TABLE 2-5 shows the placement of the right-sided chest leads.

Table 2.5 ■ THE RIGHT-SIDED CHEST LEAD PLACEMENT

Chest Leads	Position
V _{1R}	Fourth intercostal space to left of sternum
V _{2R}	Fourth intercostal space to right of sternum
V _{3R}	Directly between V _{2R} and V _{4R}
V _{4R}	Fifth intercostal space at right midclavicular line
V _{5R}	Level with V _{4R} at right anterior axillary line
V _{6R}	Level with V _{5R} at right midaxillary line



Clinical Tip:

Patients with an acute inferior MI should have right-sided ECGs to assess for possible right ventricular infarction.

THE 15-LEAD ECG

Areas of the heart that are not well visualized by the six chest leads include the wall of the right ventricle and the posterior wall of the left ventricle. A 15-lead ECG, which includes the standard 12 leads plus leads V_{4R}, V₈, and V₉, increases the chance of detecting an MI in these areas (FIG. 2-11). TABLE 2-6 shows the chest lead placement for the 15-lead ECG.




Figure 2.11 ■ Electrode placement for V_{4R}, V₈, and V₉.

Table 2.6 ■ CHEST LEAD PLACEMENT FOR A 15-LEAD ECG

Chest Leads	Electrode Placement	View of Heart
V _{4R}	Fifth intercostal space in right anterior midclavicular line	Right ventricle
V ₈	Posterior fifth intercostal space in left midscapular line	Posterior wall of left ventricle
V ₉	Directly between V ₈ and spinal column at posterior fifth intercostal space	Posterior wall of left ventricle



Clinical Tip:

Use a 15-lead ECG when the 12-lead tracing is normal but the history still suggests an acute infarction.

RECORDING OF THE ECG

Paper tracings of the ECG patterns from a monitoring lead, inscribed on graph paper, are called rhythm strips. Rhythm strips are valuable because they permit analysis of the ECG in detail. Several different factors can be measured and compared because the graph paper moves past a heated stylus at a standard, constant speed, usually 25 mm/sec (FIG. 2-12).

The vertical lines on the graph paper measure time—one small box equals 0.04 sec and one large box equals 0.20 sec. Therefore these boxes are useful in measuring the duration of various events. Horizontal lines on the graph paper measure voltage, 0.10 mV per each small box. Voltage measurement, however, is only relevant to calibrated tracings, such as a 12-lead or 15-lead ECG produces.




Figure 2.12 ■ ECG graph paper values.

COMPONENTS OF AN ECG TRACING

An ECG tracing specifically reflects electrical activity in the heart. A single cardiac cycle inscribes various deflections on the graph paper. This electrical activity is described as:

Wave: A deflection, either positive or negative, away from the baseline (isoelectric line) of the ECG tracing

Complex: Several waves

Segment: A straight line between waves or complexes

Interval: A segment and a wave



Clinical Tip:

Between waves and cycles, the ECG records a baseline (isoelectric line), which indicates the absence of net electrical activity (Fig. 2-13).

The various patterns described in Figure 2-13 are derived from electrical impulses that originate in the SA node and spread throughout the heart. These electrical components are described as:

P Wave: First wave seen. Small, rounded, and upright (positive); indicates atrial depolarization (and contraction).

PR Interval: Distance between beginning of P wave and beginning of QRS complex. Measures time




Figure 2.13 ■ The electrical pattern of the cardiac cycle shows waves and intervals.

during which a depolarization wave travels from the atria to the ventricles.

QRS Complex: Three deflections following P wave. Indicates ventricular depolarization (and contraction).

Q WAVE: First negative deflection

R WAVE: First positive deflection

S WAVE: First negative deflection after R wave

ST Segment: Distance between end of S wave and beginning of T wave. Measures time between ventricular depolarization and beginning of repolarization.

T Wave: Rounded, upright (positive) wave following QRS complex. Represents ventricular repolarization.

QT Interval: Measured from beginning of QRS complex to end of T wave. Represents total ventricular activity.

U Wave: Small, rounded, upright wave following T wave. Most easily seen with a slow heart rate. Indicates repolarization of Purkinje fibers.

METHODS FOR CALCULATING HEART RATE

Heart rate is calculated as the number of times the heart beats per minute (bpm). On an ECG tracing the bpm is usually calculated as the number of QRS complexes. Included are extra beats such as premature ventricular contractions (PVC), premature atrial contractions



Figure 2.14 ■ Counting large boxes for heart rate. The rate is 60 bpm.

(PAC), and premature junctional contractions (PJC). The rate is measured from the R-R interval, the distance between one R wave and the next. If the atrial rate (the number of P waves) and the ventricular rate (the number of QRS complexes) vary, the analysis may show them as different rates, one atrial and one ventricular.

Method 1: Count Large Boxes

Regular rhythms can be quickly determined by counting the number of large graph boxes between two R waves (FIG. 2-14). That number is divided into 300 to calculate beats per minute. The rates for the first one to six large boxes can be easily memorized.

Remember: 60 sec/min divided by 0.20 sec/large box = 300 large boxes/min. This method usually gives an approximate heart rate but is not as accurate as method 2, explained below.

Method 2: Count Small Boxes

The most accurate way to measure a regular rhythm is to count the number of small boxes between two R waves. That number is divided into 1500 to calculate

Table 2.7 ■ METHODS 1 AND 2 FOR CALCULATING HEART RATE

Number of Large Boxes	Rate/Min	Number of Small Boxes	Rate/Min
1	300	2	750
2	150	3	500
3	100	4	375
4	75	5	300
5	60	6	250
6	50	7	214
7	43	8	186
8	38	9	167
9	33	10	150
10	30	11	136
11	27	12	125
12	25	13	115
13	23	14	107
14	21	15	100
15	20	16	94

bpm. Remember: 60 sec/min divided by 0.04 sec/small box = 1500 small boxes/min.

Examples: If there are three small boxes between two R waves, $1500/3 = 500$ bpm. If there are five small boxes between two R waves, $1500/5 = 300$ bpm.

TABLE 2-7 describes methods 1 and 2 for calculating the heart rate.



Clinical Tip:

The approximate rate per minute is rounded to the next-highest number.

Method 3: Six-second ECG Rhythm Strip

The best method for measuring irregular heart rates with varying R-R intervals is to count the number of R waves in a 6-second (sec) strip (including extra beats such as PVCs, PACs, and PJCs) and multiply by 10 (FIG. 2-15). This gives the average number of beats per minute.




Figure 2.15 ■ Using 6-sec ECG rhythm strip to calculate heart rate: $7 \times 10 = 70$ bpm.



Clinical Tip:

If a rhythm is extremely irregular, it is best to count the number of R-R intervals per 60 sec (1 min).

RHYTHM STRIP ANALYSIS

Arrhythmia interpretation is easiest when you use a consistent analytical approach. Each of the many possible arrhythmias can be described according to the chosen format, and the “rules” for known arrhythmias can be used as a comparison for analyzing an unknown arrhythmia. Although you can use many formats, try to use one approach consistently, and perform all steps in the analysis process to avoid interpretation errors.

Using the format in this text, we begin by analyzing rate, regularity, P waves, PR interval, QRS interval, and QT interval. Next, we check for dropped beats,

pauses, or both. Finally, we look for any grouping of QRS complexes (TABLE 2-8).

INSTRUCTIONS FOR ANALYZING ECG PRACTICE AND TEST STRIPS

Use the following guidelines when analyzing the ECG practice and test strips in this book. They apply to Unit II (Chaps. 3-10) and Unit III (Chaps. 11-14).

- All of the ECG strips were recorded in lead II.
- All ECG strips are 6 sec in length. Notice the 1-sec marks across the top of each strip.
- Rate is determined by the methods you learned in this chapter. Method 2, the small-box method (p 15), is used to measure the rate for regular rhythms. Method 3, the 6-sec method (p 15), is used to determine the rate for irregular rhythms. With a few exceptions, the ventricular rate (R-R interval) is the rate given in the answer section.

Table 2.8 ■ ANALYZING AN ECG RHYTHM

Component	Characteristic
Rate	The bpm is commonly the ventricular rate. If atrial and ventricular rates differ, as in a third-degree block, measure both rates. Normal: 60–100 bpm Slow (bradycardia): <60 bpm Fast (tachycardia): >100 bpm
Regularity	Measure R-R intervals and P-P intervals. Regular: Consistent intervals Regularly irregular: Repeating pattern Irregular: No pattern
P Waves	If present: Same in size, shape, position? Does each QRS have a P wave? Normal: Upright (positive) and uniform Inverted: Negative Notched: P'
PR Interval	None: Rhythm is junctional or ventricular. Constant: Intervals are the same. Variable: Intervals differ.
QRS Interval	Normal: 0.12–0.20 sec and constant Wide: >0.10 sec
QT Interval	None: Absent Beginning of QRS to end of T wave Varies with HR. Normal: Less than half the R-R interval
Dropped beats	Occur in AV blocks. Occur in sinus arrest.
Pause	Compensatory: Complete pause following a PAC, PJC, or PVC Noncompensatory: Incomplete pause following a PAC, PJC, or PVC
QRS Complex grouping	Bigeminy: Repeating pattern of normal complex followed by a premature complex Trigeminy: Repeating pattern of 2 normal complexes followed by a premature complex Quadrigeminy: Repeating pattern of three normal complexes followed by a premature complex Couplets: 2 consecutive premature complexes Triplets: 3 consecutive premature complexes

PAC = premature atrial contraction; PJC = premature junctional contraction; PVC = premature ventricular contraction.

Remember to include any extra beats such as PVCs, PACs, or PJs.

- To determine rhythm, use calipers or an ECG ruler to measure the distance between R-R intervals. If the distance is the same throughout the ECG strip the rhythm is regular. In an irregular rhythm, the R-R intervals are not spaced evenly. Measure each R-R interval on the entire 6-sec ECG strip to make an accurate interpretation of a regular or irregular rhythm.
- Identify and examine the P waves. Notice whether they are normal, absent, or retrograde. Also see whether a pacemaker spike precedes them. Remember, there can be variations of the P wave in each ECG strip.
- Measure the PR intervals. Notice whether each interval is the same or varies in time. If the PR interval is not present, indicate “none.”
- Measure the QRS complexes. Indicate whether they appear notched, as in a bundle branch block. Also notice whether a pacemaker spike precedes them.
- Interpret the tracing in detail. Identify any extra, dropped, or grouped complexes or other important features.
- Check your answers against the answer key at the end of the chapter.

Rhythms and Their Analysis




Sinoatrial Node Arrhythmias

Sinus rhythms all originate in the sinoatrial (SA) node. The ECG features common to all of them are upright P waves, all similar in appearance; normal-duration PR intervals; and normal-duration QRS complexes if no ventricular conduction disturbances are

present. Sinus rhythms described here, in addition to normal sinus rhythm, are sinus bradycardia, sinus tachycardia, sinus arrhythmia, sinus pause (sinus arrest), and SA block. All ECG strips, including the practice strips, were recorded in lead II.

NORMAL SINUS RHYTHM (NSR) ■



Rate: Normal (60–100 bpm)

Rhythm: Regular

P Waves: Normal (upright and uniform)

PR Interval: Normal (0.12–0.20 sec)

QRS: Normal (0.06–0.10 sec)



Clinical Tip:

A normal ECG does not exclude heart disease.




Clinical Tip:

This rhythm is generated by the sinus node and its rate is within normal limits (60–100 bpm).

SINUS BRADYCARDIA ■

- The SA node discharges more slowly than in NSR.



Rate: Slow (<60 bpm)

Rhythm: Regular

P Waves: Normal (upright and uniform)

PR Interval: Normal (0.12–0.20 sec)

QRS: Normal (0.06–0.10 sec)



Clinical Tip:

Sinus bradycardia is normal in athletes and during sleep. In acute MI, it may be protective and beneficial or the slow rate may compromise cardiac output. Certain medications, such as beta blockers, may also cause sinus bradycardia.

SINUS TACHYCARDIA ■

- The SA node discharges more frequently than in NSR.



Rate: Fast (>100 bpm)

Rhythm: Regular

P Waves: Normal (upright and uniform)

PR Interval: Normal (0.12–0.20 sec)

QRS: Normal (0.06–0.10 sec)



Clinical Tip:

Sinus tachycardia may be caused by exercise, anxiety, fever, hypoxemia, hypovolemia, or cardiac failure.

SINUS ARRHYTHMIA ■

- The SA node discharges irregularly.
- The R-R interval is irregular.



Rate: Usually normal (60–100 bpm); frequently increases with inspiration and decreases with expiration; may be < 60 bpm

Rhythm: Irregular; varies with respiration; difference between shortest R-R and longest R-R intervals is > 0.12 sec

P Waves: Normal (upright and uniform)

PR Interval: Normal (0.12–0.20 sec)

QRS: Normal (0.06–0.10 sec)




Clinical Tip:

The pacing rate of the SA node varies with respiration, especially in children and elderly people.

SINUS PAUSE (SINUS ARREST) ■

- The SA node fails to discharge and then resumes.
- Electrical activity resumes either when the SA node resets itself or when a lower latent pacemaker begins to discharge.
- The pause (arrest) time interval is not a multiple of the normal P-P interval.



Rate: Normal to slow; determined by duration and frequency of sinus pause (arrest)

Rhythm: Irregular whenever a pause (arrest) occurs

P Waves: Normal (upright and uniform) except in areas of pause (arrest)

PR Interval: Normal (0.12–0.20 sec)

QRS: Normal (0.06–0.10 sec)




Clinical Tip:

Cardiac output may decrease, causing syncope or dizziness.

SINOATRIAL (SA) BLOCK ■

- The block occurs in some multiple of the P-P interval.
- After the dropped beat, cycles continue on time.



Rate: Normal to slow; determined by duration and frequency of SA block

Rhythm: Irregular whenever an SA block occurs

P Waves: Normal (upright and uniform) except in areas of dropped beats

PR Interval: Normal (0.12–0.20 sec)

QRS: Normal (0.06–0.10 sec)




Clinical Tip:

Cardiac output may decrease, causing syncope or dizziness.

ECG PRACTICE STRIPS ■

For instructions on analyzing these practice strips, please see the guidelines given at the end of Chapter 2.

ECG 3•1




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 3•2



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 3•3




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 3•4




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 3•5



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 3•6




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 3•7




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 3•8




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 3•9



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

Answers to Chapter 3
ECG PRACTICE STRIPS ■

■ **ECG 3•1**

Rate: 35 bpm
Rhythm: Regular
P Waves: Normal
PR Interval: 0.16 sec
QRS: 0.08 sec
Interpretation: Sinus bradycardia

■ **ECG 3•2**

Rate: 63 bpm
Rhythm: Regular
P Waves: Normal
PR Interval: 0.20 sec
QRS: 0.10 sec
Interpretation: Normal sinus rhythm with U wave

■ **ECG 3•3**

Rate: 136 bpm
Rhythm: Regular
P Waves: Normal
PR Interval: 0.16 sec
QRS: 0.10 sec
Interpretation: Sinus tachycardia

■ **ECG 3•4**

Rate: 80 bpm
Rhythm: Irregular
P Waves: Normal
PR Interval: 0.16 sec
QRS: 0.10 sec
Interpretation: Sinus rhythm with
ST segment depression and one
PVC at beat 6

■ ECG 3•5

Rate: 150 bpm
Rhythm: Regular
P Waves: Normal
PR Interval: 0.12 sec
QRS: 0.10 sec
Interpretation: Sinus tachycardia

■ ECG 3•6

Rate: 50 bpm
Rhythm: Irregular
P Waves: Normal
PR Interval: 0.16 sec
QRS: 0.10 sec
Interpretation: Sinus pause (sinus arrest)

■ ECG 3•7

Rate: 70 bpm
Rhythm: Irregular
P Waves: Normal
PR Interval: 0.16 sec
QRS: 0.10 sec
Interpretation: Sinus arrhythmia

■ ECG 3•8

Rate: 70 bpm
Rhythm: Irregular
P Waves: Normal
PR Interval: 0.16 sec
QRS: 0.10 sec
Interpretation: Sinoatrial block

■ ECG 3•9

Rate: 75 bpm
Rhythm: Regular
P Waves: Normal
PR Interval: 0.16 sec
QRS: 0.10 sec
Interpretation: Normal sinus rhythm


Atrial Arrhythmias

Atrial rhythms originate in the atria. Their common ECG features are P waves that differ in appearance from sinus P waves, and normal-duration QRS complexes if no ventricular conduction disturbances are present. Atrial rhythms to be described here are wandering atrial pacemaker (WAP), multifocal atrial tachycardia (MAT), premature atrial contractions (PAC), atrial tachycardia, supraventricular tachycardia (SVT), paroxysmal supraventricular tachycardia (PSVT), atrial flutter (A-flutter), atrial fibrillation (A-fib), and Wolff-Parkinson-White (WPW) Syndrome. All ECG strips, including the practice strips, were recorded in lead II.

cardia (MAT), premature atrial contractions (PAC), atrial tachycardia, supraventricular tachycardia (SVT), paroxysmal supraventricular tachycardia (PSVT), atrial flutter (A-flutter), atrial fibrillation (A-fib), and Wolff-Parkinson-White (WPW) Syndrome. All ECG strips, including the practice strips, were recorded in lead II.

WANDERING ATRIAL PACEMAKER (WAP) ■

- The pacemaker site transfers from the SA node to other latent pacemaker sites in the atria and the AV junction and then moves back to the SA node.



Rate: Normal (60–100 bpm)

Rhythm: Irregular

P Waves: At least three different forms, determined by focus in atria

PR Interval: Variable; determined by focus

QRS: Normal (0.06–0.10 sec)



Clinical Tip:

Wandering atrial pacemaker may occur in normal hearts as a result of fluctuations in vagal tone. It may also be seen in patients with heart disease or COPD.

MULTIFOCAL ATRIAL TACHYCARDIA (MAT) ■

- This form of WAP is associated with a ventricular response of >100 bpm.
- MAT may be confused with atrial fibrillation (A-fib); however, MAT has a visible P wave.



Rate: Fast (>100 bpm)

Rhythm: Irregular

P Wave: At least three different forms, determined by focus in atria

PR Interval: Variable; determined by focus

QRS: Normal (0.06–0.10 sec)




Clinical Tip:

MAT is commonly seen in patients with chronic obstructive pulmonary disease but may also occur in acute MI.

PREMATURE ATRIAL CONTRACTION (PAC) ■

- A single contraction occurs earlier than the next expected sinus contraction.
- After the PAC, sinus rhythm usually resumes.



Rate: Depends on rate of underlying rhythm

Rhythm: Irregular whenever a PAC occurs

P Waves: Present; in the PAC, may have a different shape

PR Interval: Varies in the PAC; otherwise normal (0.12–0.20 sec)

QRS: Normal (0.06–0.10 sec)



Clinical Tip:

In patients with heart disease, frequent PACs may precede PSVT, A-fib, or A-flutter.

ATRIAL TACHYCARDIA ■

- A rapid atrial rate overrides the SA node and becomes the dominant pacemaker.
- Some ST wave and T wave abnormalities may be present.



Rate: 150–250 bpm

Rhythm: Regular


P Waves: Normal (upright and uniform) but differ in shape from sinus P waves

PR Interval: May be short (<0.12 sec) in rapid rates

QRS: Normal (0.06–0.10 sec) but can be aberrant at times

SUPRAVENTRICULAR TACHYCARDIA (SVT) ■

- This arrhythmia has such a fast rate that the P waves may not be seen.



Rate: 150–250 bpm

Rhythm: Regular

P Waves: Frequently buried in preceding T waves and difficult to see

PR Interval: Usually not possible to measure

QRS: Normal (0.06–0.10 sec) but may be wide if abnormally conducted through ventricles



Clinical Tip:

SVT may be related to caffeine intake, nicotine, stress, or anxiety in healthy adults.




Clinical Tip:

Some patients may experience angina, hypotension, light headedness, palpitations, and intense anxiety.

PAROXYSMAL SUPRAVENTRICULAR TACHYCARDIA (PSVT) ■

- PSVT is a rapid rhythm that starts and stops suddenly.
- For accurate interpretation, the beginning or end of the PSVT must be seen.
- PSVT is sometimes called paroxysmal atrial tachycardia (PAT).



Rate: 150–250 bpm

Rhythm: Irregular

P Waves: Frequently buried in preceding T waves and difficult to see

PR Interval: Usually not possible to measure

QRS: Normal (0.06–0.10 sec) but may be wide if abnormally conducted through ventricles




Clinical Tip:

The patient may feel palpitations, dizziness, lightheadedness, or anxiety.

ATRIAL FLUTTER ■

- The AV node conducts impulses to the ventricles at a 2:1, 3:1, 4:1, or greater ratio (rarely 1:1).
- The degree of AV block may be consistent or variable.



Rate: Atrial: 250–350 bpm; ventricular: variable.

Rhythm: Atrial: regular; ventricular: variable

P Waves: Flutter waves have a saw-toothed appearance; some may not be visible, being buried in the QRS

PR Interval: Variable

QRS: Usually normal (0.06–0.10 sec), but may appear widened if flutter waves are buried in QRS



Clinical Tip:

The presence of A-flutter may be the first indication of cardiac disease.




Clinical Tip:

Signs and symptoms depend on the ventricular response rate.

ATRIAL FIBRILLATION ■

- Rapid, erratic electrical discharge comes from multiple atrial ectopic foci.
- No organized atrial depolarization are detectable.



Rate: Atrial: ≥ 350 bpm; ventricular: variable

Rhythm: Irregular

P Waves: No true P waves; chaotic atrial activity

PR Interval: None

QRS: Normal (0.06–0.10 sec)



Clinical Tip:

Atrial fibrillation is often a chronic arrhythmia associated with underlying heart disease.




Clinical Tip:

Signs and symptoms depend on the ventricular response rate.

WOLFF-PARKINSON-WHITE (WPW) SYNDROME ■

- In WPW an accessory conduction pathway is present between the atria and the ventricles. Electrical impulses may be rapidly conducted to the ventricles.
- These rapid impulses create a slurring of the initial portion of the QRS; the slurred effect is called a delta wave.



Rate: Depends on rate of underlying rhythm

Rhythm: Regular unless associated with A-fib

P Waves: Normal (upright and uniform) unless A fib is present

PR Interval: Short (< 0.12 sec)

QRS: Wide (> 0.10 sec); delta wave present




Clinical Tip:

WPW is associated with narrow-complex tachycardias, including A-flutter and A-fib.

ECG PRACTICE STRIPS ■


For instructions on analyzing these practice strips, please see the guidelines given at the end of Chapter 2.

ECG 4•1




Rate: _____ Rhythm: _____
 P Waves: _____ PR Interval: _____ QRS: _____
 Interpretation: _____

ECG 4•2




Rate: _____ Rhythm: _____
 P Waves: _____ PR Interval: _____ QRS: _____
 Interpretation: _____

ECG 4•3



Rate: _____ Rhythm: _____
 P Waves: _____ PR Interval: _____ QRS: _____
 Interpretation: _____

ECG 4•4



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 4•5



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 4•6



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 4•7




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 4•8




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 4•9



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

Answers to Chapter 4

ECG PRACTICE STRIPS ■

■ ECG 4•1

Rate: 214 bpm
Rhythm: Regular
P Waves: Buried in T waves
PR Interval: Not possible to measure
QRS: 0.08 sec
Interpretation: Supraventricular tachycardia

■ ECG 4•2

Rate: 94 bpm
Rhythm: Regular
P Waves: Flutter waves
PR Interval: Not possible to measure
QRS: 0.10 sec
Interpretation: Atrial flutter with 3:1 block (every third flutter wave is buried in the QRS)

■ ECG 4•3

Rate: 214 bpm
Rhythm: Regular
P Waves: Not clearly visible
PR Interval: Not measurable
QRS: 0.08 sec
Interpretation: Supraventricular tachycardia with ST segment depression

■ ECG 4•4

Rate: 140 bpm
Rhythm: Irregular
P Waves: Buried in T waves in beats 1 through 10, normal in beats 11 through 14
PR Interval: None in beats 1 through 10, 0.16 sec in beats 11 through 14
QRS: 0.10 sec
Interpretation: Paroxysmal supraventricular tachycardia (supraventricular tachycardia converting to normal sinus rhythm)

■ ECG 4•5

Rate: 60 bpm
Rhythm: Irregular
P Waves: None
PR Interval: None
QRS: 0.10 sec
Interpretation: Atrial fibrillation

■ ECG 4•6

Rate: 50 bpm
Rhythm: Irregular
P Waves: Normal
PR Interval: 0.20 sec
QRS: 0.08 sec
Interpretation: Sinus bradycardia with two PACs at beats 2 and 4

■ ECG 4•7

Rate: 68 bpm
Rhythm: Regular
P Waves: Flutter waves
PR Interval: Not possible to measure
QRS: 0.20 sec with notched appearance
Interpretation: Atrial flutter with a bundle branch block

■ ECG 4•8

Rate: 180 bpm
Rhythm: Irregular
P Waves: None
PR Interval: None
QRS: 0.08 sec
Interpretation: Atrial fibrillation

■ ECG 4•9


Rate: 120 bpm
Rhythm: Irregular
P Waves: Normal in first three beats
PR Interval: 0.16 sec in first three beats
QRS: 0.10 sec
Interpretation: Paroxysmal supraventricular tachycardia (sinus bradycardia converting to supraventricular tachycardia)

Junctional Arrhythmias

The internodal pathways in the heart merge with the cells of the atrioventricular (AV) junction, which include the AV node. The AV junction is the origin of junctional rhythms. Because it has pacemaker capabilities, it can act as a back-up to the SA node. The ECG features common to all junctional rhythms include P waves that are absent, inverted, buried in the QRS, or

retrograde (after the QRS). A junctional rhythm can also have a PR interval that is absent, short, or retrograde. The rhythms described here are junctional and accelerated junctional rhythms, junctional tachycardia, junctional escape beat, and premature junctional contraction (PJC). All ECG strips, including the practice strips, were recorded in lead II.

JUNCTIONAL RHYTHM ■



Rate: 40–60 bpm

Rhythm: Regular

P Waves: Absent, inverted, buried, or retrograde

PR Interval: None, short, or retrograde


QRS: Normal (0.06–0.10 sec)



Clinical Tip:

The presence of sinus node disease that is causing an inappropriate slowing of the sinus node may exacerbate this rhythm. Young healthy adults, especially those with increased vagal tone during sleep, are often noted to have periods of junctional rhythm that is completely benign, not requiring intervention.

ACCELERATED JUNCTIONAL RHYTHM ■



Rate: 61–100 bpm


Rhythm: Regular

P Waves: Absent, inverted, buried, or retrograde

PR Interval: None, short, or retrograde

QRS: Normal (0.06–0.10 sec)

JUNCTIONAL TACHYCARDIA ■



Rate: 101–180 bpm

Rhythm: Regular

P Waves: Absent, inverted, buried, or retrograde

PR Interval: None, short, or retrograde

QRS: Normal (0.06–0.10 sec)



Clinical Tip:

Signs and symptoms of decreased cardiac output may be seen in response to the rapid rate.

JUNCTIONAL ESCAPE BEAT ■

- An escape complex comes later than the next expected sinus complex.



Rate: Depends on rate of underlying rhythm

Rhythm: Irregular whenever an escape beat occurs

P Waves: None, inverted, buried, or retrograde in the escape beat

PR Interval: None, short, or retrograde

QRS: Normal (0.06–0.10 sec)

PREMATURE JUNCTIONAL CONTRACTION (PJC) ■

- Enhanced automaticity in the AV junction produces PJCs.



Rate: Depends on rate of underlying rhythm

Rhythm: Irregular whenever a PJC occurs

P Waves: Absent, inverted, buried, or retrograde in the PJC

PR Interval: None, short, or retrograde

QRS: Normal (0.06–0.10 sec)




Clinical Tip:

Before deciding that isolated PJCs are insignificant, consider the cause.

ECG PRACTICE STRIPS ■

For instructions on analyzing these practice strips, please see the guidelines given at the end of Chapter 2.

ECG 5•1




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 5•2




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 5•3




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 5•4



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 5•5




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 5•6




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 5•7




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 5•8




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 5•9



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

Answers to Chapter 5

ECG PRACTICE STRIPS ■

■ **ECG 5•1**

Rate: 75 bpm
 Rhythm: Regular
 P Waves: Inverted or absent
 PR Interval: 0.16 sec with inverted P waves
 QRS: 0.10 sec
 Interpretation: Accelerated junctional rhythm

■ **ECG 5•2**

Rate: 130 bpm
 Rhythm: Irregular
 P Waves: Retrograde
 PR Interval: None
 QRS: 0.10 sec
 Interpretation: Junctional tachycardia with two uni-
 form PVCs at beats 6 and 11

■ **ECG 5•3**

Rate: 70 bpm
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: 0.10 sec
 Interpretation: Junctional rhythm
 with bigeminal PJC's at beats 2, 4,
 and 6

■ **ECG 5•4**

Rate: 38 bpm
 Rhythm: Regular
 P Waves: Inverted
 PR Interval: 0.16 sec
 QRS: 0.10 sec
 Interpretation: Junctional rhythm

■ ECG 5•5

Rate: 150 bpm
 Rhythm: Regular
 P Waves: None
 PR Interval: None
 QRS: 0.10 sec
 Interpretation: Junctional tachycardia

■ ECG 5•6

Rate: 47 bpm
 Rhythm: Regular
 P Waves: Inverted
 PR Interval: 0.10 sec
 QRS: 0.10 sec
 Interpretation: Junctional rhythm with
 ST segment elevation

■ ECG 5•7

Rate: 160 bpm
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: 0.10 sec
 Interpretation: Accelerated junctional rhythm with
 bigeminal uniform PVCs (R on T phenomenon*)

■ ECG 5•8

Rate: 150 bpm
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: 0.10 sec
 Interpretation: Junctional tachycardia with
 uniform PVCs (R on T phenomenon*) at
 beats 2, 5, and 7

■ ECG 5•9

Rate: 110 bpm
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: 0.10 sec
 Interpretation: Junctional tachycardia with
 ST segment depression and uniform PVCs
 at beats 5 and 9

*R on T phenomenon is discussed in Chapter 6.

Ventricular Arrhythmias

All arrhythmias that originate in the ventricles depolarize the ventricles abnormally and at a slower speed. For this reason, the ECG feature common to all ventricular rhythms is a QRS complex wider than 0.10 sec in duration. The P waves are either absent or, if visible, have no consistent relationship to the QRS interval (the length of the QRS complex). The ventricular rhythms

discussed are idioventricular and accelerated idioventricular rhythm, premature ventricular contraction (PVC), monomorphic ventricular tachycardia (VT), polymorphic VT, torsade de pointes, ventricular fibrillation (VF), pulseless electrical activity (PEA), and asystole. All ECG strips, including the practice strips, were recorded in lead II.

IDIOVENTRICULAR RHYTHM ■



Rate: 20–40 bpm

Rhythm: Regular

P Waves: None

PR Interval: None


QRS: Wide (>0.10 sec),
bizarre appearance



Clinical Tip:

Diminished cardiac output is expected because of the slow heart rate. An idioventricular rhythm may be called an agonal rhythm when the heart rate drops below 20 bpm. An agonal rhythm is generally a terminal event and is usually the last rhythm before asystole.

ACCELERATED IDIOVENTRICULAR RHYTHM ■



Rate: 41–100 bpm

Rhythm: Regular

P Waves: None

PR Interval: None

QRS: Wide (>0.10 sec),
bizarre appearance




Clinical Tip:

Idioventricular rhythms appear when supraventricular pacing sites are suppressed or absent.

PREMATURE VENTRICULAR CONTRACTION (PVC) ■

- PVCs result from an irritable ventricular focus.
- PVCs may be uniform (same form) or multiform (different forms).
- Usually a PVC is followed by a full compensatory pause because the sinus node timing is not interrupted. Normally the sinus rate produces the next sinus impulse on time. In contrast, a PVC may be followed by a noncompensatory pause if the PVC enters the sinus node and resets its timing; this enables the following sinus P wave to appear earlier than expected.



Rate: Depends on rate of underlying rhythm

Rhythm: Irregular whenever a PVC occurs

P Waves: None associated with
the PVC

PR Interval: None associated with
the PVC


QRS: Wide (>0.10 sec),
bizarre appearance




Clinical Tip:

Patients may sense the occurrence of PVCs as skipped beats. Because the ventricles are only partially filled, the PVC frequently does not generate a pulse.

PREMATURE VENTRICULAR CONTRACTION: UNIFORM ■



PREMATURE VENTRICULAR CONTRACTION: MULTIFORM ■




PREMATURE VENTRICULAR CONTRACTION: VENTRICULAR BIGEMINY ■

- In ventricular bigeminy, the PVC occurs with every other beat.




PREMATURE VENTRICULAR CONTRACTION: VENTRICULAR TRIGEMINY ■

- In ventricular trigeminy, the PVC occurs with every third beat.




PREMATURE VENTRICULAR CONTRACTION: VENTRICULAR QUADRIGEMINY ■

- In ventricular quadrigeminy, the PVC occurs with every fourth beat.




PREMATURE VENTRICULAR CONTRACTION: COUPLETS ■

- In a rhythm with PVC couplets, the PVCs occur in pairs.



PREMATURE VENTRICULAR CONTRACTION: R ON T PHENOMENON ■

- The PVCs occur so early that they fall on the T wave of the preceding beat.
- These PVCs occur during the refractory period of the ventricles, a vulnerable period because the cardiac cells have not fully repolarized.



Rate: Depends on rate of underlying rhythm

Rhythm: Irregular whenever a PVC occurs

P Waves: None associated with the PVC

PR Interval: None associated with the PVC

QRS: Wide (>0.10 sec), bizarre appearance




Clinical Tip:

In an acute ischemic situation, R on T phenomenon may be especially dangerous because the ventricles may be more vulnerable to VT or VF.

PREMATURE CONTRACTION: INTERPOLATED PVC ■

- The PVC occurs between two regular complexes; it may appear sandwiched in between two normal beats.
- An interpolated PVC does not interfere with the normal cardiac cycle.



Rate: Depends on rate of underlying rhythm

Rhythm: Irregular whenever a PVC occurs


P Waves: None associated with the PVC

PR Interval: None associated with the PVC

QRS: Wide (>0.10 sec), bizarre appearance

VENTRICULAR TACHYCARDIA (VT): MONOMORPHIC ■

- In monomorphic VT, QRS complexes have the same shape and amplitude.



Rate: 100–250 bpm

Rhythm: Regular

P Waves: None or not associated with the QRS

PR Interval: None

QRS: Wide (>0.10 sec), bizarre appearance



Clinical Tip:

It is important to confirm the presence or absence of pulses because monomorphic VT may be perfusing or nonperfusing.



Clinical Tip:

Monomorphic VT will probably deteriorate into VF or unstable VT if sustained and not treated.

VENTRICULAR TACHYCARDIA (VT): POLYMORPHIC ■

- In polymorphic VT, QRS complexes vary in shape and amplitude.
- The QT interval is normal or long.



Rate: 100–250 bpm

Rhythm: Regular or irregular

P Waves: None or not associated with the QRS

PR Interval: None

QRS: Wide (>0.10 sec), bizarre appearance



Clinical Tip:

It is important to confirm the presence or absence of pulses because polymorphic VT may be perfusing or nonperfusing.




Clinical Tip:

Consider electrolyte abnormalities as a possible cause.

TORSADE DE POINTES ■

- The QRS reverses polarity and the strip shows a spindle effect.
- This rhythm is an unusual variant of polymorphic VT with long QT intervals.
- In French the term means “twisting of points.”



Rate: 200–250 bpm

Rhythm: Irregular

P Waves: None

PR Interval: None

QRS: Wide (>0.10 sec), bizarre appearance



Clinical Tip:

Torsade de pointes may deteriorate to VF or asystole.




Clinical Tip:

Frequent causes are drugs that prolong QT interval, and electrolyte abnormalities such as hypomagnesemia.

VENTRICULAR FIBRILLATION (VF) ■

- Chaotic electrical activity occurs with no ventricular depolarization or contraction.

The amplitude and frequency of the fibrillatory activity can be used to define the type of fibrillation as coarse, medium, or fine. Small baseline undulations are considered fine; large ones are coarse.



Rate: Indeterminate

Rhythm: Chaotic

P Waves: None

PR Interval: None

QRS: None



Clinical Tip:

There is no pulse or cardiac output. Rapid intervention is critical. The longer the delay, the less the chance of conversion.

PULSELESS ELECTRICAL ACTIVITY (PEA) ■

- The monitor shows an identifiable electrical rhythm, but no pulse is detected.
- The rhythm may be sinus, atrial, junctional, or ventricular.
- PEA is also called electromechanical dissociation (EMD).



Rate: Reflects underlying rhythm

Rhythm: Reflects underlying rhythm

P Waves: Reflect underlying rhythm

PR Interval: Reflects underlying rhythm

QRS: Reflects underlying rhythm




Clinical Tip:

Potential causes of PEA are trauma, tension pneumothorax, thrombosis (pulmonary or coronary), cardiac tamponade, toxins, hypo- or hyperkalemia, hypovolemia, hypoxia, hypoglycemia, hypothermia, and hydrogen ion (acidosis).

ASYSTOLE ■

- Electrical activity in the ventricles is completely absent.



Rate: None

Rhythm: None

P Waves: None

PR Interval: None

QRS: None



Clinical Tip:

Rule out other causes such as loose leads, no power, or signal gain too low.




Clinical Tip:

Seek to identify the underlying cause as in PEA. Also, search to identify underlying VF.

ECG PRACTICE STRIPS ■

For instructions on analyzing these practice strips, please see the guidelines given at the end of Chapter 2.

ECG 6•1




Rate: _____ **Rhythm:** _____

P Waves: _____ **PR Interval:** _____ **QRS:** _____

Interpretation: _____

ECG 6•2




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 6•3




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 6•4



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____


Interpretation: _____

ECG 6•5




Rate: _____ Rhythm: _____
P Waves: _____ PR Interval: _____ QRS: _____
Interpretation: _____

ECG 6•6




Rate: _____ Rhythm: _____
P Waves: _____ PR Interval: _____ QRS: _____
Interpretation: _____

ECG 6•7



Rate: _____ Rhythm: _____
P Waves: _____ PR Interval: _____ QRS: _____
Interpretation: _____

ECG 6•8



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 6•9



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

Answers to Chapter 6

ECG PRACTICE STRIPS ■

■ **ECG 6•1**

Rate: 167 bpm
 Rhythm: Regular
 P Waves: None
 PR Interval: None
 QRS: Wide—greater than 0.10 sec
 Interpretation: Ventricular tachycardia—
 monomorphic

■ **ECG 6•2**

Rate: 80 bpm (counting PVCs), 41 bpm in under-
 lying rate
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.10 sec
 Interpretation: Sinus bradycardia with interpolated
 multiform PVCs

■ **ECG 6•3**

Rate: 214 bpm
 Rhythm: Regular
 P Waves: None
 PR Interval: None
 QRS: Wide—greater than 0.10 sec
 Interpretation: Ventricular tachycardia—
 monomorphic

■ **ECG 6•4**

Rate: 43 bpm
 Rhythm: Regular
 P Waves: None
 PR Interval: None
 QRS: Wide—greater than 0.10 sec
 Interpretation: Accelerated idioventricular
 rhythm

■ ECG 6•5

Rate: Indeterminate
 Rhythm: Chaotic
 P Waves: None
 PR Interval: None
 QRS: None
 Interpretation: Ventricular fibrillation

■ ECG 6•6

Rate: 170 bpm (counting PVCs), 150 bpm in underlying rate
 Rhythm: Irregular
 P Waves: Buried in T wave
 PR Interval: Not measurable
 QRS: 0.10 sec
 Interpretation: Supraventricular tachycardia with two uniform PVCs at beats 4 and 15

■ ECG 6•7

Rate: Indeterminate
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: Wide—greater than 0.10 sec
 Interpretation: Torsade de pointes

■ ECG 6•8

Rate: 40 bpm
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: Wide—greater than 0.10 sec
 Interpretation: Idioventricular rhythm with one PVC at beat 2

■ ECG 6•9

Rate: None
 Rhythm: None
 P Waves: None
 PR Interval: None
 QRS: None
 Interpretation: Asystole

Atrioventricular and Bundle Branch Blocks

Atrioventricular (AV) blocks reflect delay or interruption of impulses through the AV junction due to disease in this region. They are traditionally divided into three categories: first, second, and third degree. This pathological block, caused by such conditions as ischemia, necrosis, degenerative diseases of the conduction system, and drug toxicity, is different from the physiological AV block that occurs in atrial flutter and fibrillation.

Another disorder, involving bundle branch conduction through the ventricles, is bundle branch block (BBB). The problem does not occur until the signal is conducted through the ventricles, although the block originates above them.

All ECG strips, including the practice strips, were recorded in lead II.

FIRST-DEGREE AV BLOCK ■



Rate: Depends on rate of underlying rhythm

Rhythm: Regular

P Waves: Normal (upright and uniform)

PR Interval: Prolonged (>0.20 sec)

QRS: Normal (0.06–0.10 sec)



Clinical Tip:

Usually a first-degree AV block is benign, but if associated with an acute MI it may lead to further AV defects.




Clinical Tip:

Often AV block is caused by medications that prolong AV conduction; these include digoxin, calcium channel blockers, and beta blockers.

SECOND-DEGREE AV BLOCK: TYPE I (MOBITZ I OR WENCKEBACH) ■

- PR intervals become progressively longer until one P wave is totally blocked and produces no QRS complex. After a pause, during which the AV node recovers, this cycle is repeated.



Rate: Depends on rate of underlying rhythm

Rhythm: Atrial: regular; ventricular: irregular

P Waves: Normal (upright and uniform), more P waves than QRS

PR Interval: Progressively longer until one P wave is blocked and a QRS is dropped

QRS: Normal (0.06–0.10 sec)



Clinical Tip:

This rhythm may be caused by medication such as beta blockers, digoxin, and calcium channel blockers. Ischemia involving the right coronary artery is another cause.

SECOND-DEGREE AV BLOCK: TYPE II (MOBITZ II) ■

- Conduction ratio (P waves to QRS complexes) is commonly 2:1, 3:1, 4:1, or variable.
- QRS complexes are usually wide because this block usually involves both bundle branches.



Rate: Atrial: usually 60–100 bpm; ventricular: slower than atrial rate

Rhythm: Atrial regular and ventricular may be regular or irregular

P Waves: Normal (upright and uniform); more P waves than QRSs

PR Interval: Normal or prolonged but constant

QRS: May be normal, but usually wide (>0.10 sec) if the bundle branches are involved



Clinical Tip:

Resulting bradycardia can compromise cardiac output and lead to complete AV block. This rhythm often occurs with cardiac ischemia or an MI.

THIRD-DEGREE AV BLOCK ■

- Conduction between the atria and the ventricles is totally absent because of complete electrical block at or below the AV node. This is known as AV dissociation.
- “Complete heart block” is another name for this rhythm.



Rate: Atrial: 60–100 bpm; ventricular: 40–60 bpm if escape focus is junctional, <40 bpm if escape focus is ventricular

Rhythm: Usually regular, but atria and ventricles act independently

P Waves: Normal (upright and uniform); may be superimposed on QRS complexes or T waves

PR Interval: Varies greatly

QRS: Normal if ventricles are activated by junctional escape focus; wide if escape focus is ventricular




Clinical Tip:

Third-degree AV block may be associated with ischemia involving the left coronary arteries.

BUNDLE BRANCH BLOCK (BBB) ■

- Either the left or the right ventricle may depolarize late, creating a “wide” or “notched” QRS complex.



Rate: Depends on rate of underlying rhythm

Rhythm: Regular

P Waves: Normal (upright and uniform)

PR Interval: Normal (0.12–0.20 sec)

QRS: Wide (>0.10 sec) with or without a notched appearance




Clinical Tip:

Bundle branch block commonly occurs in coronary artery disease.

ECG PRACTICE STRIPS ■

For instructions on analyzing these practice strips, please see the guidelines given at the end of Chapter 2.

ECG 7•1




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 7•2




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 7•3



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 7•4




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 7•5



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 7•6



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 7•7



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 7•8




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 7•9



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

*Answers to Chapter 7***ECG PRACTICE STRIPS** ■■ **ECG 7•1**

Rate: Atrial 125 bpm, ventricular 44 bpm
 Rhythm: Atrial regular, ventricular regular
 P Waves: Normal
 PR Interval: 0.12 sec and constant
 QRS: 0.10 sec
 Interpretation: Second-degree AV block
 Type II, 3:1 conduction

■ **ECG 7•2**

Rate: 80 bpm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.32 sec
 QRS: 0.10 sec
 Interpretation: First-degree AV block with a PJC at
 beat 6

■ **ECG 7•3**

Rate: 60 bpm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: Progressive lengthening
 QRS: Wide—greater than 0.10 sec
 Interpretation: Second-degree AV
 block Type I (Wenckebach) with
 wide QRS

■ **ECG 7•4**

Rate: 70 bpm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: Progressive lengthening
 QRS: 0.10 sec
 Interpretation: Second-degree AV block
 Type I (Wenckebach)

■ **ECG 7•5**

Rate: 40 bpm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.16 sec and constant
 QRS: 0.08 sec
 Interpretation: Second-degree AV block Type II with
 inverted T waves

■ **ECG 7•6**

Rate: 56 bpm
 Rhythm: Regular
 P Waves: Normal but not associated with QRS
 PR Interval: Variable
 QRS: Wide—greater than 0.10 sec with notched
 appearance
 Interpretation: Third-degree AV block with a bundle
 branch block

■ **ECG 7•7**

Rate: 65 bpm
 Rhythm: Regular
 P Waves: Normal
 PR Interval: 0.20 sec
 QRS: 0.16 sec with notched appearance
 Interpretation: Normal sinus rhythm with a bundle
 branch block

■ **ECG 7•8**

Rate: 90 bpm (counting PVCs), 94 in underlying rate
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.28 sec
 QRS: 0.10 sec
 Interpretation: Normal sinus rhythm with first-
 degree AV block with multiform PVCs at beats
 4, 7, and 9

■ **ECG 7•9**

Rate: 56 bpm
 Rhythm: Regular
 P Waves: Normal but not associated with QRS
 PR Interval: Variable
 QRS: 0.10 sec
 Interpretation: Third-degree AV block

Artificial Cardiac Pacemakers

As you found in Chapter 1, myocardial fibers possess highly specialized electrical properties. The components of automaticity, excitability, and conductivity allow a certain number of specialized cardiac cells, called pacemaker cells, to go into action. These cells actually do pace the heart. Each specialized area of the heart has its own intrinsic rate:

- Sinoatrial node: The dominant pacemaker of the heart. Intrinsic rate is 60–100 bpm.
- Atrioventricular node: Intrinsic rate is 40–60 bpm.
- Purkinje system: The bundle branches terminate with this network of fibers. They spread electrical impulses rapidly throughout the ventricular walls. Intrinsic rate is 20–40 bpm.

People with strong, healthy hearts, such as athletes, can sustain an adequate blood flow in their cardiovascular system with heart rates as low as 50 to 60 bpm. But in a person with an unhealthy heart, the natural pacemakers can discharge too slowly or stop altogether. A slow heart rate may be caused by factors such as coronary artery disease or by the side effects of certain medications.

Heart failure can result from an insufficient amount of oxygenated blood being pumped throughout the cardiovascular system. The person begins to suffer from hypotension, bradycardia, diaphoresis, decreased level of consciousness, and possibly an AV block. In this case an artificial pacemaker can save a person's life.

ARTIFICIAL PACEMAKER

An artificial cardiac pacemaker is a device used to electronically stimulate the heart in place of the heart's own natural pacemakers and conduction system. It is composed of one or more electrodes implanted in the heart and connected to a power source that generates regular, timed stimuli. It may be preset to stimulate the heart's activity continuously or intermittently.

Some of the conditions for insertion of a pacemaker are continuous or intermittent third-degree AV block, second-degree AV block Type II, chronic A-fib with bradycardic ventricular response, and sick sinus syndrome (a condition marked by sinus block, severe sinus bradycardia, or alternating periods of bradycardia and tachycardia).

Several types of pacemakers exist, but the most common types used for the conditions just described are the ventricular and dual-chambered pacemakers. Ventricular pacemakers stimulate only the ventricle, while dual-chambered types stimulate both the atrium and ventricle, if needed. The following bullet points explain terms you will find associated with artificial cardiac pacemakers.

Temporary Pacemaker

- A temporary pacemaker is commonly used in an emergency. It paces the heart through epicardial, transvenous, or transcutaneous routes. The pulse generator is located externally. Indications may include symptomatic bradycardia, AV heart block (second-degree Type II or third degree), change in mental status, or pulmonary edema.

Permanent Pacemaker

- A permanent pacemaker may be indicated if there is a chronic or intermittent AV block, sinus arrest, or sick sinus syndrome. It is surgically implanted, usually under local anesthesia. Its circuitry is sealed in an airtight case and the pacemaker is implanted in the body. It uses sensing and pacing device leads.

Single-Chamber Pacemaker

- One lead is placed in the heart and paces a single chamber (either atrium or ventricle).

Table 8.1 ■ PACEMAKER CODES

Chamber Paced	Chamber Sensed	Response to Sensing	Programmable Functions	Response to Tachycardia
A = Atrium V = Ventricle D = Dual (atrium and ventricle) O = None	A = Atrium V = Ventricle D = Dual (atrium and ventricle) O = None	T = Triggers pacing I = Inhibits pacing D = (triggers and inhibits) O = None	P = Basic programs (rate and output) M = Multiple programs C = Communication (i.e., telemetry) R = Rate response O = None	P = Pacing S = Shock D = Dual (pace and shock) O = None

Dual-Chamber Pacemaker

- One lead is placed in the right atrium and the other in the right ventricle. The atrial electrode generates a spike that should be followed by a P wave, and the ventricular electrode generates a spike followed by a wide QRS complex.

PACEMAKER MODES

- Fixed rate (asynchronous): Discharges at a preset rate (usually 70–80 bpm) regardless of the patient's own electrical activity.
- Demand (synchronous): Discharges only when the patient's heart rate drops below the pacemaker's preset (base) rate.



Clinical Tip:

Patients with pacemakers may receive defibrillation, but avoid placing the defibrillator paddles or pads less than 5 inches from the pacemaker battery pack.

UNDERSTANDING PACEMAKER CODES

An international code was developed in 1974 to identify the preprogrammed pacing, sensing, and response functions of a permanent pacemaker. The code initially consisted of three letters, but in 1980 the system was modified with the addition of two more letters (designating two more functions). The first three letters are the most commonly known because they are used for symptomatic bradycardia.

Understanding the codes used in a permanent pacemaker help to ensure that the pacemaker is set

properly. A simple classification of the five-code system is described in TABLE 8-1.

ARTIFICIAL PACEMAKER RHYTHM

- An artificial pacemaker may vary in its rate, rhythm, P waves, P-R interval, and QRS complex.

Rate: Varies according to preset pacemaker rate.

Rhythm: Regular for asynchronous pacemaker; irregular for demand pacemaker unless 100% paced with no intrinsic beats.

P waves: None produced by ventricular pacemaker. Sinus P waves may be seen but are unrelated to QRS. Atrial or dual-chamber pacemaker should have P waves following each atrial spike.

P-R interval: None for ventricular pacer. Atrial or dual-chamber pacemaker produces constant PR intervals.


QRS: Wide (greater than 0.10 sec) following each ventricular spike in a pacemaker rhythm. The patient's own electrical activity may generate a QRS complex that looks different from the paced QRSs. If atrially paced only, the QRS complex may be within normal limits.




Clinical Tip:

Once an impulse is generated by the pacemaker it appears as a spike, either above or below the baseline (isoelectric line), on the ECG. The spike indicates that the pacemaker has fired.


SINGLE-CHAMBER PACEMAKER RHYTHM—ATRIAL



SINGLE-CHAMBER PACEMAKER RHYTHM—VENTRICULAR



DUAL-CHAMBER PACEMAKER RHYTHM—ATRIAL AND VENTRICULAR



PACEMAKER MALFUNCTIONS

At times a pacemaker may have a malfunction. Common problems are listed in TABLE 8-2.




Clinical Tip:

A pacemaker is said to be in capture when a spike produces an ECG wave or complex.

Table 8.2 ■ **PACEMAKER MALFUNCTIONS**

Malfunction	Reason
Failure to pace	Pacemaker spikes are absent. The cause may be a dead battery, a disruption in the connecting wires, or improper programming.
Failure to capture	Pacemaker spikes are present, but no P wave or QRS follows the spike. Turning up the pacemaker's voltage often corrects this problem. Lead wires should also be checked—a dislodged or broken lead wire may deliver some, but not all, of the needed energy.
Failure to sense	The pacemaker fires because it fails to detect the heart's intrinsic beats, resulting in abnormal complexes. The cause may be a dead battery, decrease of P wave or QRS voltage, or damage to a pacing lead wire. One serious potential consequence may be an R on T phenomenon.
Oversensing	The pacemaker may be too sensitive and misinterpret muscle movement or other events in the cardiac cycle as depolarization. This error resets the pacemaker inappropriately, increasing the amount of time before the next discharge.

FAILURE TO CAPTURE



FAILURE TO SENSE




OVERSENSING



ECG PRACTICE STRIPS ■

For instructions on analyzing these practice strips, please see the guidelines given at the end of Chapter 2.

ECG 8•1




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 8•2



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 8•3




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 8•4




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 8•5




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 8•6




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 8•7




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 8•8



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 8•9



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

Answers to Chapter 8

ECG PRACTICE STRIPS ■

■ ECG 8•1

Rate: 50 bpm
 Rhythm: Regular
 P Waves: Normal following pacemaker spike
 PR Interval: 0.20 sec
 QRS: 0.10 sec
 Interpretation: Pacemaker—atrial, with ST segment depression and inverted T waves

■ ECG 8•2

Rate: 80 bpm
 Rhythm: Irregular
 P Waves: None following pacemaker spike
 PR Interval: None
 QRS: Wide—greater than 0.10 sec following pacemaker spike
 Interpretation: Pacemaker—atrial and ventricular, with one PVC at beat 5. Notice that there is no P wave generated with the atrial spike. This would be a failure to capture with the atrial spike.

■ ECG 8•3

Rate: 80 bpm
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: Wide—greater than 0.10 sec following pacemaker spike
 Interpretation: Pacemaker—ventricular, with PVCs at beats 4 and 6

■ ECG 8•4

Rate: 40 bpm
 Rhythm: Regular
 P Waves: Normal at beats 2 through 4
 PR Interval: 0.20 sec at beats 2 through 4
 QRS: Wide—greater than 0.10 sec at beat 1; 0.10 sec at beats 2 through 4
 Interpretation: Pacemaker—ventricular evolving to sinus complexes, with no ventricular pacing, at beats 3 and 4

■ ECG 8•5

Rate: 65 bpm
 Rhythm: Regular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.10 sec
 Interpretation: Pacemaker—atrial, with nonpaced P wave at beat 3

■ ECG 8•6

Rate: 100 bpm
 Rhythm: Regular
 P Waves: Normal with low voltage following pacemaker spike
 PR Interval: 0.16 sec
 QRS: Wide—greater than 0.10 sec with notched appearance following pacemaker spike
 Interpretation: Pacemaker—atrial and ventricular

■ ECG 8•7

Rate: 125 bpm
 Rhythm: Regular
 P Waves: None
 PR Interval: None
 QRS: Wide—greater than 0.10 sec following pacemaker spike
 Interpretation: Pacemaker—ventricular

■ ECG 8•8

Rate: 60 bpm
 Rhythm: Regular
 P Waves: None following pacemaker spike
 PR Interval: None
 QRS: Wide—greater than 0.10 sec following pacemaker spike
 Interpretation: Pacemaker—atrial and ventricular, with no atrial conduction. The atrial spike shows failure to capture.

■ ECG 8•9

Rate: 80 bpm
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: Wide—greater than 0.10 sec following pacemaker spike
 Interpretation: Pacemaker—ventricular, with one junctional complex with ST segment depression and inverted T wave at beat 6


Artifact

In an ECG tracing, waves, segments, and complexes are produced by the heart's electrical activity. However, deflections are occasionally produced by other influences, known as ECG artifact. Some common causes of artifact are loose electrodes; patient movement and

muscle tremors; ECG calibration marks; 60-cycle electrical interference; and malfunction of the ECG machine, patient cable, or lead wires. Artifact can complicate the interpretation of an ECG tracing, so it is important to recognize and eliminate it whenever possible.

LOOSE ELECTRICAL CONNECTION ■

- Bizarre, irregular deflections in the baseline (isoelectric line) of the ECG tracing may be caused by poor electrical contact, a loose electrode, or a broken wire. Make sure all the electrical and patient cable connections are secure.




VARIATION WITH RESPIRATION ■

- Patient movement may occur with respiration. The resulting undulations move the entire tracing up and down, reflecting the rhythm of the patient's breathing.



60-CYCLE INTERFERENCE ■

- Improperly grounded electrical equipment or other electrical interference can cause a phenomenon called 60-cycle interference. If this occurs, check to make sure all electrical equipment is properly grounded and that the patient cable electrical connections are clean.



MUSCLE ARTIFACT ■

- Shivering, tense muscles, seizures, patient movement, and disorders such as Parkinson's disease may cause muscle tremor artifact.




Clinical Tip:

Never confuse muscle artifact with A-fib if the rhythm is regular.

ECG PRACTICE STRIPS ■

For instructions on analyzing these practice strips, please see the guidelines given at the end of Chapter 2.

ECG 9•1




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 9•2




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 9•3




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 9•4



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 9•5




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 9•6



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 9•7




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 9•8




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 9•9



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

Answers to Chapter 9

ECG PRACTICE STRIPS ■

■ ECG 9•1

Rate: 60 bpm (counting PVCs), 66 bpm in underlying rate
 Rhythm: Irregular
 P Waves: Present, but hard to see because of artifact
 PR Interval: Not possible to measure
 QRS: 0.10 sec
 Interpretation: Normal sinus rhythm with uniform PVCs and muscle artifact

■ ECG 9•2

Rate: 80 bpm
 Rhythm: Irregular
 P Waves: Normal in beats 5 through 8
 PR Interval: 0.16 sec in beats 5 through 8
 QRS: 0.10 sec in beats 5 through 8
 Interpretation: Normal sinus rhythm beginning with muscle artifact and adjusting to a normal baseline in beats 5 through 8

■ ECG 9•3

Rate: 80 bpm
 Rhythm: Irregular
 P Waves: Normal but with artifact
 PR Interval: 0.16 sec
 QRS: 0.10 sec
 Interpretation: Normal sinus rhythm with two PACs at beats 4 and 7 and 60-cycle interference

■ ECG 9•4

Rate: 80 bpm
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: 0.10 sec
 Interpretation: Atrial fibrillation with 60-cycle interference

■ **ECG 9•5**

Rate: 38 bpm
Rhythm: Regular
P Waves: Present, but hard to see because of artifact
PR Interval: Not possible to measure
QRS: 0.10 sec
Interpretation: Sinus bradycardia with muscle artifact

■ **ECG 9•6**

Rate: None
Rhythm: None
P Waves: None
PR Interval: None
QRS: None
Interpretation: Loose electrodes

■ **ECG 9•7**

Rate: Not possible to measure
Rhythm: Irregular
P Waves: None
PR Interval: None
QRS: Wide—greater than 0.10 sec
Interpretation: Torsade de pointes with 60-cycle interference

■ **ECG 9•8**

Rate: Not possible to measure
Rhythm: Irregular
P Waves: None
PR Interval: None
QRS: Wide—greater than 0.10 sec
Interpretation: Torsade de pointes with muscle artifact

■ **ECG 9•9**

Rate: 90 bpm
Rhythm: Irregular
P Waves: None
PR Interval: None
QRS: 0.10 sec
Interpretation: Atrial fibrillation with muscle artifact

The 12-Lead ECG and Acute Myocardial Infarction

A standard 12-lead ECG provides views of the heart from 12 different angles. This diagnostic test helps to identify pathological conditions, especially bundle branch blocks and T wave changes associated with ischemia, injury, and infarction. The 12-lead ECG also uses ST segment analysis to pinpoint the specific location of a myocardial infarction (MI).

Multiple views give a more comprehensive picture of the heart's electrical activity than a rhythm strip and can be used to assess left ventricular function. Patients with other conditions, such as electrolyte imbalances or adverse conditions caused by certain medications, may also benefit from a 12-lead ECG.

The 12-lead ECG is the type most commonly used in clinical settings. The following list highlights some of its important aspects:

- The 12-lead ECG consists of the six limb leads—I, II, III, aVR, aVL, and aVF—and the six chest leads—V₁, V₂, V₃, V₄, V₅, and V₆.
- The limb leads record electrical activity in the heart's frontal plane. This view shows the middle of the heart from top to bottom. Electrical activity is recorded from the anterior-to-posterior axis.
- The chest leads record electrical activity in the heart's horizontal plane. This transverse view shows the middle of the heart from right to left, dividing it into upper and lower portions. Electrical activity is recorded from either a superior or an inferior approach.
- Measurements are central to 12-lead ECG analysis. The height and depth of waves can offer important diagnostic information in certain conditions, including MI and ventricular hypertrophy.
- The direction of ventricular depolarization is an important factor in determining the axis of the heart.
- In an MI, multiple leads are necessary to recognize its presence and determine its location. If large areas of the heart are affected, the patient can develop cardiogenic shock and fatal arrhythmias.

- ECG signs of an MI are best seen in the reflecting leads—those facing the affected surface of the heart. Reciprocal leads are in the same plane but opposite the area of infarction; they show a “mirror image” of the electrical complex.
- Prehospital EMS systems may use 12-lead ECGs to discover signs of acute MI, such as ST segment elevation, in preparation for in-hospital administration of thrombolytic drugs.
- Once a 12-lead ECG is performed, a 15-lead, or right-sided, ECG may be used for an even more comprehensive view if the right ventricle or the posterior portion of the heart appears to be affected.



Clinical Tip:

Always compare the patient's current 12-lead ECG with the previous one.



Clinical Tip:

Monitor the patient, not just the ECG, for clinical improvement.

TROUBLESHOOTING ECG PROBLEMS

Without proper assessment and treatment, a patient with an abnormal ECG could have a potentially fatal outcome. An accurate and properly monitored patient ECG is extremely important, so remember the following troubleshooting tips:

- Place leads in the correct position. Incorrect placement can give false readings.
- Avoid placing leads over bony areas.
- In patients with large breasts, place the electrodes under the breast. The most accurate tracings are obtained through the smallest amount of fat tissue.




Figure 10.1 ■ Normal R-wave progression in chest leads V_1 - V_6 .

- Apply tincture of benzoin to the electrode sites if the patient is diaphoretic. The electrodes will adhere to the skin better.
- Shave hair at the electrode site if it interferes with contact between the electrode and skin.
- Discard old electrodes and use new ones if the gel on the back of the electrode dries.

R WAVE PROGRESSION

Normal ventricular depolarization in the heart progresses from right to left and from front to back (FIG. 10-1). In a normal heart, the R wave becomes taller and the S wave smaller as electrical activity crosses the heart from right to left. This phenomenon is called R wave progression and is noted on the chest leads.

Alteration in the normal R wave progression may be seen in left ventricular hypertrophy, COPD, left BBB, or anteroseptal MI.

ELECTRICAL AXIS DEVIATION

The electrical axis is the sum total of all electrical currents generated by the ventricular myocardium during depolarization (FIG. 10-2). Analysis of the axis may help to determine the location and extent of cardiac injury, such as ventricular hypertrophy, BBB, or changes in the position of the heart in the chest (from, e.g., pregnancy or ascites).

The direction of the QRS complex in leads I and aVF determines the axis quadrant in relation to the heart.



Clinical Tip:

Extreme right axis deviation is also called indeterminate, "no man's land," and "northwest."





Figure 10.2 ■ Electrical axis deviation of the heart.

ISCHEMIA, INJURY, AND INFARCTION

Ischemia, injury, and infarction are three stages that cardiac tissue goes through when a complete blockage occurs in a coronary artery. The location of the myocardial infarction is a critical factor in determining the most appropriate treatment and predicting probable complications.

Each coronary artery delivers blood to specific areas of the heart. Blockages at different sites can damage various parts of the heart as shown in FIGURES 10-3 and 10-4. Characteristic ECG changes occur in different leads with each type of MI (FIG. 10-5) and can be correlated to the blockages shown in Figures 10-3 and 10-4.



Anterior view

Figure 10.3 ■ Anterior wall of the heart.




Figure 10.4 ■ Septal, lateral, and inferior walls of the heart.

I lateral	aVR	V ₁ septal	V ₄ anterior
II inferior	aVL lateral	V ₂ septal	V ₄ lateral
III inferior	aVF inferior	V ₃ anterior	V ₄ lateral

Figure 10.5 ■ Location of MI by ECG leads.



Clinical Tip:

Lead aVR may not show any change in an MI.



Clinical Tip:

An MI may not be limited to just one region of the heart. For example, if there are changes in leads V₃ and V₄ (anterior) and leads I, aVL, V₅, and V₆ (lateral), the MI is called an anterolateral infarction.

PROGRESSION OF AN ACUTE MYOCARDIAL INFARCTION

An acute MI is a continuum that extends from the normal state to a full infarction (FIG. 10-6):

- Ischemia—Lack of oxygen to the cardiac tissue, represented by ST segment depression, T wave inversion, or both
- Injury—Arterial occlusion with ischemia, represented by ST segment elevation




Figure 10.6 ■ Progression of an acute MI.

- Infarction—Death of tissue, represented by a pathological Q wave



Clinical Tip:

Once the acute MI has ended, the ST segment returns to baseline and the T wave becomes upright, but the Q wave remains abnormal because of scar formation.

ST SEGMENT ELEVATION AND DEPRESSION

A normal ST segment represents early ventricular repolarization. Displacement of the ST segment can be caused by various conditions, shown in FIGURES 10-7 through 10-9.




Figure 10.7 ■ ST segment at baseline.

Primary Causes of ST Segment Elevation

- ST segment elevation exceeding 1 mm in the limb leads and 2 mm in the chest leads indicates an evolving acute MI until there is proof to the contrary.

- Early repolarization (normal variant in young adults)
- Pericarditis
- Ventricular aneurysm
- Pulmonary embolism
- Intracranial hemorrhage




Figure 10.8 ■ Elevated ST segment.

Primary Causes of ST Segment Depression

- Myocardial ischemia
- Left ventricular hypertrophy
- Intraventricular conduction defects
- Medication (e.g., digitalis)
- Reciprocal changes in leads opposite the area of acute injury




Figure 10.9 ■ Depressed ST segment.

THE NORMAL 12-LEAD ECG

FIGURE 10-10 shows an example of a normal 12-lead ECG.



Clinical Tip:

A normal ECG does not rule out any acute coronary syndrome.

ANTERIOR MYOCARDIAL INFARCTION

- Occlusion of the left coronary artery—left anterior descending branch
- ECG changes: ST segment elevation with tall T waves and taller-than-normal R waves in leads V₃ and V₄; reciprocal changes in II, III, and aVF

FIGURE 10-11 shows ECG changes typical of an anterior MI.



Clinical Tip:

Anterior MI frequently involves a large area of the myocardium and can present with cardiogenic shock, second-degree AV block Type II, or third-degree AV block.




Figure 10.10 ■ Normal 12-lead ECG.




Figure 10.11 ■ Anterior myocardial infarction.

INFERIOR MYOCARDIAL INFARCTION

- Occlusion of the right coronary artery—posterior descending branch

- ECG changes: ST segment elevation in leads II, III, and aVF; reciprocal ST segment depression in I and aVL

FIGURE 10-12 shows a tracing typical of inferior MI.




Figure 10.12 ■ Inferior myocardial infarction.



Clinical Tip:

Be alert for symptomatic sinus bradycardia, AV blocks, hypotension, and hypoperfusion.

LATERAL MYOCARDIAL INFARCTION

- Occlusion of the left coronary artery—circumflex branch
- ECG changes: ST segment elevation in leads I, aVL, V₅, and V₆; reciprocal ST segment depression in V₁, V₂, and V₃

FIGURE 10-13 shows characteristic ECG changes in lateral MI.



Clinical Tip:

Lateral MI is often associated with anterior or inferior wall MI. Be alert for changes that may indicate cardiogenic shock or congestive heart failure.




Figure 10.13 ■ Lateral myocardial infarction.

SEPTAL MYOCARDIAL INFARCTION

- Occlusion of the left coronary artery—left anterior descending branch
- ECG changes: pathological Q waves; absence of normal R waves in leads V₁ and V₂

FIGURE 10-14 shows a tracing characteristic of septal MI.




Figure 10.14 ■ Septal myocardial infarction.



Clinical Tip:

Septal MI is often associated with an anterior wall MI.

POSTERIOR MYOCARDIAL INFARCTION

- Occlusion of the right coronary artery (posterior descending branch) or the left circumflex artery
- Usually produces tall R waves and ST segment depression in leads V_1 , V_2 , V_3 , and V_4 . Complications may include left ventricular dysfunction.
- You may need to view the true posterior leads, V_8 and V_9 (used in the 15-lead ECG) for definite diagnosis of an acute posterior MI. In these leads you will see ST segment elevation.

FIGURE 10-15 shows typical ECG changes in posterior MI.



Clinical Tip:

Diagnosis may require a 15-lead ECG because a standard 12-lead does not look directly at the posterior wall.

LEFT BUNDLE BRANCH BLOCK

- QRS complex greater than 0.10 sec
- QRS predominantly negative in leads V_1 and V_2




Figure 10.15 ■ Posterior myocardial infarction.

- QRS predominantly positive in V_5 and V_6 and often notched
- Absence of small, normal Q waves in I, aVL, V_5 , and V_6
- Wide monophasic R waves in I, aVL, V_1 , V_5 , and V_6

FIGURE 10-16 shows an example of left BBB.




Figure 10.16 ■ Left bundle branch block.



Clinical Tip:

Patients may have underlying heart disease, including coronary artery disease, hypertension, cardiomyopathy, and ischemia.

RIGHT BUNDLE BRANCH BLOCK

- QRS complex greater than 0.10 sec
- QRS axis normal or deviated to the right
- Broad S wave in leads I, aVL, V_5 , and V_6
- RSR' pattern in lead V_1 with R' taller than R
- qRS pattern in V_5 and V_6
- ST-T distorted and in opposite direction to terminal portion of QRS (this is not ST elevation or ST depression).

FIGURE 10-17 shows an example of right BBB.



Clinical Tip:

Patients may have underlying right ventricular hypertrophy, pulmonary edema, cardiomyopathy, congenital heart disease, or rheumatic heart disease.



Clinical Tip:

In bundle branch blocks, the ST segment and T waves are distorted and are in the opposite direction to the terminal portion of the QRS. This is not ST elevation or ST depression.





Figure 10.17 ■ Right bundle branch block.

UNIT THREE

ECG Practice Tests




ECG Practice Test One

For instructions on analyzing these practice test strips, please see the guidelines given at the end of Chapter 2. To help you with your success in ECG interpretation, hints have been provided throughout this chapter. By the time you tackle Chapters 12 through 14, you will have enough experience to proceed without any hints.

TEST STRIP SECTION ONE ■

ECG 11•1




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____


Interpretation: _____

ECG 11•2



Rate: _____ Rhythm: _____
 P Waves: _____ PR Interval: _____ QRS: _____
 Interpretation: _____

ECG 11•3



Hint: Notice the single spike marks.


Rate: _____ Rhythm: _____
 P Waves: _____ PR Interval: _____ QRS: _____
 Interpretation: _____

ECG 11•4



Rate: _____ Rhythm: _____
 P Waves: _____ PR Interval: _____ QRS: _____
 Interpretation: _____

ECG 11•5




Hint: Check to see if there is a full compensatory or noncompensatory pause.

Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•6




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•7




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•8



Hint: Measure carefully to see if the rhythm is regular or irregular.


Rate: _____ Rhythm: _____
 P Waves: _____ PR Interval: _____ QRS: _____
 Interpretation: _____

ECG 11•9



Rate: _____ Rhythm: _____
 P Waves: _____ PR Interval: _____ QRS: _____
 Interpretation: _____

ECG 11•10



Rate: _____ Rhythm: _____
 P Waves: _____ PR Interval: _____ QRS: _____
 Interpretation: _____

ECG 11•11



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•12




Hint: The PR intervals become progressively longer until one P wave is blocked.

Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•13



Hint: Notice that the P waves are normal (upright and uniform).

Rate: _____ Rhythm: _____
 P Waves: _____ PR Interval: _____ QRS: _____
 Interpretation: _____

ECG 11•14



Rate: _____ Rhythm: _____
 P Waves: _____ PR Interval: _____ QRS: _____
 Interpretation: _____

ECG 11•15



Rate: _____ Rhythm: _____
 P Waves: _____ PR Interval: _____ QRS: _____
 Interpretation: _____

ECG 11•16



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•17



Hint: Notice that the PR interval is prolonged.

Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•18




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____


ECG 11•19



Hint: Notice the rate and the inverted P waves.

Rate: _____ Rhythm: _____
 P Waves: _____ PR Interval: _____ QRS: _____
 Interpretation: _____

ECG 11•20




Rate: _____ Rhythm: _____
 P Waves: _____ PR Interval: _____ QRS: _____
 Interpretation: _____

ECG 11•21



Rate: _____ Rhythm: _____
 P Waves: _____ PR Interval: _____ QRS: _____
 Interpretation: _____

ECG 11•22



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•23




Hint: Notice that the blocks are an even interval of the underlying rhythm; after the dropped beats, the cycles continue on time.

Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•24




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____


Interpretation: _____

ECG 11•25



Rate: _____ Rhythm: _____
 P Waves: _____ PR Interval: _____ QRS: _____
 Interpretation: _____

ECG 11•26



Rate: _____ Rhythm: _____
 P Waves: _____ PR Interval: _____ QRS: _____
 Interpretation: _____


ECG 11•27



Hint: Conduction between the atria and the ventricles is absent.

Rate: _____ Rhythm: _____
 P Waves: _____ PR Interval: _____ QRS: _____
 Interpretation: _____

ECG 11•28




Hint: Notice that the first sinus beat occurs after the pacemaker voltage is increased.

Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•29



Hint: Notice whether the rhythm is regular or irregular.

Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____


Interpretation: _____

ECG 11•30



Rate: _____ Rhythm: _____
 P Waves: _____ PR Interval: _____ QRS: _____
 Interpretation: _____

ECG 11•31



Hint: Notice that the premature beat lands directly on the T wave.

Rate: _____ Rhythm: _____
 P Waves: _____ PR Interval: _____ QRS: _____
 Interpretation: _____

ECG 11•32



Rate: _____ Rhythm: _____
 P Waves: _____ PR Interval: _____ QRS: _____
 Interpretation: _____

ECG 11•33



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•34



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•35




Hint: Notice that beat 3 is premature.

Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•36



Rate: _____ Rhythm: _____
 P Waves: _____ PR Interval: _____ QRS: _____
 Interpretation: _____

ECG 11•37



Hint: An elevated T wave usually indicates hyperkalemia.

Rate: _____ Rhythm: _____
 P Waves: _____ PR Interval: _____ QRS: _____
 Interpretation: _____

ECG 11•38



Rate: _____ Rhythm: _____
 P Waves: _____ PR Interval: _____ QRS: _____
 Interpretation: _____

ECG 11•39




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•40




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•41



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____



Hint: Notice that the P waves have a saw-toothed appearance.

ECG 11•42




Hint: Chaotic electrical activity occurs with no ventricular depolarization or contraction.

Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•43




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•44




Hint: Notice that the rhythm converts to a different arrhythmia.

Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•45




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•46




Hint: This rhythm has no pulse.

Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•47




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•48



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•49




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•50



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•51




Hint: *This fast rhythm can start or stop suddenly.*

Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•52




Hint: Notice that each premature beat has a different form.

Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•53




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•54




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•55




Hint: *Electrical activity in the ventricles is completely absent.*

Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•56



Hint: *Notice that the waves are of the same shape and amplitude.*

Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____


Interpretation: _____

ECG 11•57




Rate: _____ Rhythm: _____
 P Waves: _____ PR Interval: _____ QRS: _____
 Interpretation: _____

ECG 11•58



Rate: _____ Rhythm: _____
 P Waves: _____ PR Interval: _____ QRS: _____
 Interpretation: _____

ECG 11•59



Hint: The P waves in this arrhythmia can be absent, inverted, or retrograde.

Rate: _____ Rhythm: _____
 P Waves: _____ PR Interval: _____ QRS: _____
 Interpretation: _____

ECG 11•60



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•61



Rate: _____ Rhythm: _____


P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____



Hint: Notice the spindle effect in the rhythm.

ECG 11•62




Hint: Notice whether the rhythm is regular or irregular.

Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•63



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•64



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•65



Hint: Notice that the T wave is elevated.

Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•66




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•67




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•68




 **Hint:** Notice that every third complex is a premature beat with a full compensatory pause.

Rate: _____ Rhythm: _____
 P Waves: _____ PR Interval: _____ QRS: _____
 Interpretation: _____


ECG 11•69



 **Hint:** Notice that the dropped beats are at even intervals with the underlying rhythm.

Rate: _____ Rhythm: _____
 P Waves: _____ PR Interval: _____ QRS: _____
 Interpretation: _____

ECG 11•70




Hint: Notice that the P waves are buried in the T waves.

Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•71




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•72




Hint: Notice the pacemaker spike before each P wave.

Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•73




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•74



Hint: Notice that beats 2, 6, and 8 are all premature.

Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 11•75



Hint: Notice the dual spike marks.

Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

Answers to Practice Test One

■ ECG 11•1

Rate: 38 bpm
 Rhythm: Regular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.10 sec
 Interpretation: Sinus bradycardia

■ ECG 11•2

Rate: 136 bpm
 Rhythm: Regular
 P Waves: Normal but encroach on preceding T waves
 PR Interval: 0.12 sec
 QRS: 0.08 sec
 Interpretation: Sinus tachycardia

■ ECG 11•3

Rate: 68 bpm
 Rhythm: Regular
 P Waves: Present following pacemaker spike
 PR Interval: 0.16 sec
 QRS: 0.06 sec
 Interpretation: Pacemaker—atrial; ventricular sensed

■ ECG 11•4

Rate: 38 bpm
 Rhythm: Regular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.10 sec
 Interpretation: Sinus bradycardia with ST segment depression and inverted T waves

■ ECG 11•5

Rate: 90 bpm (counting PVCs), 75 bpm in now underlying rhythm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.10 sec
 Interpretation: Normal sinus rhythm with two multi-form PVCs with noncompensatory pauses

■ ECG 11•6

Rate: 90 bpm
 Rhythm: Irregular
 P Waves: Flutter waves
 PR Interval: None
 QRS: 0.08 sec
 Interpretation: Atrial flutter with variable block

■ ECG 11•7

Rate: 150 bpm
 Rhythm: Regular
 P Waves: Normal
 PR Interval: 0.12 sec
 QRS: 0.08 sec
 Interpretation: Sinus tachycardia

■ ECG 11•8

Rate: 60 bpm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.20 sec
 QRS: 0.10 sec
 Interpretation: Sinus arrhythmia with respiration artifact

■ ECG 11•9

Rate: 88 bpm
 Rhythm: Regular
 P Waves: Normal
 PR Interval: 0.20 sec
 QRS: 0.10 sec
 Interpretation: Normal sinus rhythm with ST segment depression and inverted T waves

■ ECG 11•10

Rate: 54 bpm
 Rhythm: Regular
 P Waves: None
 PR Interval: None
 QRS: 0.12 sec
 Interpretation: Junctional rhythm with bundle branch block

■ ECG 11•11

Rate: 88 bpm
 Rhythm: Regular
 P Waves: None
 PR Interval: None
 QRS: 0.16 sec
 Interpretation: Accelerated idioventricular rhythm

■ ECG 11•12

Rate: 60 bpm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: Progressive prolongation until dropped QRS
 QRS: 0.10 sec
 Interpretation: Second-degree AV block Type I (Wenckebach)

■ **ECG 11•13**

Rate: 78 bpm
Rhythm: Regular
P Waves: Normal
PR Interval: 0.16 sec
QRS: 0.10 sec
Interpretation: Normal sinus rhythm

■ **ECG 11•14**

Rate: 88 bpm
Rhythm: Regular
P Waves: Normal
PR Interval: 0.20 sec
QRS: 0.08 sec
Interpretation: Normal sinus rhythm with ST segment elevation

■ **ECG 11•15**

Rate: 50 bpm
Rhythm: Irregular
P Waves: None
PR Interval: None
QRS: 0.20 sec following pacemaker spike
Interpretation: Pacemaker—ventricular with intermittent loss of capture

■ **ECG 11•16**

Rate: 120 bpm (counting PVCs), underlying rhythm is 125 bpm
Rhythm: Irregular
P Waves: Normal
PR Interval: 0.16 sec
QRS: 0.08 sec
Interpretation: Sinus tachycardia with ST segment depression and couplet PVCs

■ **ECG 11•17**

Rate: 68 bpm
Rhythm: Regular
P Waves: Normal
PR Interval: 0.28 sec
QRS: 0.10 sec
Interpretation: Normal sinus rhythm with first-degree AV block

■ **ECG 11•18**

Rate: 50 bpm
Rhythm: Irregular
P Waves: None
PR Interval: None
QRS: 0.10 sec
Interpretation: Atrial fibrillation with ST segment depression and slow ventricular response

■ **ECG 11•19**

Rate: 71 bpm
Rhythm: Regular
P Waves: Inverted
PR Interval: 0.08 sec
QRS: 0.08 sec
Interpretation: Accelerated junctional rhythm

■ **ECG 11•20**

Rate: 70 bpm
Rhythm: Irregular
P Waves: Normal
PR Interval: 0.32 sec for beats 2 through 6
QRS: 0.10 sec
Interpretation: Normal sinus rhythm first-degree AV block changing to second-degree AV block type I

■ **ECG 11•21**

Rate: 80 bpm (counting PVCs), 88 bpm for underlying rhythm
Rhythm: Irregular
P Waves: Normal
PR Interval: 0.16 sec
QRS: 0.10 sec
Interpretation: Normal sinus rhythm with PVCs at beats 2 and 7

■ **ECG 11•22**

Rate: 90 bpm (counting PVCs), 75 bpm for underlying rhythm
Rhythm: Irregular
P Waves: Normal
PR Interval: 0.16 sec
QRS: 0.10 sec
Interpretation: Normal sinus rhythm with multiform PVCs at beats 6 and 8

■ **ECG 11•23**

Rate: 50 bpm, 68 bpm for underlying rhythm
Rhythm: Irregular
P Waves: Normal
PR Interval: 0.16 sec
QRS: 0.10 sec
Interpretation: Normal sinus rhythm with sinus block (two blocked beats)

■ **ECG 11•24**

Rate: Indeterminate
Rhythm: Chaotic
P Waves: None
PR Interval: None
QRS: None
Interpretation: Ventricular fibrillation—coarse fibrillatory waves

■ **ECG 11•25**

Rate: 68 bpm
 Rhythm: Regular
 P Waves: Normal
 PR Interval: 0.12 sec
 QRS: 0.12 sec with notched appearance
 Interpretation: Normal sinus rhythm with a bundle branch block

■ **ECG 11•26**

Rate: Indeterminate
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: Variable
 Interpretation: Ventricular fibrillation with defibrillation converting to atrial fibrillation

■ **ECG 11•27**

Rate: Atrial 90 bpm, ventricular 35 bpm
 Rhythm: Atrial irregular, ventricular regular
 P Waves: Normal but not associated with QRS
 PR Interval: Variable
 QRS: 0.14 sec
 Interpretation: Third-degree AV block

■ **ECG 11•28**

Rate: 70 bpm in second half of strip
 Rhythm: Irregular
 P Waves: Normal following pacemaker spike
 PR Interval: 0.16 sec
 QRS: 0.10 sec
 Interpretation: Pacemaker—atrial with failure to capture at first three pacemaker spikes. When the current is increased, there is capture at pacemaker spikes 4 through 7.

■ **ECG 11•29**

Rate: 100 bpm
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: 0.12 sec
 Interpretation: Atrial fibrillation with a bundle branch block

■ **ECG 11•30**

Rate: 90 bpm; 75 bpm in underlying rhythm, over 100 bpm in VT
 Rhythm: Irregular
 P Waves: Not seen due to artifact
 PR Interval: Indeterminate due to artifact
 QRS: 0.08 sec for beats 1 through 4
 Interpretation: Probably normal sinus rhythm with muscle artifact converting to ventricular tachycardia at beat 5

■ **ECG 11•31**

Rate: 60 bpm (counting PVCs), 38 bpm in underlying rhythm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.10 sec
 Interpretation: Sinus bradycardia with interpolated PVCs (R on T)

■ **ECG 11•32**

Rate: 56 bpm
 Rhythm: Regular
 P Waves: Normal
 PR Interval: 0.12 sec
 QRS: 0.10 sec
 Interpretation: Sinus bradycardia with respiration artifact

■ **ECG 11•33**

Rate: 70 bpm
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: 0.08 sec
 Interpretation: Atrial fibrillation with ST segment depression

■ **ECG 11•34**

Rate: 40 bpm
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: 0.10 sec
 Interpretation: Atrial fibrillation with slow ventricular response

■ **ECG 11•35**

Rate: 70 bpm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.10 sec
 Interpretation: Normal sinus rhythm with one PAC at beat 3

■ **ECG 11•36**

Rate: 80 bpm
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: 0.08 sec
 Interpretation: Accelerated junctional rhythm with ST segment elevation and PJC's at beats 2 and 4

■ **ECG 11•37**

Rate: 79 bpm
Rhythm: Regular
P Waves: Normal
PR Interval: 0.12 sec
QRS: 0.08 sec
Interpretation: Normal sinus rhythm with peaked T waves

■ **ECG 11•38**

Rate: 88 bpm
Rhythm: Regular
P Waves: Normal
PR Interval: 0.12 sec
QRS: 0.10 sec
Interpretation: Normal sinus rhythm with ST segment elevation and inverted T waves

■ **ECG 11•39**

Rate: Atrial 79 bpm, ventricular 47 bpm
Rhythm: Atrial regular, ventricular regular
P Waves: Normal but not associated with QRS
PR Interval: Variable
QRS: 0.16 sec
Interpretation: Third-degree AV block

■ **ECG 11•40**

Rate: 50 bpm
Rhythm: Irregular
P Waves: Flutter waves
PR Interval: None
QRS: 0.08 sec
Interpretation: Atrial flutter with variable block

■ **ECG 11•41**

Rate: 80 bpm (counting PVCs), 71 bpm in underlying rhythm
Rhythm: Irregular
P Waves: Normal
PR Interval: 0.14 sec
QRS: 0.10 sec
Interpretation: Normal sinus rhythm with U waves and two uniform PVCs with noncompensatory pauses

■ **ECG 11•42**

Rate: Indeterminate
Rhythm: Chaotic
P Waves: None
PR Interval: None
QRS: None
Interpretation: Ventricular fibrillation

■ **ECG 11•43**

Rate: 50 bpm
Rhythm: Regular
P Waves: Normal
PR Interval: 0.20 sec
QRS: 0.08 sec
Interpretation: Sinus bradycardia

■ **ECG 11•44**

Rate: 100 bpm
Rhythm: Irregular
P Waves: Flutter waves changing to fibrillatory waves
PR Interval: Indeterminate
QRS: 0.08 sec
Interpretation: Atrial flutter converting to atrial fibrillation

■ **ECG 11•45**

Rate: 30 bpm
Rhythm: Irregular
P Waves: Normal
PR Interval: 0.16 sec
QRS: 0.10 sec
Interpretation: Sinus pause (sinus arrest) with 4.28-sec pause (arrest)

■ **ECG 11•46**

Rate: 150 bpm before and after defibrillation
Rhythm: Regular in section before and after defibrillation
P Waves: None
PR Interval: None
QRS: Wide—greater than 0.10 sec
Interpretation: After checking the patient, there are no pulses, ventricular tachycardia defibrillated, converting back to ventricular tachycardia with no pulses

■ **ECG 11•47**

Rate: 90 bpm
Rhythm: Irregular
P Waves: Normal
PR Interval: 0.24 sec
QRS: 0.08 sec
Interpretation: Normal sinus rhythm with first-degree AV block including ST segment elevation and bigeminal PJC's

■ **ECG 11•48**

Rate: 110 bpm
Rhythm: Irregular
P Waves: None
PR Interval: None
QRS: 0.10 sec
Interpretation: Atrial fibrillation with rapid ventricular response

■ **ECG 11•49**

Rate: 80 bpm
 Rhythm: Regular
 P Waves: Present with low amplitude following pacemaker spike
 PR Interval: 0.20 sec
 QRS: 0.18 sec following pacemaker spike
 Interpretation: Pacemaker—atrial and ventricular

■ **ECG 11•50**

Rate: 76 bpm in first section; 170–180 bpm in second section
 Rhythm: Irregular
 P Waves: Normal in first five beats
 PR Interval: 0.12 sec in first five beats
 QRS: 0.10 sec in the first five beats changing to a wide QRS—greater than 0.10 sec in the last seven ventricular beats
 Interpretation: Normal sinus rhythm converting to ventricular tachycardia

■ **ECG 11•51**

Rate: 150 bpm; 214 bpm in first section; 100 bpm in second section
 Rhythm: Irregular
 P Waves: Buried in T wave in beats 1 through 9, none in beat 10, and normal in beats 11 through 15
 PR Interval: None in beats 1 through 10, 0.16 sec in beats 11 through 15
 QRS: 0.08 sec
 Interpretation: Paroxysmal supraventricular tachycardia (supraventricular tachycardia converting to normal sinus rhythm)

■ **ECG 11•52**

Rate: 70 bpm (counting PVCs), 68 bpm in underlying rhythm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.08 sec
 Interpretation: Normal sinus rhythm with multiform PVCs at beats 2 and 5 with full compensatory pauses

■ **ECG 11•53**

Rate: 80 bpm
 Rhythm: Irregular
 P Waves: Present with low amplitude
 PR Interval: 0.20 sec
 QRS: 0.08 sec
 Interpretation: Normal sinus rhythm with two PACs at beats 4 and 7

■ **ECG 11•54**

Rate: Indeterminate
 Rhythm: Chaotic
 P Waves: None
 PR Interval: None
 QRS: First five beats wide—greater than 0.10 sec; for remainder of rhythm—no true QRS complexes
 Interpretation: Ventricular tachycardia deteriorating into ventricular fibrillation with coarse fibrillatory waves

■ **ECG 11•55**

Rate: None
 Rhythm: None
 P Waves: None
 PR Interval: None
 QRS: None
 Interpretation: Asystole

■ **ECG 11•56**

Rate: 214 bpm
 Rhythm: Regular
 P Waves: None
 PR Interval: None
 QRS: Wide—greater than 0.10 sec
 Interpretation: Ventricular tachycardia—monomorphic

■ **ECG 11•57**

Rate: 80 bpm
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: 0.12 sec following pacemaker spike
 Interpretation: Pacemaker—atrial and ventricular, with no capture of the P waves and with a short run of ventricular tachycardia

■ **ECG 11•58**

Rate: 110 bpm
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: 0.08 sec
 Interpretation: Atrial fibrillation with rapid ventricular response

■ **ECG 11•59**

Rate: 48 bpm
 Rhythm: Regular
 P Waves: Absent or inverted
 PR Interval: Varies
 QRS: 0.10 sec
 Interpretation: Junctional rhythm

■ **ECG 11•60**

Rate: Indeterminate
Rhythm: Irregular
P Waves: None
PR Interval: None
QRS: Wide—greater than 0.10 sec
Interpretation: Ventricular tachycardia—
polymorphic vs. artifact: assess patient

■ **ECG 11•61**

Rate: Indeterminate
Rhythm: Irregular
P Waves: None
PR Interval: None
QRS: Wide—greater than 0.10 sec
Interpretation: Torsade de pointes

■ **ECG 11•62**

Rate: 83 bpm
Rhythm: Regular
P Waves: Normal but distorted due to artifact
PR Interval: Indeterminate
QRS: 0.10 sec
Interpretation: Normal sinus rhythm with muscle
artifact

■ **ECG 11•63**

Rate: 40 bpm
Rhythm: Irregular
P Waves: Normal in beat 2
PR Interval: 0.20 sec in beat 2
QRS: 0.10 sec in beat 2; wide—greater than
0.10 sec—in beats 1, 3, and 4 following
pacemaker spike
Interpretation: Pacemaker—ventricular with
intrinsic sinus complex at beat 2

■ **ECG 11•64**

Rate: 136 bpm
Rhythm: Regular
P Waves: Normal
PR Interval: 0.12 sec
QRS: 0.08 sec
Interpretation: Sinus tachycardia with artifact
after beat 3

■ **ECG 11•65**

Rate: 125 bpm
Rhythm: Regular
P Waves: Normal
PR Interval: 0.20 sec
QRS: 0.10 sec
Interpretation: Sinus tachycardia with peaked
T waves

■ **ECG 11•66**

Rate: 60 bpm
Rhythm: Irregular
P Waves: Normal but not associated with QRS
PR Interval: Variable
QRS: 0.16 sec
Interpretation: Third-degree AV block with
couplet PVCs

■ **ECG 11•67**

Rate: 115 bpm
Rhythm: Regular
P Waves: Normal
PR Interval: 0.12 sec
QRS: 0.10 sec
Interpretation: Sinus tachycardia

■ **ECG 11•68**

Rate: 90 bpm
Rhythm: Irregular
P Waves: Normal
PR Interval: 0.16 sec
QRS: 0.08 sec
Interpretation: Normal sinus rhythm with ventricular
trigeminy

■ **ECG 11•69**

Rate: 60 bpm
Rhythm: Irregular
P Waves: Normal
PR Interval: 0.16 sec
QRS: 0.10 sec
Interpretation: Normal sinus rhythm with sinoatrial
blocks after beats 3 and 4

■ **ECG 11•70**

Rate: 250 bpm
Rhythm: Regular
P Waves: Buried in T waves
PR Interval: Not measurable
QRS: 0.08 sec
Interpretation: Supraventricular
tachycardia

■ **ECG 11•71**

Rate: 60 bpm
Rhythm: Irregular
P Waves: None
PR Interval: None
QRS: 0.16 sec
Interpretation: Atrial fibrillation with
bundle branch block

■ ECG 11•72

Rate: 94 bpm
Rhythm: Regular
P Waves: Normal following pacemaker spike
PR Interval: 0.16 sec
QRS: 0.10 sec
Interpretation: Pacemaker—atrial; ventricular

■ ECG 11•73

Rate: Indeterminate
Rhythm: Irregular
P Waves: None
PR Interval: None
QRS: Wide—greater than 0.10 sec
Interpretation: Ventricular tachycardia—
polymorphic

■ ECG 11•74

Rate: 90 bpm
Rhythm: Irregular
P Waves: Normal
PR Interval: 0.12 sec
QRS: 0.10 sec
Interpretation: Normal sinus rhythm with PJs
at beats 2, 6, and 8

■ ECG 11•75

Rate: 75 bpm
Rhythm: Regular
P Waves: Present following pacemaker spike
PR Interval: 0.20 sec
QRS: 0.16 sec following pacemaker spike
Interpretation: Pacemaker—atrial and ventricular

ECG Practice Test Two

For instructions on analyzing these practice test strips, please see the guidelines given at the end of chapter 2.

TEST STRIP SECTION TWO ■

ECG 12•1




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•2




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•3



Rate: _____ Rhythm: _____
P Waves: _____ PR Interval: _____ QRS: _____
Interpretation: _____

ECG 12•4




Rate: _____ Rhythm: _____
P Waves: _____ PR Interval: _____ QRS: _____
Interpretation: _____

ECG 12•5



Rate: _____ Rhythm: _____
P Waves: _____ PR Interval: _____ QRS: _____
Interpretation: _____

ECG 12•6




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•7



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•8




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•9



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•10



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•11




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•12



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•13



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•14



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•15



Rate: _____ Rhythm: _____
P Waves: _____ PR Interval: _____ QRS: _____
Interpretation: _____

ECG 12•16



Rate: _____ Rhythm: _____
P Waves: _____ PR Interval: _____ QRS: _____
Interpretation: _____

ECG 12•17



Rate: _____ Rhythm: _____
P Waves: _____ PR Interval: _____ QRS: _____
Interpretation: _____

ECG 12•18




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•19




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•20



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•21




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•22




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•23




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•24




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•25




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•26




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•27




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•28




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•29




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•30



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•31



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•32



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•33




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•34




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•35



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•36



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•37



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•38



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•39




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•40




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•41




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•42



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•43



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•44



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•45




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•46




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•47




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•48



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•49




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•50




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•51




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•52




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•53




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•54




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•55



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•56



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•57



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•58




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•59




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•60



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•61



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•62




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•63




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•64




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•65




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•66



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•67



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•68



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•69



Rate: _____ Rhythm: _____
P Waves: _____ PR Interval: _____ QRS: _____
Interpretation: _____

ECG 12•70




Rate: _____ Rhythm: _____
P Waves: _____ PR Interval: _____ QRS: _____
Interpretation: _____

ECG 12•71



Rate: _____ Rhythm: _____
P Waves: _____ PR Interval: _____ QRS: _____
Interpretation: _____

ECG 12•72




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•73




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•74




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 12•75



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

Answers to Practice Test Two

■ ECG 12•1

Rate: 68 bpm
 Rhythm: Regular
 P Waves: Flutter waves
 PR Interval: None
 QRS: 0.08 sec
 Interpretation: Atrial flutter with 4:1 block

■ ECG 12•2

Rate: 70 bpm
 Rhythm: Irregular
 P Waves: Vary in form
 PR Interval: 0.16 sec
 QRS: 0.10 sec
 Interpretation: Wandering atrial pacemaker

■ ECG 12•3

Rate: 80 bpm (counting PVCs), 68 bpm in underlying rhythm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.10 sec
 Interpretation: Normal sinus rhythm with three-beat ventricular tachycardia (triplet PVCs)

■ ECG 12•4

Rate: 120 bpm, 115 bpm in underlying rhythm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.08 sec
 Interpretation: Sinus tachycardia with PACs at beats 6 and 12

■ ECG 12•5

Rate: 60 bpm
 Rhythm: Regular
 P Waves: Normal following pacemaker spike
 PR Interval: 0.16 sec
 QRS: 0.10 sec
 Interpretation: Pacemaker—atrial paced, ventricular sensed with ST segment depression and inverted T waves

■ ECG 12•6

Rate: 188 bpm
 Rhythm: Regular
 P Waves: Not clearly visible, probably buried in T waves
 PR Interval: 0.06 sec
 QRS: 0.10 sec
 Interpretation: Atrial tachycardia

■ ECG 12•7

Rate: Indeterminate
 Rhythm: Chaotic
 P Waves: None
 PR Interval: None
 QRS: None
 Interpretation: Ventricular fibrillation with defibrillation and return to ventricular fibrillation

■ ECG 12•8

Rate: Indeterminate
 Rhythm: Chaotic
 P Waves: None
 PR Interval: None
 QRS: None
 Interpretation: Coarse ventricular fibrillation deteriorating to fine ventricular fibrillation

■ **ECG 12•9**

Rate: 80 bpm (counting PJs)
Rhythm: Irregular
P Waves: Normal except none in PJs
PR Interval: 0.16 sec in sinus beats
QRS: 0.10 sec
Interpretation: Normal sinus rhythm with bigeminal PJs

■ **ECG 12•10**

Rate: 38 bpm
Rhythm: Regular
P Waves: Normal
PR Interval: 0.16 sec
QRS: 0.08 sec
Interpretation: Sinus bradycardia

■ **ECG 12•11**

Rate: 130 bpm; 50 bpm in first section, 240 bpm in second section
Rhythm: Irregular
P Waves: None in beats 1, 2, and 4; inverted in beat 3; buried in T waves in beats 5 through 13
PR Interval: None except 0.16 sec in beat 3
QRS: 0.08 sec
Interpretation: Paroxysmal supraventricular tachycardia (junctional rhythm converting to supraventricular tachycardia)

■ **ECG 12•12**

Rate: Indeterminate
Rhythm: Irregular
P Waves: None
PR Interval: None
QRS: None
Interpretation: Ventricular fibrillation deteriorating to asystole

■ **ECG 12•13**

Rate: 70 bpm
Rhythm: Irregular
P Waves: Normal
PR Interval: 0.16 sec
QRS: 0.10 sec
Interpretation: Normal sinus rhythm with PVCs at beats 4 and 6

■ **ECG 12•14**

Rate: 30 bpm
Rhythm: Irregular
P Waves: Normal
PR Interval: 0.16 sec
QRS: 0.10 sec
Interpretation: Sinus bradycardia with a 3.48 sec pause (sinus arrest)

■ **ECG 12•15**

Rate: 100 bpm
Rhythm: Irregular
P Waves: Flutter waves
PR Interval: None
QRS: 0.08 sec
Interpretation: Atrial flutter with 4:1 conduction and couplet PVCs

■ **ECG 12•16**

Rate: 40 bpm
Rhythm: Irregular
P Waves: Normal
PR Interval: 0.16 sec
QRS: 0.10 sec
Interpretation: Sinus bradycardia with sinus arrhythmia

■ **ECG 12•17**

Rate: 90 bpm
Rhythm: Irregular
P Waves: None
PR Interval: None
QRS: 0.12 sec
Interpretation: Atrial fibrillation with bundle branch block

■ **ECG 12•18**

Rate: 136 bpm
Rhythm: Regular
P Waves: Inverted
PR Interval: 0.12 sec
QRS: 0.06 sec
Interpretation: Junctional tachycardia with inverted T waves

■ **ECG 12•19**

Rate: 80 bpm
Rhythm: Irregular
P Waves: Normal with beats 1, 3, 5, and 7
PR Interval: 0.16 sec with beats 1, 3, 5, and 7
QRS: 0.12 sec
Interpretation: Normal sinus rhythm with bundle branch block, and bigeminal PVCs

■ **ECG 12•20**

Rate: 125 bpm
Rhythm: Regular
P Waves: Normal
PR Interval: 0.12 sec
QRS: 0.08 sec
Interpretation: Sinus tachycardia

■ **ECG 12•21**

Rate: 90 bpm
 Rhythm: Irregular
 P Waves: Flutter waves
 PR Interval: None
 QRS: 0.10 sec
 Interpretation: Atrial fibrillation with PVCs at beats 3, 5, and 7 changing to atrial flutter

■ **ECG 12•22**

Rate: Indeterminate
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: None
 Interpretation: Coarse ventricular fibrillation deteriorating to asystole

■ **ECG 12•23**

Rate: 110 bpm
 Rhythm: Irregular
 P Waves: Present but encroaching on previous T waves
 PR Interval: 0.16 sec
 QRS: 0.10 sec
 Interpretation: Sinus tachycardia with sinus block

■ **ECG 12•24**

Rate: 125 bpm
 Rhythm: Regular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.10 sec
 Interpretation: Sinus tachycardia with ST segment depression

■ **ECG 12•25**

Rate: 88 bpm
 Rhythm: Regular
 P Waves: None
 PR Interval: None
 QRS: Wide—greater than 0.10 sec
 Interpretation: Accelerated idioventricular rhythm

■ **ECG 12•26**

Rate: 150 bpm
 Rhythm: Irregular
 P Waves: Normal in first nine beats, then buried in T waves in last six beats
 PR Interval: 0.12 sec for first nine beats, then indeterminate for last six beats
 QRS: 0.08 sec
 Interpretation: Paroxysmal supraventricular tachycardia (sinus arrhythmia converting to supraventricular tachycardia)

■ **ECG 12•27**

Rate: 68 bpm
 Rhythm: Regular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.06 sec
 Interpretation: Normal sinus rhythm with ST segment depression

■ **ECG 12•28**

Rate: 60 bpm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.28 sec in first two beats, 0.16 sec in rest of beats
 QRS: 0.10 sec
 Interpretation: Normal sinus rhythm with first-degree AV block in beats 1 and 2 changing to sinus bradycardia

■ **ECG 12•29**

Rate: 75 bpm
 Rhythm: Regular
 P Waves: None
 PR Interval: None
 QRS: 0.10 sec
 Interpretation: Accelerated junctional rhythm with ST segment depression and inverted T waves

■ **ECG 12•30**

Rate: 48 bpm
 Rhythm: Regular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.10 sec
 Interpretation: Sinus bradycardia

■ **ECG 12•31**

Rate: Indeterminate
 Rhythm: Chaotic
 P Waves: None
 PR Interval: None
 QRS: Wide—greater than 0.10 sec
 Interpretation: Ventricular fibrillation converting to ventricular tachycardia

■ **ECG 12•32**

Rate: 120 bpm
 Rhythm: Irregular
 P Waves: Normal in first 6 beats
 PR Interval: 0.16 sec
 QRS: 0.10 sec
 Interpretation: Normal sinus rhythm with ST segment depression followed by a PVC at beat 7, with a run of supraventricular tachycardia following the PVC

■ **ECG 12•33**

Rate: 100 bpm
 Rhythm: Irregular
 P Waves: Buried in T waves in first six beats; normal in last four beats
 PR Interval: 0.08–0.12 sec
 QRS: 0.08–0.12 sec
 Interpretation: Paroxysmal supraventricular tachycardia (supraventricular tachycardia with a bundle branch block converting to normal sinus rhythm.)

■ **ECG 12•34**

Rate: 90 bpm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.10 sec
 Interpretation: Normal sinus rhythm with ST segment depression and PACs at beats 5 and 8

■ **ECG 12•35**

Rate: 70 bpm
 Rhythm: Irregular
 P Waves: Flutter waves
 PR Interval: None
 QRS: 0.10 sec
 Interpretation: Atrial flutter with a 4:1 block and one PVC at beat 3 with a full compensatory pause (every fourth flutter wave is buried in the QRS)

■ **ECG 12•36**

Rate: 40 bpm
 Rhythm: Regular
 P Waves: None
 PR Interval: None
 QRS: 0.10 sec
 Interpretation: Junctional rhythm with ST segment depression

■ **ECG 12•37**

Rate: 80 bpm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.10 sec
 Interpretation: Normal sinus rhythm with ST segment elevation and a three-beat run of ventricular tachycardia

■ **ECG 12•38**

Rate: 100 bpm
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: 0.10 sec
 Interpretation: Atrial fibrillation with ST segment depression and inverted T waves

■ **ECG 12•39**

Rate: 70 bpm
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: 0.04 sec following pacemaker spike, 0.12 in second complex
 Interpretation: Pacemaker—ventricular, with respiratory artifact and one PVC at beat 2

■ **ECG 12•40**

Rate: 88 bpm
 Rhythm: Regular
 P Waves: Pacemaker spike, with small inverted P waves following spike
 PR Interval: None
 QRS: 0.08 sec
 Interpretation: Pacemaker—atrial paced, ventricular sensed

■ **ECG 12•41**

Rate: 20 bpm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.20 sec
 QRS: 0.16 sec
 Interpretation: Sinus arrest with a 4.36 sec pause and respiratory artifact

■ **ECG 12•42**

Rate: Indeterminate
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: Wide—greater than 0.10 sec
 Interpretation: Ventricular tachycardia—polymorphic

■ **ECG 12•43**

Rate: 110 bpm
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: 0.08 sec
 Interpretation: Atrial fibrillation with a four-beat multiform ventricular tachycardia

■ **ECG 12•44**

Rate: 70 bpm
 Rhythm: Irregular
 P Waves: Flutter waves
 PR Interval: None
 QRS: 0.16 sec
 Interpretation: Atrial flutter with variable block and a bundle branch block

■ **ECG 12•45**

Rate: 115 bpm
 Rhythm: Regular
 P Waves: None
 PR Interval: None
 QRS: 0.12 sec with notched appearance
 Interpretation: Junctional tachycardia with a bundle branch block

■ **ECG 12•46**

Rate: Indeterminate
 Rhythm: Chaotic
 P Waves: None
 PR Interval: None
 QRS: None
 Interpretation: Ventricular fibrillation—fine fibrillatory waves

■ **ECG 12•47**

Rate: 155 bpm in the beginning, 68 bpm after cardioversion
 Rhythm: Irregular
 P Waves: Normal in last three beats
 PR Interval: 0.16 sec in last three beats
 QRS: Wide—greater than 0.10 sec in first three beats; 0.10 sec in last three beats
 Interpretation: Monomorphic ventricular tachycardia with synchronized cardioversion converting to normal sinus rhythm

■ **ECG 12•48**

Rate: 70 bpm
 Rhythm: Irregular
 P Waves: Normal with 60-cycle interference
 PR Interval: 0.20 sec
 QRS: 0.10 sec
 Interpretation: Normal sinus rhythm with bigeminal uniform PVCs and 60-cycle interference

■ **ECG 12•49**

Rate: 70 bpm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.08 sec
 Interpretation: Normal sinus rhythm with two 1.28 second pauses (sinus arrest)

■ **ECG 12•50**

Rate: 150 bpm
 Rhythm: Regular
 P Waves: Normal but different in shape from sinus P waves
 PR Interval: 0.12 sec
 QRS: 0.08 sec
 Interpretation: Atrial tachycardia

■ **ECG 12•51**

Rate: 40 bpm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: Variable
 QRS: 0.16 sec for first four beats
 Interpretation: Third-degree AV block converting to coarse ventricular fibrillation

■ **ECG 12•52**

Rate: 100 bpm
 Rhythm: Regular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.08 sec
 Interpretation: Normal sinus rhythm with ST segment depression

■ **ECG 12•53**

Rate: 136 bpm
 Rhythm: Regular
 P Waves: Buried in T waves
 PR Interval: Not able to measure
 QRS: 0.10 sec
 Interpretation: Supraventricular tachycardia

■ **ECG 12•54**

Rate: 60 bpm
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: 0.08 sec
 Interpretation: Atrial fibrillation

■ **ECG 12•55**

Rate: 90 bpm
 Rhythm: Irregular
 P Waves: Normal except that the P wave for beat 6 (a PAC) is buried in the preceding T wave
 PR Interval: 0.16 sec
 QRS: 0.12 sec
 Interpretation: Normal sinus rhythm with a bundle branch block and a PAC at beat 6

■ **ECG 12•56**

Rate: 160 bpm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.08 sec
 Interpretation: Sinus tachycardia with ST segment depression, inverted T waves, and three uniform PVCs at beats 2, 10, and 13 with full compensatory pauses

■ **ECG 12•57**

Rate: 50 bpm
 Rhythm: Irregular
 P Waves: Flutter waves
 PR Interval: None
 QRS: 0.10 sec for beats 1 and 5
 Interpretation: Atrial flutter with a three-beat accelerated ventricular rhythm at beats 2 through 4

■ **ECG 12•58**

Rate: 110 bpm
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: Variable
 Interpretation: Unknown underlying rhythm with multiformed accelerated ventricular rhythm and ventricular couplet

■ **ECG 12•59**

Rate: 46 bpm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.12 sec
 Interpretation: Sinus bradycardia with sinus arrhythmia with a bundle branch block

■ **ECG 12•60**

Rate: 56 bpm
 Rhythm: Regular
 P Waves: None
 PR Interval: None
 QRS: 0.18 sec
 Interpretation: Accelerated idioventricular rhythm

■ **ECG 12•61**

Rate: 50 bpm
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: 0.08 sec
 Interpretation: Junctional rhythm with ST segment depression and a PJC at beat 3

■ **ECG 12•62**

Rate: 83 bpm
 Rhythm: Regular
 P Waves: Normal
 PR Interval: 0.20 sec
 QRS: 0.08 sec
 Interpretation: Normal sinus rhythm with an interpolated PVC between beats 3 and 4

■ **ECG 12•63**

Rate: Atrial 107 bpm, ventricular 35 bpm
 Rhythm: Atrial regular, ventricular regular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.08 sec
 Interpretation: Second-degree AV block Type II with 3:1 block

■ **ECG 12•64**

Rate: 110 bpm; 61 bpm in first section, 170 bpm in next section
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: 0.16 sec
 Interpretation: Paroxysmal supraventricular tachycardia (accelerated junctional rhythm with bundle branch block followed by supraventricular tachycardia)

■ **ECG 12•65**

Rate: 60 bpm
 Rhythm: Irregular
 P Waves: Retrograde
 PR Interval: None
 QRS: 0.08 sec
 Interpretation: Junctional rhythm with multiform bigeminal PVCs

■ **ECG 12•66**

Rate: 130 bpm
 Rhythm: Regular
 P Waves: None
 PR Interval: None
 QRS: Wide—greater than 0.10 sec
 Interpretation: Ventricular tachycardia—monomorphic

■ **ECG 12•67**

Rate: 90 bpm (counting PVCs), 65 bpm for underlying rhythm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.28 sec
 QRS: 0.12 sec
 Interpretation: Normal sinus rhythm with first-degree AV block, inverted T waves, and interpolated PVCs at beats 4, 6, and 9

■ **ECG 12•68**

Rate: 50 bpm
 Rhythm: Regular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.08 sec
 Interpretation: Sinus bradycardia

■ **ECG 12•69**

Rate: 80 bpm
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: 0.16 sec
 Interpretation: Accelerated idioventricular rhythm
 with bigeminal PVCs

■ **ECG 12•70**

Rate: 120 bpm (counting PVCs), 115 bpm in underlying rhythm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.08 sec
 Interpretation: Sinus tachycardia with couplet PVCs
 at beats 7 and 8

■ **ECG 12•71**

Rate: 39 bpm in first section, 166 bpm in last section
 Rhythm: Irregular
 P Waves: Normal for first three beats
 PR Interval: 0.16 sec for first three beats
 QRS: 0.10 sec for first three beats; wide—greater
 than 0.10 sec—for remaining beats
 Interpretation: Sinus bradycardia converting to
 monomorphic ventricular tachycardia

■ **ECG 12•72**

Rate: 160 bpm
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: 0.08 sec
 Interpretation: Atrial fibrillation with rapid ventricular
 response

■ **ECG 12•73**

Rate: 100 bpm
 Rhythm: Regular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.10 sec
 Interpretation: Normal sinus rhythm

■ **ECG 12•74**

Rate: 80 bpm
 Rhythm: Irregular
 P Waves: Flutter waves
 PR Interval: None
 QRS: Difficult to measure because of flutter waves,
 probably 0.08 sec
 Interpretation: Atrial flutter with variable block

■ **ECG 12•75**

Rate: 60 bpm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: Variable
 QRS: 0.10 sec
 Interpretation: Third-degree AV block with multiform
 PVCs at beats 2 and 5

ECG Practice Test Three

For instructions on analyzing these practice test strips, please see the guidelines given at the end of chapter 2.


TEST STRIP SECTION THREE ■

ECG 13•1




Rate: _____ Rhythm: _____
P Waves: _____ PR Interval: _____ QRS: _____
Interpretation: _____

ECG 13•2



Rate: _____ Rhythm: _____
P Waves: _____ PR Interval: _____ QRS: _____
Interpretation: _____

ECG 13•3




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•4




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•5




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•6




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•7




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•8




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•9



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•10



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•11



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•12



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•13



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•14



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•15




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•16



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•17




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•18



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•19




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•20




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•21



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•22




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•23




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•24



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•25



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•26



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____


Interpretation: _____

ECG 13•27




Rate: _____ Rhythm: _____
 P Waves: _____ PR Interval: _____ QRS: _____
 Interpretation: _____

ECG 13•28




Rate: _____ Rhythm: _____
 P Waves: _____ PR Interval: _____ QRS: _____
 Interpretation: _____

ECG 13•29



Rate: _____ Rhythm: _____
 P Waves: _____ PR Interval: _____ QRS: _____
 Interpretation: _____

ECG 13•30




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•31



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•32




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____


Interpretation: _____

ECG 13•33



Rate: _____ Rhythm: _____
P Waves: _____ PR Interval: _____ QRS: _____
Interpretation: _____

ECG 13•34




Rate: _____ Rhythm: _____
P Waves: _____ PR Interval: _____ QRS: _____
Interpretation: _____

ECG 13•35



Rate: _____ Rhythm: _____
P Waves: _____ PR Interval: _____ QRS: _____
Interpretation: _____

ECG 13•36




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•37



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•38




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•39




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•40



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•41




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•42




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•43




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•44



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•45




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•46



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•47



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•48



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•49



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•50



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•51



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•52




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•53




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•54



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•55




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•56



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•57



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•58




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•59




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•60




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•61




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•62



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•63




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•64




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•65



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•66



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•67




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•68



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•69




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•70




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•71



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•72



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•73




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•74




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 13•75



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

Answers to Practice Test Three

■ ECG 13•1

Rate: 50 bpm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.10 sec
 Interpretation: Sinus bradycardia with ST segment depression, inverted T waves, and a PAC at beat 2

■ ECG 13•2

Rate: 140 bpm
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: 0.08 sec
 Interpretation: Atrial fibrillation with rapid ventricular response

■ ECG 13•3

Rate: None
 Rhythm: None
 P Waves: None
 PR Interval: None
 QRS: None
 Interpretation: Loose electrode artifact

■ ECG 13•4

Rate: 110 bpm
 Rhythm: Irregular
 P Waves: Normal for beats 1, 2, and 11
 PR Interval: 0.12 sec for beats 1, 2, and 11
 QRS: 0.08 sec for beats 1, 2, and 11
 Interpretation: Paroxysmal supraventricular tachycardia (normal sinus rhythm converting to supraventricular tachycardia and back to a sinus complex at beat 11)

■ ECG 13•5

Rate: 180 bpm
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: 0.08 sec
 Interpretation: Atrial fibrillation with rapid ventricular response and ST segment depression

■ ECG 13•6

Rate: Indeterminate
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: None
 Interpretation: Asystole converting to ventricular fibrillation

■ ECG 13•7

Rate: 60 bpm, 38 bpm in underlying rhythm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.36 sec
 QRS: 0.08 sec
 Interpretation: Sinus bradycardia with first-degree AV block, inverted T waves, and multiform interpolated PVCs at beats 3 and 5

■ ECG 13•8

Rate: 30 bpm; 48 in second half of strip
 Rhythm: Irregular
 P Waves: Only two visible, of same morphology
 PR Interval: Variable
 QRS: 0.14 sec
 Interpretation: Ventricular fibrillation converting to third-degree AV block

■ **ECG 13•9**

Rate: 150 bpm
Rhythm: Regular
P Waves: Normal but encroaching on previous T waves
PR Interval: 0.12 sec
QRS: 0.08 sec
Interpretation: Sinus tachycardia

■ **ECG 13•10**

Rate: 50 bpm
Rhythm: Irregular
P Waves: Normal
PR Interval: 0.16 sec
QRS: 0.10 sec
Interpretation: Sinus bradycardia with PJs at beats 2 and 4

■ **ECG 13•11**

Rate: 34 bpm
Rhythm: Regular
P Waves: Flutter waves
PR Interval: None
QRS: 0.10 sec
Interpretation: Atrial flutter with 8:1 block

■ **ECG 13•12**

Rate: 88 bpm
Rhythm: Regular
P Waves: None
PR Interval: None
QRS: 0.16 sec following pacemaker spike
Interpretation: Pacemaker—ventricular

■ **ECG 13•13**

Rate: 170 bpm
Rhythm: Irregular
P Waves: None
PR Interval: None
QRS: 0.08
Interpretation: Atrial fibrillation with ST segment elevation and couplet PVCs

■ **ECG 13•14**

Rate: 80 bpm
Rhythm: Irregular
P Waves: Normal
PR Interval: 0.16 sec
QRS: 0.12 sec in positive complexes, difficult to measure in beats 3 and 6
Interpretation: Normal sinus rhythm with bundle branch block and ventricular trigeminy

■ **ECG 13•15**

Rate: 110 bpm
Rhythm: Irregular
P Waves: Normal
PR Interval: 0.12 sec
QRS: 0.10 sec
Interpretation: Sinus tachycardia with inverted T waves and one PVC at beat 7

■ **ECG 13•16**

Rate: 48 bpm
Rhythm: Regular
P Waves: Inverted
PR Interval: 0.12 sec
QRS: 0.10 sec
Interpretation: Junctional rhythm

■ **ECG 13•17**

Rate: 40 bpm
Rhythm: Irregular
P Waves: Normal but not associated with QRS
PR Interval: Variable
QRS: 0.12 sec
Interpretation: Third-degree AV block

■ **ECG 13•18**

Rate: 70 bpm
Rhythm: Irregular
P Waves: None
PR Interval: None
QRS: 0.12 sec
Interpretation: Junctional rhythm with bundle branch block and bigeminal PJs at beats 2, 4, and 6

■ **ECG 13•19**

Rate: 71 bpm
Rhythm: Regular
P Waves: Normal
PR Interval: 0.16 sec
QRS: 0.12 sec with notched appearance
Interpretation: Normal sinus rhythm with bundle branch block

■ **ECG 13•20**

Rate: 50 bpm
Rhythm: Irregular
P Waves: None
PR Interval: None
QRS: 0.12 sec
Interpretation: Atrial fibrillation with ST segment depression and slow ventricular response

■ **ECG 13•21**

Rate: 65 bpm
 Rhythm: Regular
 P Waves: Normal
 PR Interval: 0.20 sec
 QRS: 0.08 sec
 Interpretation: Normal sinus rhythm with a U wave

■ **ECG 13•22**

Rate: 78 bpm
 Rhythm: Regular
 P Waves: None
 PR Interval: None
 QRS: 0.20 sec following pacemaker spike
 Interpretation: Pacemaker—ventricular

■ **ECG 13•23**

Rate: 90 bpm (counting PVCs), 94 bpm in underlying rhythm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.12 sec
 QRS: 0.10 sec
 Interpretation: Normal sinus rhythm with ventricular trigeminy

■ **ECG 13•24**

Rate: 60 bpm, 79 bpm in underlying rhythm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.10 sec
 Interpretation: Normal sinus rhythm with 2.56 second pause (sinus arrest)

■ **ECG 13•25**

Rate: 40 bpm
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: 0.08 sec
 Interpretation: Atrial fibrillation with slow ventricular response

■ **ECG 13•26**

Rate: 110 bpm
 Rhythm: Irregular
 P Waves: Fibrillatory and flutter waves
 PR Interval: None
 QRS: 0.08 sec
 Interpretation: Atrial fibrillation converting to atrial flutter with 4:1 block

■ **ECG 13•27**

Rate: 20 bpm
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: Wide—greater than 0.10 sec
 Interpretation: Idioventricular rhythm (Agonal rhythm)

■ **ECG 13•28**

Rate: 58 bpm
 Rhythm: Regular
 P Waves: Normal
 PR Interval: 0.32 sec
 QRS: 0.10 sec
 Interpretation: Sinus bradycardia with first-degree AV block

■ **ECG 13•29**

Rate: 79 bpm
 Rhythm: Regular
 P Waves: None following pacemaker spike
 PR Interval: None
 QRS: 0.16 sec following pacemaker spike
 Interpretation: Pacemaker—atrial and ventricular, with failure to capture P waves and failure to fire a ventricular spike at beat 4

■ **ECG 13•30**

Rate: 90 bpm (counting PVCs), 88 bpm in underlying rhythm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.12 sec
 QRS: 0.08 sec
 Interpretation: Normal sinus rhythm with two uniform PVCs at beats 3 and 7

■ **ECG 13•31**

Rate: 50 bpm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.20 sec
 QRS: 0.10 sec
 Interpretation: Sinus bradycardia with junctional escape at beats 1 and 3, ST segment depression, and inverted T waves

■ **ECG 13•32**

Rate: 75 bpm
 Rhythm: Regular
 P Waves: Normal following pacemaker spike
 PR Interval: 0.16 sec
 QRS: 0.10 sec
 Interpretation: Pacemaker—atrial paced, ventricular sensed

■ **ECG 13•33**

Rate: Atrial 100 bpm, ventricular 58 bpm
Rhythm: Atrial regular, ventricular regular
P Waves: Normal but not associated with QRS
PR Interval: Variable
QRS: 0.10 sec
Interpretation: Third-degree AV block

■ **ECG 13•34**

Rate: 80 bpm (counting PVCs); 38 bpm in underlying rhythm
Rhythm: Irregular
P Waves: Normal
PR Interval: 0.20 sec
QRS: 0.12 sec
Interpretation: Sinus bradycardia with a bundle branch block and bigeminal PVCs

■ **ECG 13•35**

Rate: 150 bpm
Rhythm: Regular
P Waves: Flutter waves
PR Interval: None
QRS: 0.08 sec
Interpretation: Atrial flutter with 2:1 block

■ **ECG 13•36**

Rate: 100 bpm
Rhythm: Irregular
P Waves: None
PR Interval: None
QRS: 0.08 sec
Interpretation: Atrial fibrillation

■ **ECG 13•37**

Rate: None
Rhythm: None
P Waves: None
PR Interval: None
QRS: None
Interpretation: Asystole

■ **ECG 13•38**

Rate: 80 bpm (counting PVCs), 62 bpm in underlying rhythm
Rhythm: Irregular
P Waves: Normal
PR Interval: 0.20 sec
QRS: 0.08 sec
Interpretation: Normal sinus rhythm with three-beat ventricular tachycardia (triplet PVCs at beats 3, 4, and 5)

■ **ECG 13•39**

Rate: 71 bpm
Rhythm: Regular
P Waves: Normal
PR Interval: 0.20 sec
QRS: 0.12 sec with notched appearance
Interpretation: Normal sinus rhythm with a bundle branch block

■ **ECG 13•40**

Rate: 60 bpm
Rhythm: Irregular
P Waves: None
PR Interval: None
QRS: 0.12 sec
Interpretation: Junctional rhythm with bundle branch block and interpolated PJC at beat 4

■ **ECG 13•41**

Rate: 100 bpm (counting PVCs); 75 bpm in underlying rhythm
Rhythm: Irregular
P Waves: Normal
PR Interval: 0.16 sec
QRS: 0.10 sec
Interpretation: Normal sinus rhythm with couplet PVCs

■ **ECG 13•42**

Rate: 80 bpm
Rhythm: Irregular
P Waves: Normal
PR Interval: 0.14 sec
QRS: 0.10 sec in beats 1, 4, and 8; 0.16 sec in beats 2, 3, 5, 6, and 7
Interpretation: Sinus rhythm with multiform PVCs at beats 2, 3, 5, 6, and 7

■ **ECG 13•43**

Rate: Atrial 79 bpm, ventricular 48 bpm
Rhythm: Atrial regular, ventricular regular
P Waves: Normal but not associated with QRS
PR Interval: Variable
QRS: 0.16 sec
Interpretation: Third-degree AV block

■ **ECG 13•44**

Rate: 115 bpm
Rhythm: Regular
P Waves: Retrograde
PR Interval: None
QRS: 0.08 sec
Interpretation: Junctional tachycardia

■ **ECG 13•45**

Rate: 140 bpm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.10 sec
 QRS: 0.08 sec
 Interpretation: Paroxysmal supraventricular tachycardia (sinus tachycardia converting to supraventricular tachycardia)

■ **ECG 13•46**

Rate: 50 bpm (counting PVCs), 38 bpm in underlying rhythm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.10 sec
 Interpretation: Sinus bradycardia with interpolated PVC at beat 4

■ **ECG 13•47**

Rate: 94 bpm
 Rhythm: Regular
 P Waves: Normal
 PR Interval: 0.12 sec
 QRS: 0.08 sec
 Interpretation: Normal sinus rhythm with ST segment depression

■ **ECG 13•48**

Rate: 80 bpm; 70 bpm in underlying rhythm, 90 bpm in last section
 Rhythm: Irregular
 P Waves: Normal for beats 1 through 4
 PR Interval: 0.16 sec for beats 1 through 4
 QRS: 0.10 sec for beats 1 through 4; wide—greater than 0.10 sec—for beats 5 through 8
 Interpretation: Normal sinus rhythm converting to an accelerated idioventricular rhythm

■ **ECG 13•49**

Rate: 80 bpm (counting PVCs), 75 bpm in underlying rhythm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.10 sec in beats 1, 2, 3, 5, 7, and 8; 0.20 sec in beats 4 and 6
 Interpretation: Normal sinus rhythm with uniform PVCs at beats 4 and 6 with full compensatory pauses

■ **ECG 13•50**

Rate: 75 bpm
 Rhythm: Regular
 P Waves: Unclear, may be buried in terminal portion of QRS
 PR Interval: None
 QRS: 0.16 sec
 Interpretation: Accelerated idioventricular rhythm

■ **ECG 13•51**

Rate: 75 bpm
 Rhythm: Regular
 P Waves: Normal for first four beats
 PR Interval: 0.20 sec for first four beats
 QRS: 0.08 sec
 Interpretation: Normal sinus rhythm with muscle artifact

■ **ECG 13•52**

Rate: 40 bpm
 Rhythm: Irregular
 P Waves: Normal in first two beats
 PR Interval: 0.20 sec for first two beats
 QRS: 0.10 sec for first two beats; 0.16 sec—following pacemaker spikes at beats 3 and 4
 Interpretation: Sinus bradycardia converting to a ventricular pacemaker

■ **ECG 13•53**

Rate: None
 Rhythm: None
 P Waves: None
 PR Interval: None
 QRS: None
 Interpretation: Asystole

■ **ECG 13•54**

Rate: 38 bpm
 Rhythm: Regular
 P Waves: Inverted in first two beats, upright in last two beats
 PR Interval: 0.16 sec
 QRS: 0.10 sec
 Interpretation: Wandering atrial pacemaker

■ **ECG 13•55**

Rate: 41 bpm
 Rhythm: Regular
 P Waves: Normal
 PR Interval: 0.24 sec
 QRS: 0.20 sec
 Interpretation: Sinus bradycardia with first-degree AV block and bundle branch block

■ **ECG 13•56**

Rate: 90 bpm
Rhythm: Irregular
P Waves: Normal
PR Interval: 0.16 sec
QRS: 0.10 sec
Interpretation: Sinus arrhythmia

■ **ECG 13•57**

Rate: 110 bpm
Rhythm: Irregular
P Waves: Normal except in PACs
PR Interval: 0.14 sec
QRS: 0.08 sec
Interpretation: Sinus tachycardia with atrial quadrigeminy (PACs at beats 4 and 8)

■ **ECG 13•58**

Rate: 90 bpm; 88 bpm in paced section
Rhythm: Regular in paced section, then irregular
P Waves: None
PR Interval: None
QRS: 0.16 sec
Interpretation: Pacemaker—ventricular, with polymorphic ventricular tachycardia at end of strip

■ **ECG 13•59**

Rate: Indeterminate
Rhythm: Irregular
P Waves: None
PR Interval: None
QRS: Wide—greater than 0.10 sec—with bizarre appearance
Interpretation: Torsade de pointes

■ **ECG 13•60**

Rate: 70 bpm
Rhythm: Irregular
P Waves: Different forms
PR Interval: Variable
QRS: 0.10 sec
Interpretation: Wandering atrial pacemaker

■ **ECG 13•61**

Rate: 60 bpm
Rhythm: Regular
P Waves: Normal following pacemaker spike
PR Interval: 0.28 sec
QRS: 0.14 sec
Interpretation: Pacemaker—atrial paced with first-degree AV block and ventricular sensed

■ **ECG 13•62**

Rate: Indeterminate
Rhythm: Irregular
P Waves: None
PR Interval: None
QRS: Wide—greater than 0.10 sec
Interpretation: Ventricular tachycardia deteriorating to ventricular fibrillation

■ **ECG 13•63**

Rate: 38 bpm
Rhythm: Regular
P Waves: Normal
PR Interval: 0.16 sec
QRS: 0.10 sec
Interpretation: Sinus bradycardia

■ **ECG 13•64**

Rate: 50 bpm; 78 bpm in underlying rhythm
Rhythm: Irregular
P Waves: None or inverted for first five beats
PR Interval: 0.16 sec for beats 2 through 4
QRS: 0.10 sec for first five beats
Interpretation: Junctional rhythm converting to ventricular fibrillation

■ **ECG 13•65**

Rate: 80 bpm (counting PVCs), 75 bpm in underlying rhythm
Rhythm: Irregular
P Waves: Normal
PR Interval: 0.16 sec
QRS: 0.10 sec
Interpretation: Normal sinus rhythm with ventricular trigeminy

■ **ECG 13•66**

Rate: 100 bpm (counting PVCs), 92 bpm in underlying rhythm
Rhythm: Irregular
P Waves: Present, but distorted due to muscle artifact
PR Interval: 0.20 sec
QRS: 0.08 sec
Interpretation: Normal sinus rhythm with three-beat ventricular tachycardia (triplet PVCs) and muscle artifact

■ **ECG 13•67**

Rate: 86 bpm (counting PVCs), 68 bpm in underlying rhythm
Rhythm: Irregular
P Waves: Normal following pacemaker spike
PR Interval: 0.20 sec
QRS: 0.08 sec
Interpretation: Pacemaker—atrial paced, ventricular sensed, with PVCs at beats 3, 5, and 8

■ **ECG 13•68**

Rate: 110 bpm (counting PVCs), 115 bpm in underlying rhythm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.10 sec
 Interpretation: Sinus tachycardia with ST segment depression and ventricular couplet at beats 6 and 7

■ **ECG 13•69**

Rate: 70 bpm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: Variable
 QRS: 0.08 sec
 Interpretation: Second-degree AV block Type I (Wenckebach)

■ **ECG 13•70**

Rate: 33 bpm
 Rhythm: Regular
 P Waves: None
 PR Interval: None
 QRS: 0.14 sec
 Interpretation: Idioventricular rhythm

■ **ECG 13•71**

Rate: 88 bpm
 Rhythm: Regular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: Wide—greater than 0.10 sec—notched appearance
 Interpretation: Normal sinus rhythm with a bundle branch block

■ **ECG 13•72**

Rate: 50 bpm
 Rhythm: Regular
 P Waves: Normal
 PR Interval: 0.28 sec
 QRS: 0.08 sec
 Interpretation: Sinus bradycardia with first-degree AV block

■ **ECG 13•73**

Rate: 75 bpm
 Rhythm: Regular
 P Waves: None
 PR Interval: None
 QRS: 0.16 sec
 Interpretation: Pacemaker—ventricular

■ **ECG 13•74**

Rate: 68 bpm
 Rhythm: Regular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.06 sec
 Interpretation: Normal sinus rhythm

■ **ECG 13•75**


Rate: 80 bpm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.10 sec
 Interpretation: Normal sinus rhythm with bigeminal PACs

ECG Practice Test Four

For instructions on analyzing these practice test strips, please see the guidelines given at the end of chapter 2.


TEST STRIP SECTION FOUR ■

ECG 14•1




Rate: _____ Rhythm: _____
P Waves: _____ PR Interval: _____ QRS: _____
Interpretation: _____

ECG 14•2



Rate: _____ Rhythm: _____
P Waves: _____ PR Interval: _____ QRS: _____
Interpretation: _____

ECG 14•3



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•4



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•5



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•6



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•7



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•8



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•9




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•10



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•11



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•12



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•13




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•14




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•15




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•16



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•17




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•18



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•19



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•20




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•21



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•22



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•23



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•24




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•25



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•26




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•27



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•28




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•29



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•30




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•31



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•32




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•33




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•34



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•35



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•36



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•37




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•38




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•39



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•40



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•41




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•42



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•43




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•44



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•45




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•46



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•47



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•48




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•49



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•50



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•51



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•52



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•53



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•54



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•55



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•56




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•57



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•58




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•59




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•60



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•61




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•62



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•63




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•64




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•65



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•66




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•67




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•68



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•69



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•70




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•71




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•72



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•73




Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•74



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

ECG 14•75



Rate: _____ Rhythm: _____

P Waves: _____ PR Interval: _____ QRS: _____

Interpretation: _____

Answers to Practice Test Four

■ ECG 14•1

Rate: 88 bpm
 Rhythm: Regular
 P Waves: Normal
 PR Interval: 0.18 sec
 QRS: 0.08 sec
 Interpretation: Normal sinus rhythm with ST segment elevation

■ ECG 14•2

Rate: None
 Rhythm: None
 P Waves: None
 PR Interval: None
 QRS: None
 Interpretation: Asystole and pacemaker with failure to capture

■ ECG 14•3

Rate: 40 bpm (counting PVCs), 34 bpm in underlying rhythm
 Rhythm: Irregular, regular in underlying rhythm
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.16 sec in first beat, 0.08 sec in remaining beats
 Interpretation: Sinus bradycardia with a PVC at beat 1

■ ECG 14•4

Rate: 150 bpm (counting PVCs), 150 bpm in underlying rhythm
 Rhythm: Irregular, regular in underlying rhythm
 P Waves: None
 PR Interval: None
 QRS: 0.08 sec
 Interpretation: Junctional tachycardia with couplet PVCs

■ ECG 14•5

Rate: 150 bpm
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: 0.08 sec
 Interpretation: Atrial fibrillation with rapid ventricular response

■ ECG 14•6

Rate: 70 bpm
 Rhythm: Irregular
 P Waves: Flutter waves
 PR Interval: None
 QRS: 0.10 sec
 Interpretation: Atrial flutter with variable block with atrial fibrillation from beats 2 to 4

■ ECG 14•7

Rate: 110 bpm
 Rhythm: Irregular
 P Waves: Variable
 PR Interval: Variable
 QRS: 0.10 sec in beats 1, 2, 4, 5, 6; 0.12 sec in beats 3, 7, and remaining beats
 Interpretation: Possible sinus tachycardia with ST segment depression and PVC in beat 3 and polymorphic ventricular tachycardia

■ ECG 14•8

Rate: 63 bpm
 Rhythm: Regular
 P Waves: Buried in artifact
 PR Interval: Indeterminate
 QRS: 0.08 sec
 Interpretation: Indeterminate rhythm; possibly normal sinus rhythm with muscle artifact

■ **ECG 14•9**

Rate: 80 bpm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.08 sec
 Interpretation: Normal sinus rhythm with ventricular bigeminy

■ **ECG 14•10**

Rate: 47 bpm
 Rhythm: Regular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.10 sec
 Interpretation: Sinus bradycardia

■ **ECG 14•11**

Rate: 80 bpm (counting PVCs), 69 bpm in underlying rhythm
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: 0.12 sec
 Interpretation: Junctional rhythm with a bundle branch block two multiform PVCs at beats 4 and 6 and muscle artifact at the end of the rhythm

■ **ECG 14•12**

Rate: 120 bpm
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: 0.10 sec
 Interpretation: Atrial fibrillation with rapid ventricular response and ST segment depression

■ **ECG 14•13**

Rate: 120 bpm (counting PVCs), 125 in underlying rhythm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.08 sec in narrow beats, 0.12 sec in wide beats
 Interpretation: Sinus tachycardia with ST segment depression and PVCs at beats 2, 6, 7, and 11

■ **ECG 14•14**

Rate: Indeterminate
 Rhythm: Flat line, then chaotic
 P Waves: None
 PR Interval: None
 QRS: None
 Interpretation: Asystole converting to ventricular fibrillation

■ **ECG 14•15**

Rate: Indeterminate
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: Wide—greater than 0.10 sec with bizarre appearance
 Interpretation: Possible ventricular tachycardia or CPR artifact or other artifact

■ **ECG 14•16**

Rate: 70 bpm; 120 bpm in initial section, then 39 bpm in remaining section
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.10 sec
 Interpretation: Sinus tachycardia changing to sinus bradycardia

■ **ECG 14•17**

Rate: Atrial 75 bpm, ventricular 40 bpm
 Rhythm: Atrial regular, ventricular irregular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.12 sec
 Interpretation: Second-degree AV block Type II

■ **ECG 14•18**

Rate: 20 bpm
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: Wide—greater than 0.10 sec
 Interpretation: Idioventricular rhythm (agonal rhythm)

■ **ECG 14•19**

Rate: 110 bpm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.08 sec
 Interpretation: Sinus tachycardia with 0.92 second pause (sinoatrial block)

■ **ECG 14•20**

Rate: 79 bpm
 Rhythm: Regular
 P Waves: Normal
 PR Interval: 0.20 sec
 QRS: 0.08 sec
 Interpretation: Normal sinus rhythm with ST segment elevation

■ **ECG 14•21**

Rate: Atrial 75 bpm, ventricular 48 bpm
 Rhythm: Atrial regular, ventricular regular
 P Waves: Normal, some obscured by QRS complexes
 PR Interval: Variable, not associated with QRS complexes
 QRS: 0.20 sec
 Interpretation: Third-degree AV block

■ **ECG 14•22**

Rate: 75 bpm
 Rhythm: Regular
 P Waves: Normal following pacemaker spike
 PR Interval: 0.20 sec
 QRS: 0.20 sec following pacemaker spike
 Interpretation: Pacemaker—atrial and ventricular

■ **ECG 14•23**

Rate: 50 bpm
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: 0.12 sec
 Interpretation: Atrial fibrillation with bundle branch block and slow ventricular response

■ **ECG 14•24**

Rate: Indeterminate
 Rhythm: Irregular
 P Waves: Present, low amplitude in first two beats only
 PR Interval: 0.20 sec in first two beats only
 QRS: Wide—0.16 sec in first two beats
 Interpretation: Sinus bradycardia with a bundle branch block deteriorating into ventricular fibrillation

■ **ECG 14•25**

Rate: 214 bpm
 Rhythm: Regular
 P Waves: None visible
 PR Interval: None
 QRS: 0.06 sec
 Interpretation: Supraventricular tachycardia with muscle artifact

■ **ECG 14•26**

Rate: 80 bpm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.10 sec in normal beats, 0.16 sec in wide beats
 Interpretation: Normal sinus rhythm with bigeminal multiform PVCs

■ **ECG 14•27**

Rate: 80 bpm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.12 sec
 QRS: 0.10 sec
 Interpretation: Normal sinus rhythm with PJC at beat 2, and uniform PVCs at beats 4 and 7

■ **ECG 14•28**

Rate: 110 bpm; 80 bpm in paced portion
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: 0.20 sec
 Interpretation: Pacemaker—ventricular, with PVC at beat 3 and the rhythm deteriorates into monomorphic ventricular tachycardia

■ **ECG 14•29**

Rate: 68 bpm after defibrillation
 Rhythm: Irregular
 P Waves: Normal in last three beats
 PR Interval: 0.16 sec in last three beats
 QRS: 0.08 sec in last three beats
 Interpretation: Ventricular fibrillation defibrillated and converting to normal sinus rhythm

■ **ECG 14•30**

Rate: 79 bpm
 Rhythm: Regular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.08 sec
 Interpretation: Normal sinus rhythm with ST segment depression

■ **ECG 14•31**

Rate: 130 bpm
 Rhythm: Irregular
 P Waves: Different forms
 PR Interval: Variable
 QRS: 0.06 sec
 Interpretation: Multifocal atrial tachycardia

■ **ECG 14•32**

Rate: 75 bpm
 Rhythm: Regular
 P Waves: Normal
 PR Interval: 0.12 sec
 QRS: 0.12 sec with notched appearance
 Interpretation: Normal sinus rhythm with a bundle branch block

■ **ECG 14•33**

Rate: Indeterminate
 Rhythm: Chaotic
 P Waves: None
 PR Interval: None
 QRS: None
 Interpretation: Ventricular fibrillation—coarse fibrillatory waves transitioning to fine fibrillatory waves

■ **ECG 14•34**

Rate: 50 bpm
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: Wide—greater than 0.10 sec following pacemaker spike
 Interpretation: Pacemaker—ventricular, with oversensing of muscle artifact after beats 2 and 4

■ **ECG 14•35**

Rate: 80 bpm (counting PVCs), 75 bpm in underlying rhythm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.10 sec in normal beats, 0.16 sec in wide beats
 Interpretation: Normal sinus rhythm with ventricular quadrigeminy

■ **ECG 14•36**

Rate: 50 bpm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.08 sec
 Interpretation: Sinus bradycardia with sinus arrhythmia with inverted T waves

■ **ECG 14•37**

Rate: 144 bpm
 Rhythm: Irregular
 P Waves: Inverted at beat 7
 PR Interval: 0.16 sec
 QRS: Wide—greater than 0.10 sec except for beat 7, which is 0.10 sec
 Interpretation: Monomorphic ventricular tachycardia with one supraventricular complex at beat 7

■ **ECG 14•38**

Rate: 30 bpm
 Rhythm: Irregular
 P Waves: Flutter waves
 PR Interval: None
 QRS: 0.08 sec
 Interpretation: Atrial flutter with variable block and slow ventricular response

■ **ECG 14•39**

Rate: 94 bpm
 Rhythm: Regular
 P Waves: Normal
 PR Interval: 0.20 sec
 QRS: 0.08 sec
 Interpretation: Normal sinus rhythm

■ **ECG 14•40**

Rate: 68 bpm
 Rhythm: Regular
 P Waves: Inverted
 PR Interval: 0.12 sec
 QRS: 0.04 sec
 Interpretation: Accelerated junctional rhythm with ST segment depression and flattened T waves

■ **ECG 14•41**

Rate: 130 bpm
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: 0.08 sec
 Interpretation: Atrial fibrillation with rapid ventricular response

■ **ECG 14•42**

Rate: 60 bpm; 38 bpm in first section, 98 bpm in second section
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.10 sec
 Interpretation: Sinus bradycardia changing to normal sinus rhythm

■ **ECG 14•43**

Rate: 41 bpm
 Rhythm: Regular
 P Waves: Normal
 PR Interval: 0.20 sec
 QRS: 0.12 sec
 Interpretation: Sinus bradycardia with bundle branch block

■ **ECG 14•44**

Rate: 115 bpm
 Rhythm: Regular
 P Waves: Normal
 PR Interval: 0.12 sec
 QRS: 0.06 sec
 Interpretation: Sinus tachycardia with peaked T waves

■ **ECG 14•45**

Rate: 79 bpm
 Rhythm: Regular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.12 sec with notched appearance
 Interpretation: Normal sinus rhythm with a bundle branch block

■ **ECG 14•46**

Rate: 70 bpm; 75 bpm in underlying rhythm
 Rhythm: Irregular, regular starting with second beat
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.10 sec
 Interpretation: Normal sinus rhythm with a 1.52 second pause (sinoatrial block)

■ **ECG 14•47**

Rate: 100 bpm; 75 bpm in underlying rhythm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.10 sec
 Interpretation: Normal sinus rhythm with two runs of three-beat ventricular tachycardia (sets of triplet PVCs)

■ **ECG 14•48**

Rate: 100 bpm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.10 sec in normal beats, 0.16 sec in wide beats
 Interpretation: Normal sinus rhythm with ST segment depression and bigeminal uniform PVCs

■ **ECG 14•49**

Rate: 80 bpm
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: 0.12 sec
 Interpretation: Atrial fibrillation with bundle branch block and controlled ventricular response

■ **ECG 14•50**

Rate: 70 bpm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.08 sec
 Interpretation: Sinus arrhythmia

■ **ECG 14•51**

Rate: 70 bpm
 Rhythm: Irregular
 P Waves: Normal following pacemaker spike
 PR Interval: 0.20 sec
 QRS: 0.08 sec
 Interpretation: Pacemaker—atrial paced and ventricular sensed, with ST segment depression and PVCs at beats 4 and 6

■ **ECG 14•52**

Rate: 170 bpm; 110 bpm in first section, 240 bpm in last section
 Rhythm: Irregular
 P Waves: Normal for beats 1 through 8; buried in T wave in beats 9 through 17
 PR Interval: 0.22 sec in first section; thereafter not measurable
 QRS: 0.08 sec
 Interpretation: Paroxysmal supraventricular tachycardia (normal sinus rhythm changing to a supraventricular tachycardia)

■ **ECG 14•53**

Rate: 80 bpm
 Rhythm: Irregular
 P Waves: Normal, some following pacemaker spike
 PR Interval: 0.20 sec in paced beats, 0.24 sec in beats 3 and 6, 0.16 sec in beat 4
 QRS: 0.06 sec in narrow complexes, 0.12 sec in wide complexes
 Interpretation: Pacemaker—atrial paced and ventricular sensed in beats 1, 7, and 8; intrinsic sinus complexes at beats 3, 4, and 6 with T wave inversion; PVCs at beats 2 and 5

■ **ECG 14•54**

Rate: 107 bpm
 Rhythm: Regular
 P Waves: Flutter waves
 PR Interval: None
 QRS: 0.08 sec
 Interpretation: Atrial flutter with 3:1 conduction

■ **ECG 14•55**

Rate: 83 bpm
 Rhythm: Regular
 P Waves: Normal
 PR Interval: 0.20 sec
 QRS: 0.10 sec
 Interpretation: Normal sinus rhythm

■ **ECG 14•56**

Rate: 47 bpm
 Rhythm: Regular
 P Waves: Inverted
 PR Interval: 0.10 sec
 QRS: 0.10 sec
 Interpretation: Junctional rhythm

■ **ECG 14•57**

Rate: 50 bpm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.12 sec
 QRS: 0.08 sec
 Interpretation: Sinus bradycardia with sinus arrhythmia

■ **ECG 14•58**

Rate: 100 bpm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.08 sec
 Interpretation: Normal sinus rhythm with bigeminal PJs

■ **ECG 14•59**

Rate: 50 bpm (counting PVCs), 28 bpm in underlying rhythm
 Rhythm: Irregular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.10 sec
 Interpretation: Sinus bradycardia with ST elevation, T wave inversion, and interpolated PVCs (R on T) at beats 3 and 5

■ **ECG 14•60**

Rate: 65 bpm
 Rhythm: Regular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.08 sec
 Interpretation: Normal sinus rhythm with ST segment elevation

■ **ECG 14•61**

Rate: 70 bpm; 60 bpm in underlying paced rhythm
 Rhythm: Irregular
 P Waves: Normal following pacemaker spike
 PR Interval: 0.20 sec
 QRS: 0.10 sec
 Interpretation: Pacemaker—atrial paced, ventricular sensed, with PJs at beats 2 and 6

■ **ECG 14•62**

Rate: Indeterminate
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: Wide—greater than 0.10 sec
 Interpretation: Sinus pause (sinus arrest) deteriorating into ventricular tachycardia

■ **ECG 14•63**

Rate: 46 bpm
 Rhythm: Regular
 P Waves: None
 PR Interval: None
 QRS: 0.08 sec
 Interpretation: Junctional rhythm with ST segment depression

■ **ECG 14•64**

Rate: Atrial 65 bpm, ventricular 40 bpm (28 bpm in first three complexes)
 Rhythm: Atrial regular, ventricular irregular
 P Waves: Normal
 PR Interval: Variable, not associated with QRS complexes
 QRS: 0.18 sec
 Interpretation: Third-degree AV block

■ **ECG 14•65**

Rate: None; atrial 68 bpm
 Rhythm: None
 P Waves: Present and regular
 PR Interval: None
 QRS: None
 Interpretation: Ventricular asystole

■ **ECG 14•66**

Rate: Indeterminate
 Rhythm: Chaotic
 P Waves: None
 PR Interval: None
 QRS: None
 Interpretation: Ventricular fibrillation—coarse changing to fine fibrillatory waves

■ **ECG 14•67**

Rate: 40 bpm
 Rhythm: Regular
 P Waves: Normal
 PR Interval: 0.52 sec
 QRS: 0.10 sec
 Interpretation: Sinus bradycardia with first-degree AV block

■ **ECG 14•68**

Rate: Atrial 75 bpm, ventricular 48 bpm
 Rhythm: Atrial regular, ventricular regular
 P Waves: Variable (some upright, buried in QRS or in T wave)
 PR Interval: Variable, not associated with QRS complexes
 QRS: 0.12 sec
 Interpretation: Third-degree AV block

■ **ECG 14•69**

Rate: 130 bpm; 110 bpm in underlying rhythm
 Rhythm: Irregular
 P Waves: Normal in first three beats and in beats 9 to 13; different in beat 4 and not visible in beats 5 to 8
 PR Interval: 0.12 sec in first three beats and in beats 9 to 13; 0.08 sec in beat 4
 QRS: 0.08 sec
 Interpretation: Sinus tachycardia with ST segment depression and inverted T waves with a PAC in beat 4 and a short run of supraventricular tachycardia in beats 5 to 8

■ **ECG 14•70**

Rate: 50 bpm
 Rhythm: Regular
 P Waves: Normal
 PR Interval: 0.16 sec
 QRS: 0.12 sec
 Interpretation: Second-degree AV block type II with 3:1 block and a bundle branch block

■ **ECG 14•71**

Rate: Indeterminate
 Rhythm: Chaotic
 P Waves: None
 PR Interval: None
 QRS: None
 Interpretation: Ventricular fibrillation—fine fibrillatory waves

■ **ECG 14•72**

Rate: 63 bpm
 Rhythm: Regular
 P Waves: Normal
 PR Interval: 0.20 sec
 QRS: 0.16 sec with notched appearance
 Interpretation: Normal sinus rhythm with a bundle branch block

■ **ECG 14•73**

Rate: 100 bpm
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: 0.16 sec following pacemaker spike and in intrinsic beats
 Interpretation: Pacemaker—ventricular, with a fusion beat in beat 3 and wide intrinsic complexes in beats 4 and 5

■ **ECG 14•74**

Rate: 80 bpm
 Rhythm: Irregular
 P Waves: Normal except for PJC's
 PR Interval: 0.16 sec
 QRS: 0.10 sec
 Interpretation: Normal sinus rhythm with ST segment depression, inverted T waves, and bigeminal PJC's

■ **ECG 14•75**

Rate: 20 bpm
 Rhythm: Irregular
 P Waves: None
 PR Interval: None
 QRS: Wide—greater than 0.10 sec
 Interpretation: Idioventricular rhythm (agonal rhythm)

Case Studies



ECG Case Studies One

To illustrate medical standards, the following case studies are based on the medical protocols (guidelines and medications) presented in Appendices A through D. Choose the most appropriate answer to each question according to the case scenario. All of the ECG strips were recorded in lead II.




CASE STUDY ONE

A 70-year-old woman was admitted to the hospital 2 days ago and underwent a right total hip replacement. The patient has a history of hypertension, degenerative joint disease, arthritis, pulmonary embolism, and coronary artery disease. She just finished a physical therapy session and has

been resting in her hospital room. When you walk into the room to check your patient, you notice that she is unresponsive with no respiration or pulse. You immediately call a code and begin cardiopulmonary resuscitation (CPR). When the emergency team arrives, the patient is attached to the ECG monitor, which displays the rhythm shown in ECG 15-1.

ECG 15-1




- The interpretation of ECG 15-1 is
 - Ventricular fibrillation.
 - Ventricular tachycardia.
 - Accelerated idioventricular.
 - Asystole.
- Your initial treatment of the patient is to
 - Administer 300 mg of amiodarone.
 - Defibrillate at 360 J (or equivalent biphasic energy).
 - Consider 1 mg of atropine.
 - Administer synchronized cardioversion at 360 J.
- Following the appropriate treatment in question 2, ECG 15-2 now shows on the monitor as
 - Ventricular tachycardia.
 - Atrial tachycardia.
 - Ventricular fibrillation.
 - Atrial fibrillation.

ECG 15•2




4. Your next appropriate treatment of choice for this patient is to
- Provide five cycles (2 min) of CPR.
 - Defibrillate at 360 J (or equivalent biphasic energy).
 - Consider 1–2 g of magnesium sulfate.
 - Defibrillate at 200 J (or equivalent biphasic energy).
5. Following the appropriate treatment in question 4, ECG 15–3 shows on the monitor as
- Torsade de pointes.
 - Asystole.
 - Ventricular fibrillation.
 - Ventricular tachycardia.

ECG 15•3



6. Your next appropriate treatment for this patient is to
- Defibrillate at 360 J (or equivalent biphasic energy).
 - Administer epinephrine 1 mg (10 mL of 1:1000).
 - Administer vasopressin 40 U.
 - Either A or C.
7. Following the appropriate treatment in question 6, ECG 15–4 shows on the monitor as
- Atrial fibrillation.
 - Ventricular fibrillation.
 - Torsade de pointes.
 - Ventricular tachycardia.

ECG 15•4



8. The patient is still pulseless with no respiration and has not responded to any treatments given so far. Now your most appropriate treatment choice is to administer

- A. Epinephrine 1 mg (10 mL of 1:10,000).
- B. Vasopressin 40 U.
- C. Epinephrine 1 mg (10 mL of 1:1000).
- D. Either A or B.

9. Following the appropriate treatment in question 8, there is no pulse. After 2 minutes of CPR you defibrillate at 360 J, the rhythm (ECG 15-5) shows on the monitor as

- A. Ventricular fibrillation converting to a sinus rhythm.
- B. Ventricular tachycardia converting to asystole.
- C. Torsade de pointes.
- D. Atrial tachycardia converting to a sinus rhythm.


ECG 15•5



10. The patient is still not responsive but now has spontaneous respirations of 12 breaths per minute, a palpable carotid pulse, and a blood pressure (BP) of 100/70 mm Hg. Please identify the last rhythm in this case study that shows on the monitor (ECG 15-6):

- A. Sinus rhythm at 75 bpm.
- B. Atrial fibrillation at 70 bpm.
- C. Junctional tachycardia at 80 bpm.
- D. Sinus arrhythmia at 90 bpm.

ECG 15•6



CASE STUDY TWO

You are at a public beach and suddenly notice an 8-year-old boy floating face down close to shore. At first you wonder if he is breathing through a snorkel, but you see no sign of either a snorkel or a mask. You become concerned for his safety because he is not lifting his head to breathe. You swim out about 10 feet and bring the boy back to the beach. You lay him on his back. There is no apparent trauma. You later find out that the boy, who is not a good swimmer, had ventured out dangerously far from shore.

1. What is your first plan of action?

- A. Turn the child on his side to expel any water he may have swallowed.
- B. Hit the child firmly between the shoulder blades to expel any water he may have swallowed.
- C. Check for responsiveness.
- D. Begin CPR.

2. After you perform the proper step in question 1, the boy is still not moving. The boy's parents are not around, but you see several bystanders. You

- A. Send a bystander to summon help, phone 911, and get an automated external defibrillator (AED) if one is available.
 - B. Wait until the parents arrive before you do anything else.
 - C. Wait for the lifeguard, who is 10 minutes away.
 - D. Begin rescue breathing.
3. You have already placed the boy supine on a flat surface and found him to be unresponsive. There are no signs of trauma or spinal injury. Your next step is to
 - A. Open the airway by using the jaw thrust method.
 - B. Begin rescue breathing.
 - C. Check for a pulse.
 - D. Open the airway by using the head tilt–chin lift method.
 4. Following the correct procedure in question 3, your next step is to
 - A. Begin chest compressions.
 - B. Begin rescue breathing.
 - C. Look, listen, and feel for adequate breathing.
 - D. Check for a pulse.
 5. Following the correct procedure in question 4, you find that the boy is not breathing. You
 - A. Give one breath.
 - B. Give two breaths.
 - C. Check for a pulse.
 - D. Begin chest compressions.
 6. If the procedure in question 5 is successful, your next step is to
 - A. Check for a carotid pulse.
 - B. Check for a pedal pulse.
 - C. Check for a radial pulse.
 - D. Check for a brachial pulse.
 7. After checking for a pulse you find none. You then begin chest compressions at a rate of
 - A. 80 per minute.
 - B. 90 per minute.
 - C. 100 per minute.
 - D. 120 per minute.
 8. You compress the chest at
 - A. 1.0–2.0 in.
 - B. 1.5–2.0 in.
 - C. One third the depth of the chest.
 - D. One third to one half the depth of the chest.
 9. The proper compression-to-ventilation ratio for a child with one rescuer performing CPR is
 - A. 3:1.
 - B. 15:2.
 - C. 15:1.
 - D. 30:2.
 10. After the fifth cycle of CPR (2 min) you stop to recheck the pulse and look for any other signs of circulation. You find a pulse of 80 bpm but no respiratory rate. You then
 - A. Continue rescue breathing at 10–12 breaths per minute.
 - B. Continue rescue breathing at 12–20 breaths per minute.
 - C. Continue rescue breathing at 24–30 breaths per minute.
 - D. None of the above.
 11. After another 4 minutes the ambulance arrives and the young boy is taken to the closest hospital. If the boy had started breathing at 16 breaths per minute with a pulse of 80 bpm before the ambulance arrived, what would have been your next step?
 - A. Monitor breathing and circulation and place the child in the recovery position.
 - B. Leave for home.
 - C. Walk around and look for the boy's parents.
 - D. Continue CPR.




CASE STUDY THREE

An 82-year-old man arrives in the emergency department complaining of chest pain and palpitations. He has an altered level of conscious-

ness, shortness of breath, and diaphoresis. There are no signs of trauma. He has a history of heart disease, hypertension, and diabetes. When the emergency team arrives, the patient is attached to the ECG monitor, which displays the rhythm shown in ECG 15–7.

ECG 15•7



1. The interpretation of ECG 15–7 is

- A. Paroxysmal supraventricular tachycardia.
- B. Supraventricular tachycardia.
- C. Ventricular tachycardia.
- D. Atrial tachycardia.

2. Your initial assessment of the patient is to

- A. Defibrillate at 360 J (or equivalent biphasic energy).
- B. Administer 300 mg of amiodarone.

- C. Administer five cycles (2 min) of CPR.
- D. Consider and treat possible causes.

3. Following the appropriate assessment in question 2, ECG 15–8 shows on the monitor as

- A. Atrial fibrillation.
- B. Ventricular tachycardia.
- C. Sinus tachycardia.
- D. Supraventricular tachycardia.

ECG 15•8




4. Your appropriate treatment choice for this patient is to

- A. Defibrillate at 360 J (or equivalent biphasic energy).
- B. Administer synchronized cardioversion at 100 J (or equivalent biphasic energy).
- C. Administer 1 mg of epinephrine.
- D. Administer 1 mg of atropine.

5. Following the appropriate treatment in question 4, the rhythm (ECG 15–9) shows on the monitor as


- A. Ventricular fibrillation.
- B. Supraventricular tachycardia.
- C. Ventricular tachycardia.
- D. Paroxysmal supraventricular tachycardia.

ECG 15•9



6. The patient's condition is still unstable. Your next appropriate treatment is to
- Administer synchronized cardioversion at 200 J (or equivalent biphasic energy).
 - Administer 1–2 g of magnesium sulfate.
 - Defibrillate at 360 J (or equivalent biphasic energy).
 - Administer 1 mg of atropine.
7. Following the appropriate treatment in question 6, ECG 15–10 shows on the monitor as
- Ventricular fibrillation.
 - Atrial fibrillation.
 - Ventricular tachycardia.
 - Sinus tachycardia.

ECG 15•10




8. Your next most appropriate treatment choice would be to
- Administer synchronized cardioversion at 300 J (or equivalent biphasic energy).
 - Administer synchronized cardioversion at 100 J (or equivalent biphasic energy).
 - Administer synchronized cardioversion at 50 J (or equivalent biphasic energy).
 - Defibrillate at 360 J (or equivalent biphasic energy).
9. Following the appropriate treatment in question 8, the rhythm (ECG 15–11) shows on the monitor as
- Ventricular tachycardia converting to a sinus rhythm.
 - Ventricular tachycardia converting to agonal rhythm.
 - Ventricular fibrillation converting to junctional rhythm.
 - Paroxysmal supraventricular tachycardia.

ECG 15•11



10. After about 30 minutes, the patient feels calm and has respirations of 18 per minute, a BP of 130/80 mm Hg, and a temperature of 98.6° F. His rhythm (ECG 15–12) shows on the monitor as
- A. Sinus arrhythmia.
 - B. Sinus bradycardia.
 - C. Normal sinus rhythm.
 - D. Atrial flutter.

ECG 15•12




CASE STUDY FOUR

You and your Emergency Medical Services team are called to respond to an unconscious person. A 48-year-old woman has been found unresponsive, sitting in a living room chair. Her husband quickly phoned 911. When you arrive, you

find him correctly administering one-person CPR to his wife. He has recently taken a CPR course at a local hospital. He tells you that his wife had an aortic valve replacement 8 months ago and has a history of hypertension. The woman remains pulseless. Once you attach the ECG monitor, ECG 15–13 shows

ECG 15•13



1. The interpretation of the rhythm in ECG 15-13 is

- A. Junctional tachycardia with pulseless electrical activity.
- B. Atrial tachycardia with pulseless electrical activity.
- C. Sinus tachycardia with pulseless electrical activity.
- D. Accelerated junctional rhythm with pulseless electrical activity.

2. Your initial treatment of the patient is to continue CPR and

- A. Defibrillate at 360 J (or equivalent biphasic energy).

- B. Consider and treat possible causes.
- C. Administer 300 mg of amiodarone.
- D. Administer 6 mg of adenosine.

3. Following the appropriate treatment in question 2, the patient still has no respiration or pulse and the monitor (ECG 15-14) shows

- A. Junctional tachycardia with pulseless electrical activity.
- B. Atrial tachycardia with pulseless electrical activity.
- C. Sinus tachycardia with pulseless electrical activity.
- D. Accelerated junctional rhythm with pulseless electrical activity.

ECG 15•14




4. Your next appropriate treatment of choice is to administer

- A. Epinephrine 1 mg.
- B. Vasopressin 40 U.
- C. Vasopressin 50 U.
- D. Either A or B.

5. Following the appropriate treatment in question 4, the patient is still not breathing but does have a pulse. The monitor (ECG 15-15) shows

- A. Accelerated junctional rhythm.
- B. Atrial tachycardia.
- C. Sinus bradycardia.
- D. Junctional tachycardia.

ECG 15•15



6. Your next appropriate treatment for this patient is to


- A. Administer 0.5 mg of atropine.
- B. Stop ventilating the patient.
- C. Begin chest compressions.
- D. Administer 6 mg of adenosine.

7. Following the correct treatment in question 6, the patient is now breathing on her own at

14 breaths per minute and has a BP of 110/70 mm Hg. The monitor displays the rhythm shown in ECG 15–16. The interpretation of this rhythm is

- A. Sinus tachycardia.
- B. Normal sinus rhythm.
- C. Junctional rhythm.
- D. Sinus bradycardia.

ECG 15•16




CASE STUDY FIVE

You are staffing the medical area for a 5 K marathon in your city. The ambient temperature is 81° F and the relative humidity is 46%. One of the runners, a 32-year-old man, crosses the finish line, walks over to the medical area, and asks for your help. He is sweating profusely and says he just needs to sit down and rest. He complains of some slight

cramping in his legs. He says he drank water along the course. He indicates that he has no past medical problems and is not taking any medication. He has participated in marathons before. He appears to be of average height and weight and says he runs about 10 miles a day. You take his vital signs and find a temperature of 98.8° F, respirations of 20 per minute, and a BP of 118/60 mm Hg. The monitor shows the rhythm in ECG 15–17.

ECG 15•17



1. The interpretation of ECG 15–17 is

- A. Junctional rhythm.
- B. Normal sinus rhythm.
- C. Sinus bradycardia.
- D. Sinus tachycardia.

2. Your initial assessment of the man's vital signs and ECG indicate that they are

- A. Abnormal.
- B. Within normal limits.
- C. Outside normal limits.
- D. None of the above.

3. Your initial treatment of the man would be to

- A. Offer him water or a sports drink.
- B. Let him leave the medical area.

- C. Call for an ambulance.
 - D. Administer 325 mg of aspirin.
4. After drinking fluids and lying down on a cot for about 30 minutes, the man has cooled off and is asking to go home. At this point he has a temperature of 98.5° F, respirations of 12 per

minute, and a BP of 120/70 mm Hg. The rhythm (ECG 15–18) shows on the monitor as

- A. Sinus tachycardia.
- B. Normal sinus rhythm.
- C. Sinus bradycardia.
- D. Sinus arrhythmia.

ECG 15•18



5. If the man refuses to go to the hospital your next appropriate management is to
- A. Let him run the race again.
 - B. Administer 325 mg of aspirin.
 - C. Obtain a signed medical release.
 - D. Call the police and restrain the patient.

if using a biphasic manual defibrillator, use the manufacturer's device-specific energy levels, usually 120–200 J).

7. The correct answer is B.

The rhythm is still chaotic with no organized activity.

8. The correct answer is D.

Administer epinephrine 1 mg (10 mL of 1:10,000) by the intravenous or intraosseous (IV/IO) method; follow with 20 mL IV flush. Repeat every 3–5 minutes; give 2.0–2.5 mg diluted in 10 mL normal saline if administering by endotracheal (ET) tube; or administer a single dose of vasopressin 40 U IV/IO to replace the first or second dose of epinephrine.

9. The correct answer is A.

Your treatment choice of defibrillation converted the chaotic electrical activity into an organized rhythm.

10. The correct answer is A.

Although the patient's ECG is normal and her heart rate has stabilized, she is still not responsive and will need to undergo tests and be medically managed for her health issues.

Answers to

CASE STUDY ONE



1. The correct answer is A.

Chaotic electrical activity occurs with no ventricular depolarization or contraction.

2. The correct answer is B.

Once the defibrillator is available, use it without delay. Defibrillate at a monophasic energy level of 360 J (or, if using a biphasic manual defibrillator, use the manufacturer's device-specific energy levels, usually 120–200 J).

3. The correct answer is C.

The rhythm is still chaotic with no organized activity.

4. The correct answer is A.

Once defibrillation is complete, CPR should resume immediately.

5. The correct answer is C.

The rhythm is still chaotic with no organized activity.

6. The correct answer is A.

Because the patient still has a shockable rhythm, defibrillate at a monophasic energy level of 360 J (or,

Answers to

CASE STUDY TWO



1. The correct answer is C.

Once the child is out of the water, gently tap his shoulder, and ask, "Are you OK?" Checking for responsiveness is the first step in CPR.

2. The correct answer is A.

The bystanders can activate the EMS system while you begin immediate steps for CPR. Because the child is a victim of asphyxial arrest (e.g., drowning) if you had been alone you would have given five cycles (2 min) of CPR before you called for help.

3. The correct answer is D.

Because there is no evidence of trauma or spinal injury, the easiest way to open the airway is by using the head tilt–chin lift method.

4. The correct answer is C.

Once you have opened the airway, it is important to look for the chest to rise, listen for any breath sounds, and feel for any airflow from the child's nose or mouth. You should take no more than 10 seconds to do this.

5. The correct answer is B.

At this point in the CPR algorithm you would give two breaths, allowing 1 second for each breath.

6. The correct answer is A.

In a child (1 yr to adolescent [12–14 yr]) you would check for a carotid pulse because the carotid artery is usually the strongest pulse point. You would check for a brachial pulse in an infant (younger than 1 yr).

7. The correct answer is C.

The compression rate is the speed of the compressions, not the actual number of compressions per minute. The compression rate in a child, if uninterrupted, would be 100 per minute.

8. The correct answer is D.

For a child you would compress at one third to one half the depth of the chest.

9. The correct answer is D.

The correct ratio for a child is 30:2 for one rescuer CPR. If two rescuers were involved the ratio would be 15:2.

10. The correct answer is B.

Your efforts have been successful at reestablishing a pulse. However, the child still needs respiratory support of 12–20 breaths per minute.

11. The correct answer is A.

If the child had regained adequate breathing and circulation, he should have been placed in the recovery position (on his side) and had his

breathing and circulation monitored until help arrived.

*Answers to***CASE STUDY THREE****1. The correct answer is C.**

The patient is showing signs and symptoms of this rapid heart rate.

2. The correct answer is D.

Always check for an underlying illness or condition such as trauma, tension pneumothorax, thrombosis (pulmonary or coronary), tamponade (cardiac), toxins, hypo- or hyperkalemia, hypovolemia, hypoxia, hypoglycemia, hypothermia, or hydrogen ion (acidosis). Correcting or managing one of these issues may help with the treatment.

3. The correct answer is B.

Because this form of VT has a regular rhythm and the complexes are formed similarly, it is called monomorphic ventricular tachycardia.

4. The correct answer is B.

Because the patient is unstable and obviously symptomatic, electrical cardioversion starting at 100 J is the treatment of choice.

5. The correct answer is C.

The patient's condition is still unstable and the ECG shows VT.

6. The correct answer is A.

Since the initial synchronized cardioversion of 100 J had no effect the next synchronized electrical charge would be 200 J.

7. The correct answer is C.

The patient is still unstable and the ECG continues to show ventricular tachycardia.

8. The correct answer is A.

The correct sequence of synchronized electrical cardioversion is 100 J, 200 J, 300 J, and 360 J.

9. The correct answer is A.

The ECG shows that the last cardioversion of 300 J converted the patient's rhythm from VT to a sinus rhythm.

10. The correct answer is C.

The electrical cardioversion finally stabilized the patient's rhythm. He will now undergo tests and be medically managed for his health issues.

*Answers to***CASE STUDY FOUR****1. The correct answer is D.**

The monitor shows an identifiable electrical rhythm, in this case accelerated junctional rhythm; however, no pulse is detectable. Therefore this arrhythmia is described as PEA.

2. The correct answer is B.

Always check for an underlying illness or disease such as trauma, tension pneumothorax, thrombosis (pulmonary or coronary), tamponade (cardiac), toxins, hypo- or hyperkalemia, hypovolemia, hypoxia, hypoglycemia, hypothermia, or hydrogen ion (acidosis). Correcting or managing one of these issues may help with the treatment.

3. The correct answer is A.

The ECG still shows a junctional rhythm with no pulse, but the heart rate is faster. The arrhythmia is still PEA.

4. The correct answer is D.

You can elect to use either epinephrine or vasopressin as your first vasopressor. Remember that if you begin with vasopressin it can be used only as a single dose. After that you must use epinephrine every 3–5 minutes.

5. The correct answer is C.

Sinus bradycardia is a sinus rhythm with a rate of less than 60 bpm.

6. The correct answer is A.

Consider atropine 0.5 mg IV if the heart rate is less than 60 bpm with a pulse.

7. The correct answer is B.

This is a normal sinus rhythm with a rate of 75 bpm. Although the patient's ECG is normal and her heart

rate has returned, she will need to undergo tests and be medically managed for her health issues.

*Answers to***CASE STUDY FIVE****1. The correct answer is B.**

Regular exercise improves the heart's ability to pump blood efficiently. Drinking water during the race has kept the runner from having serious signs and symptoms from dehydration. Therefore, his heart rate has remained normal.

2. The correct answer is B.

All vital signs and the ECG are within normal limits. The runner probably became somewhat dehydrated. In summer sports, it is not the heat, but the combination of heat and humidity, that can cause potential heat-related problems.

3. The correct answer is A.

Water and light sports drinks will help the man avoid potentially dangerous dehydration. They should also be offered along the racecourse and be consumed regularly.

4. The correct answer is C.

The man has a strong pulse with good vital signs, and sinus bradycardia shows on the monitor. The man states that his heart rate normally ranges between 50 and 58 bpm.

5. The correct answer is C.

In most cases, bradycardia in healthy, well-trained athletes does not need to be treated. Once the patient has rested and is feeling back to normal, he does not need further care. It is, however, imperative to have him sign a medical release before he leaves the medical area.

ECG Case Studies Two

To illustrate medical standards, the following case studies are based on the medical protocols (guidelines and medications) presented in Appendices A through D. Choose the most appropriate answer to each question according to the case scenario. All of the ECG strips were recorded in lead II.



CASE STUDY ONE

As a healthcare provider you are attending a workshop, required of all employees at your hospital, concerning workplace safety issues. During a short food break you notice that one of your coworkers, a 55-year-old man, is coughing. As you walk toward your coworker you notice that he is unable to speak and is making high-pitched crowing sounds. He has a panicked look on his face and is grabbing his throat with both hands.

1. What is your first plan of action?

- A. Find the instructor in charge of the workshop.
- B. Hit the man firmly between the shoulder blades.
- C. Ask, "Are you choking? Can you speak?"
- D. Perform a finger sweep.

2. After performing the correct skill in question 1 you next

- A. Perform a finger sweep.
- B. Stand behind the man, wrapping your hands around his waist, and firmly perform abdominal thrusts.
- C. Strike him between the shoulder blades five times.
- D. Ask him to try to cough up the obstructed object.

3. Following the correct procedure in question 2, the airway still is obstructed and

your coworker suddenly slumps slowly to the floor. You first

- A. Establish unresponsiveness.
- B. Find the instructor in charge of the workshop.
- C. Perform a finger sweep.
- D. Check for a carotid pulse.

4. After you have performed the correct procedure in question 3, you find that the man has lost consciousness. Your next step is to

- A. Open the airway, then look, listen, and feel for breathing.
- B. Check for a pulse and then open the airway.
- C. Begin chest compressions.
- D. Begin rescue breathing.

5. After you open the man's airway you notice a chunk of cookie at the back of his pharynx. You then

- A. Begin rescue breathing.
- B. Begin chest compressions.
- C. Check for a pulse.
- D. Perform a finger sweep.

6. After you successfully complete the correct step in question 5, the man coughs, begins breathing normally, and is responsive. You

- A. Let him leave the facility and go home.
- B. Have him drink a glass of water.
- C. Call for help and ask him to stay calm.
- D. Ask him if he wants another cookie.



CASE STUDY TWO

A 65-year-old man arrives in the Emergency Department with heart palpitations. His BP is 148/84 mm Hg, his respirations are 18 per minute, his temperature is 98.6° F, and his heart rate is fast (ECG 16–1). He says he feels as though his heart were racing. He had a similar episode during the past year and

says that, by using the Valsalva maneuver (bearing down), he was able to slow his heart rate on his own. However, that maneuver has not worked this time.

He is not taking any medications and has no prior medical history other than degenerative joint disease requiring a right total knee replacement 2 years ago. Once the emergency team arrives, the patient is attached to the ECG monitor (see ECG 16–1).

ECG 16•1




- The interpretation of ECG 16–1 is
 - Paroxysmal supraventricular tachycardia.
 - Supraventricular tachycardia.
 - Ventricular tachycardia.
 - Atrial tachycardia.
- Your initial treatment for this patient is to
 - Defibrillate at 360 J (or equivalent biphasic energy).
 - Administer 300 mg of amiodarone.
 - Administer 5 cycles (2 min) of CPR.
 - Supply oxygen, start an IV, and obtain a 12-lead ECG.
- Following your treatment in question 2, you
 - Consider and treat possible causes of the arrhythmia.
 - Defibrillate at 360 J (or equivalent biphasic energy).
 - Administer 300 mg of amiodarone.
 - Administer 5 cycles (2 min) of CPR.
- Following the appropriate assessment in question 3, the rhythm (ECG 16–2) shows on the monitor as
 - Atrial fibrillation.
 - Ventricular tachycardia.
 - Sinus tachycardia.
 - Supraventricular tachycardia.

ECG 16•2




5. The patient's vital signs are still stable except for the fast heart rate. Your next appropriate treatment choice for this patient is to
- Defibrillate at 360 J (or equivalent biphasic energy).
 - Administer 6.0 mg of adenosine.
 - Administer 1.0 mg of epinephrine.
 - Administer 1.0 mg of atropine.
6. Following the appropriate treatment in question 5, the rhythm (ECG 16-3) shows on the monitor as
- Ventricular fibrillation.
 - Supraventricular tachycardia.
 - Ventricular tachycardia.
 - Paroxysmal supraventricular tachycardia.

ECG 16•3



7. The patient is still responsive and stable, but after 1–2 minutes the rhythm in ECG 16-3 still shows on the monitor. Your next appropriate treatment for this patient is to
- Defibrillate at 360 J (or equivalent biphasic energy).
 - Administer 1–2 g of magnesium sulfate.
 - Administer 12 mg of adenosine.
 - Administer 1 mg of atropine.
8. Following the appropriate treatment in question 7, the rhythm (ECG 16-4) shows on the monitor as
- Ventricular fibrillation.
 - Atrial fibrillation.
 - Supraventricular tachycardia.
 - Sinus tachycardia.

ECG 16•4




9. After another 2 minutes the rhythm in ECG 16-4 still shows up on the monitor. Now your most appropriate treatment choice is to
- Administer 12 mg of adenosine.
 - Administer 6 mg of adenosine.
 - Defibrillate at 200 J (or equivalent biphasic energy).
 - Defibrillate at 360 J (or equivalent biphasic energy).
10. Following the appropriate treatment in question 9, the rhythm (ECG 16-5) shows on the monitor as
- Atrial fibrillation.
 - Atrial tachycardia.
 - Sinus tachycardia.
 - Conversion to normal sinus rhythm.

ECG 16•5



11. After about 30 minutes the patient feels calm and has a temperature of 98.6°F, respiration of 12 breaths per minute, and a BP of 130/80 mm Hg. His rhythm (ECG 16–6) shows on the monitor as
- A. Sinus arrhythmia.
 - B. Sinus bradycardia.
 - C. Normal sinus rhythm.
 - D. Atrial flutter.

ECG 16•6

**CASE STUDY THREE**

As a paramedic in a rural community you arrive at the scene of an emergency, where an 80-year-old woman has been found lying in an alleyway. You establish that she has no respiration or pulse. No one at the scene knows how long she has

been unresponsive. No trauma is visible and CPR was not initiated before your arrival. The weather is mild, with an ambient temperature of about 70° F. Once your Emergency Medical Services team begins CPR, you attach the ECG monitor and obtain the rhythm (ECG 16–7).

ECG 16•7



1. The interpretation of ECG 16-7 is

- A. Ventricular fibrillation.
- B. Asystole.
- C. Agonal rhythm.
- D. Third-degree block.

2. After initiating CPR you begin treatment by

- A. Administering 6 mg of adenosine.
- B. Considering and treating possible causes for the arrhythmia.
- C. Considering diltiazem or beta blockers.

D. Defibrillating at 360 J (or equivalent biphasic energy).

3. Following the appropriate treatment in question 2, the rhythm (ECG 16-8) shows on the monitor as

- A. Atrial flutter.
- B. Ventricular tachycardia.
- C. Ventricular fibrillation.
- D. Asystole.

ECG 16-8




4. Your next appropriate treatment for this patient is to administer

- A. Amiodarone 300 mg.
- B. Epinephrine 1 mg or vasopressin 40 U.
- C. Magnesium sulfate 1-2 g.
- D. Amiodarone 150 mg.

5. Following the appropriate treatment in question 4, the rhythm (ECG 16-9) shows on the monitor as

- A. Asystole.
- B. Ventricular fibrillation.
- C. Ventricular tachycardia.
- D. Third-degree block.

ECG 16-9



6. As your Emergency Medical Services team continues CPR your next appropriate treatment is to

- A. Administer 12 mg of adenosine.
- B. Defibrillate at 360 J (or equivalent biphasic energy).
- C. Administer 150 mg of amiodarone.
- D. Administer 1.0 mg of atropine.

7. Following the appropriate treatment in question 6, the rhythm (ECG 16-10) shows on the monitor as

- A. Ventricular fibrillation.
- B. Asystole.
- C. Atrial fibrillation.
- D. Agonal rhythm.

ECG 16•10



8. After 30 minutes of CPR and the right medications, the patient is still pulseless with no respiration. The next appropriate treatment choice is to

- A. Continue CPR for 2 more hours.
- B. Defibrillate at 200 J (or equivalent biphasic energy).
- C. Follow local policy to stop resuscitation efforts.
- D. Defibrillate at 360 J (or equivalent biphasic energy).




CASE STUDY FOUR

A 40-year-old female trauma patient is in a monitored room in the Intensive Care Unit of your hospital. You suddenly hear the ECG alarm go off and rush to the patient's room. You find her unre-

sponsive, with no respirations and no pulse. You call a code and begin CPR. Once the code team arrives the patient, who is already attached to the ECG monitor, has the rhythm shown in ECG 16–11.

ECG 16•11



1. The interpretation of ECG 16–11 is


- A. Ventricular fibrillation.
 - B. Agonal rhythm.
 - C. Ventricular tachycardia.
 - D. Asystole.
2. Your initial treatment of the patient is to
- A. Defibrillate at 200 J (or equivalent biphasic energy).
 - B. Defibrillate at 360 J (or equivalent biphasic energy).
 - C. Defibrillate at 50 J (or equivalent biphasic energy).
 - D. Defibrillate at 100 J (or equivalent biphasic energy).
3. Following the appropriate treatment in question 2, ECG 16–12 now shows
- A. Asystole.
 - B. Ventricular fibrillation.
 - C. Agonal rhythm.
 - D. Ventricular tachycardia.

ECG 16•12




4. The patient is still unresponsive, is not breathing, and has no pulse. Your next appropriate treatment is to
- A. Provide 5 cycles (2 min) of CPR.
 - B. Administer 1–3 g of magnesium sulfate.
 - C. Administer 150 mg of amiodarone.
 - D. Stop any further resuscitation efforts.
5. Following the appropriate treatment in question 4, ECG 16–13 shows
- A. Ventricular fibrillation.
 - B. Ventricular tachycardia.
 - C. Asystole.
 - D. Atrial fibrillation.

ECG 16•13



6. The patient is still unresponsive, with no respirations and no pulse. Your next appropriate treatment is to
- A. Perform synchronized cardioversion at 100 J (or equivalent biphasic energy).
 - B. Perform synchronized cardioversion at 200 J (or equivalent biphasic energy).
 - C. Defibrillate at 360 J (or equivalent biphasic energy).
 - D. Stop any further resuscitation.
7. Following the appropriate treatment in question 6, ECG 16–14 shows
- A. Ventricular tachycardia.
 - B. Torsade de pointes.
 - C. Ventricular fibrillation converting to a sinus rhythm.
 - D. Ventricular tachycardia converting to a sinus rhythm.

ECG 16•14



8. In about 15 minutes the patient begins breathing on her own at 12 breaths per minute. Her BP is 100/80 mm Hg and the rhythm in ECG 16–15 shows on the monitor. What is this rhythm?
- Sinus rhythm at 60 bpm.
 - Sinus rhythm with PVCs at 70 bpm.
 - Sinus rhythm at 75 bpm.
 - Sinus rhythm with PVCs at 50 bpm.

ECG 16•15

**CASE STUDY FIVE**

A 70-year-old woman arrives in the Emergency Department with a sudden onset of weakness, fatigue, and chest pain. She has a medical

history of diabetes, hypertension, and osteoporosis. Her vital signs are: temperature 98.6°F, respiration 16 breaths per minute, BP 150/90 mm Hg, and heart rate 75 bpm. Once the patient is attached to the ECG monitor, the rhythm in ECG 16–16 is seen.


ECG 16•16



1. The interpretation of ECG 16–16 is
 - A. Normal sinus rhythm.
 - B. Sinus rhythm with depressed ST segment.
 - C. Sinus rhythm with ST segment elevation.
 - D. Junctional rhythm with ST segment elevation.
2. Your initial management of the patient is to
 - A. Administer atropine 0.5 mg.
 - B. Supply oxygen, start an IV, and obtain a 12-lead ECG.
 - C. Begin CPR.
 - D. Perform cardioversion at 100 J.

3. Your initial treatment of the patient would be to
 - A. Perform cardioversion at 50 J.
 - B. Begin CPR.
 - C. Administer epinephrine 1 mg.
 - D. Administer aspirin 160–325 mg.
4. Following the appropriate treatment in question 3, the rhythm (ECG 16–17) shows as
 - A. Sinus rhythm with ST segment elevation.
 - B. Normal sinus rhythm.
 - C. Sinus rhythm with depressed ST segment.
 - D. Junctional rhythm with ST segment elevation.


ECG 16•17



5. Your next treatment of choice for this patient is to administer
 - A. Nitroglycerin by sublingual route 0.3–0.4 mg.
 - B. Nitroglycerin aerosol spray for 0.5–1.0 sec.
 - C. Aspirin 500 mg.
 - D. Either A or B.

6. Following the appropriate treatment in question 4, the rhythm (ECG 16–18) shows as
 - A. Normal sinus rhythm.
 - B. Sinus rhythm with depressed ST segment.
 - C. Sinus rhythm with ST segment elevation.
 - D. Junctional rhythm with ST segment elevation.


ECG 16•18



7. Your next appropriate treatment is to
 - A. Administer epinephrine 1 mg.
 - B. Begin the checklist for fibrinolytic therapy.
 - C. Let the patient rest for an hour.
 - D. Perform cardioversion at 100 J.

8. The patient is still complaining of chest pain and her BP is 110/60. Following the appropriate treatment in question 7, ECG 16–19 shows
 - A. Sinus rhythm with ST segment elevation.
 - B. Normal sinus rhythm.
 - C. Sinus rhythm with depressed ST segment.
 - D. Junctional rhythm with ST segment elevation.

ECG 16•19



9. Your next most appropriate treatment choice would be to
- Administer aspirin 160–325 mg.
 - Repeat nitroglycerin to a maximum of three doses.
 - Administer atropine 1 mg.
 - Administer epinephrine 1 mg.
10. If the chest pain is still not relieved and the patient's systolic BP is above 90 mm Hg, your next treatment choice is to administer
- Aspirin 160–325 mg.
 - Atropine 1 mg.
 - Epinephrine 1 mg.
 - Morphine 2–4 mg.



CASE STUDY SIX

You are monitoring the medical treatment of a 58-year-old man in the Coronary Care Unit at your hospital. Three days ago the patient underwent a triple coronary artery bypass graft. He has a history of coronary artery disease, congestive


heart failure, hypertension, and diabetes. You are at the nursing station and hear the patient's ECG monitor alarm go off. You rush to his bedside. He is conscious, but his mental status is altered and his BP is 80/50 mm Hg. You check the rhythm on his ECG monitor (ECG 16–20).

ECG 16•20



- The interpretation of the rhythm in ECG 16–20 is
 - Normal sinus rhythm.
 - Sinus bradycardia.
 - Third-degree AV block.
 - Sinus pause (sinus arrest).
- Your initial management of the patient is to
 - Administer epinephrine 1 mg.
 - Supply oxygen, start an IV, and obtain a 12-lead ECG.
 - Begin CPR.
 - Defibrillate at 360 J (or equivalent biphasic energy).
- Your first drug treatment of choice would be to administer
 - Epinephrine 5 mg.
 - Atropine 0.5 mg.
 - Adenosine 12 mg.
 - Magnesium 1–2 g.
- The patient is still conscious but symptomatic. Following the appropriate treatment in question 3, ECG 16–21 shows
 - Sinus bradycardia.
 - Third-degree AV block.
 - Second-degree AV block, Type II.
 - First-degree AV block.

ECG 16•21



5. The patient's BP is now 90/60 mm Hg and he is still confused. Your last correct treatment in question 3 was given 4 minutes ago. Your next treatment of choice is to administer
- Epinephrine 5 mg.
 - Atropine 0.5 mg.
 - Adenosine 12 mg.
 - Magnesium sulfate 1–2 g.
6. The patient now is alert, oriented, and able to follow commands. His BP is 140/90 mm Hg. Following the appropriate treatment in question 5, ECG 16–22 shows
- Sinus bradycardia.
 - Sinus tachycardia.
 - Sinus rhythm at a rate of 75 bpm.
 - Sinus rhythm at a rate of 90 bpm.

ECG 16•22



Answers to

CASE STUDY ONE



1. The correct answer is C.

When you see a person in respiratory distress displaying the universal sign for choking—one or two hands around the neck—ask, “Are you choking? Can you speak?”

2. The correct answer is B.

Stand behind the person and wrap your hands around his waist. Then make a fist with one hand. Place the thumb side of your fist in the middle of the abdomen just above the navel. Grasp your fist with your other hand. Press your fist abruptly into the abdomen using an upward, inward thrust. You have just performed the Heimlich maneuver.

3. The correct answer is A.

When a person slumps to the floor he may be losing consciousness. Before you begin any other step, establish whether the person is still responsive by gently tapping him and asking, “Are you OK?”

4. The correct answer is A.

Once you have established that the person is unresponsive, you need to open the airway and look, listen, and feel for breathing. Do not take longer than 10 seconds to perform this step.

5. The correct answer is D.

Each time the airway is opened, look for an object in the person's mouth. Only remove material you can see, in this case a piece of cookie, with a finger sweep.

Never perform a finger sweep if you do not see a foreign body in the airway.

6. The correct answer is C.

Once the foreign object is removed and the person is responsive and breathing normally, you should make sure medical help arrives and the patient is evaluated to rule out aspiration or other complications.

Answers to

CASE STUDY TWO



1. The correct answer is A.

The onset of PSVT in this case begins as a sinus rhythm then quickly escalates to a rapid SVT.

2. The correct answer is D.

It is important to give the patient oxygen and begin an IV to establish a route for medication administration. Obtaining a 12-lead ECG will identify any other irregularities in the patient's ECG.

3. The correct answer is A.

Always check for underlying illness or disease such as trauma, tension pneumothorax, thrombosis (pulmonary or coronary), tamponade (cardiac), toxins, hypo- or hyperkalemia, hypovolemia, hypoxia, hypoglycemia, hypothermia, or hydrogen ion (acidosis). Correcting or managing one of these issues may help with the treatment.

4. The correct answer is D.

This arrhythmia has such a fast rate that the P waves are not seen and are usually buried in the T waves.

5. The correct answer is B.

Because the patient is stable and the Valsalva maneuver was not successful, the next treatment choice would be 6 mg of adenosine IV given rapidly over 1–3 seconds followed by a 20-mL bolus of normal saline.

6. The correct answer is B.

The ECG still shows SVT.

7. The correct answer is C.

If within 1–2 minutes the SVT does not convert to a slower rate, administer 12 mg of adenosine IV given rapidly over 1–3 seconds followed by a 20-mL bolus of normal saline.

8. The correct answer is C.

The patient is still responsive but the ECG continues to show SVT.

9. The correct answer is A.

In another 1–2 minutes, if the rhythm has not converted, you may give a third dose of adenosine, 12 mg IV given rapidly over 1–3 seconds, followed by a 20-mL bolus of normal saline. You should not exceed a maximum total dose of 30 mg.

10. The correct answer is D.

The adenosine finally slowed and converted the rhythm to normal sinus rhythm.

11. The correct answer is C.

Although the patient's ECG is normal and his heart rate has stabilized, he will need to undergo tests and be medically managed for his health issues.

Answers to

CASE STUDY THREE



1. The correct answer is B.

The electrical activity in the heart is completely absent.

2. The correct answer is B.

In asystole it is extremely important to check for underlying illness or disease such as trauma, tension pneumothorax, thrombosis (pulmonary or coronary), tamponade (cardiac), toxins, hypo- or hyperkalemia, hypovolemia, hypoxia, hypoglycemia, hypothermia, or hydrogen ion (acidosis). Correcting or managing one of these issues may help with the treatment.

3. The correct answer is D.

There is no change in the ECG.

4. The correct answer is B.

You can elect to use either epinephrine or vasopressin as your first vasopressor. Remember that if you begin with vasopressin it can be used only as a single dose. After that you must use epinephrine every 3–5 minutes.

5. The correct answer is A.

There is still no change in the ECG.

6. The correct answer is D.

Even with CPR and emergency medications, the time is running out for this patient. Because the rhythm still shows asystole, you should give 1 mg of atropine, repeating every 3–5 minutes as needed to a total dose of no more than 3 mg.

7. The correct answer is B.

If CPR had been started by a bystander and 911 called immediately, the outcome for this patient might have

been extremely different. Unfortunately the ECG still shows asystole.

8. The correct answer is C.

All localities have different policies to terminate resuscitation efforts. Make sure you know your local protocols. To follow up on the case, you talk to your medical director the next day. She tells you that the autopsy report showed a ruptured aortic aneurysm. The patient had died suddenly even before you began resuscitative efforts.

Answers to

CASE STUDY FOUR



1. The correct answer is C.

It is important in VT to confirm the presence or absence of pulses because ventricular tachycardia may be perfusing or nonperfusing.

2. The correct answer is B.

Do not delay defibrillation. Defibrillate at a monophasic energy level of 360 J. If using a biphasic manual defibrillator, use the manufacturer's device-specific energy levels, usually 120–200 J.

3. The correct answer is D.

The ventricular rate has decreased.

4. The correct answer is A.

Between each defibrillation, it is important to provide 5 cycles (2 min) of CPR.

5. The correct answer is B.

The ventricular rate has increased.

6. The correct answer is C.

If the rhythm is still shockable, defibrillate at a monophasic energy level of 360 J. If using a biphasic manual defibrillator, use the manufacturer's device-specific energy levels, usually 120–200 J.

7. The correct answer is D.

The last defibrillation converted the patient's rhythm to a sinus rhythm.

8. The correct answer is C.

This is a normal sinus rhythm with a rate of 75 bpm. Although the patient's ECG is normal and her heart rate has stabilized, she will need to undergo tests and be medically managed for her health issues.

Answers to

CASE STUDY FIVE



1. The correct answer is C.

ST segment elevation is significant for myocardial injury and must be managed immediately.

2. The correct answer is B.

Oxygen is essential for the heart to survive, an IV allows immediate administration of medications, and a 12-lead ECG is essential in diagnosing an acute MI.

3. The correct answer is D.

Aspirin is the first choice for a patient with the initial signs of acute coronary syndrome. It must be given within minutes of the onset of signs and symptoms.

4. The correct answer is A.

Although the rate may be a little faster, there is still no change in the ECG.

5. The correct answer is D.

Nitroglycerin can be given by either an aerosol or a sublingual route.

6. The correct answer is C.

There is still no change in the ECG.

7. The correct answer is B.

The key to successful fibrinolysis is to start early.

8. The correct answer is A.

There is still no change in the ECG.

9. The correct answer is B.

Nitroglycerin administration requires a systolic blood pressure greater than 90 mm Hg.

10. The correct answer is D.

Administer morphine 2–4 mg (over 1–5 min) every 5–30 minutes until the chest pain is relieved or the patient's blood pressure falls below 90 mm Hg.

Answers to

CASE STUDY SIX



1. The correct answer is C.

Notice that the P waves are not associated with the QRS complexes.

2. The correct answer is B.

Oxygen is critical, an IV will allow immediate administration of medications, and a 12-lead ECG is essential in diagnosing an acute MI.

3. The correct answer is B.

While awaiting a pacemaker, consider atropine 0.5 mg IV every 3–5 minutes (maximum dose not to exceed 3 mg).

4. The correct answer is C.

Notice that the rhythm has changed from a third-degree AV block to a second-degree AV block Type II.

5. The correct answer is B.

Atropine appears to have stabilized the patient's condition, eliminating the need for a transcutaneous pacemaker.

6. The correct answer is C.

Although the patient's ECG is normal and his heart rate has stabilized, he will need to undergo tests and be medically managed for his health issues.

Healthcare Provider Guidelines for Cardiopulmonary Resuscitation (CPR)

CPR HEALTHCARE PROVIDER SKILL PERFORMANCE

CPR Method	Compression/ Ventilation Ratio	Rate of Compressions (min)	Depth of Compressions	Pulse Check (artery)	Hand Position for Compressions
Adult, 1 rescuer	30:2	100	1.5–2.0 in (or deeper in large patients)	Carotid	Heels of 2 hands over center of chest between nipples
Adult, 2 rescuers	30:2	100	1.5–2.0 in (or deeper in large patients)	Carotid	Heels of 2 hands over center of chest between nipples
Child, 1 rescuer	30:2	100	1/3–1/2 Depth of chest	Carotid Femoral	Heel of 1 or 2 hands over lower half of sternum at nipple line
Child, 2 rescuers	15:2	100	1/3–1/2 Depth of chest	Carotid Femoral	Heel of 1 or 2 hands over lower half of sternum at nipple line
Infant, 1 rescuer	30:2	100	1/3–1/2 Depth of chest	Brachial Femoral	2 Fingers on middle sternum, just below nipple line
Infant, 2 rescuers	15:2	100	1/3–1/2 Depth of chest	Brachial Femoral	Two thumb-encircling hands technique over lower half sternum
Newborn	3:1	120	1/3 Depth of chest	Brachial Femoral	Two thumb-encircling hands technique over lower third of sternum

CPR: Adult (Adolescent [12–14 yr] or older)

1. Ensure that the scene is safe. **Check for unresponsiveness.** Gently tap the person's shoulder. Ask, "Are you OK?"
2. **In a sudden collapse, if there is no response and you are alone, summon help, call a code, or phone 911 and get an automated external defibrillator (AED), if available.** Send a second rescuer, if available, for help.
3. **Position the person supine** on a hard, flat surface.
4. **Open the airway** by the head tilt–chin lift method or, if spinal injury is suspected, use the jaw thrust method, if possible.
5. **Look, listen, and feel for adequate breathing (no more than 10 sec).**
6. If the person is not breathing, **begin rescue breaths.** Using a bag-valve-mask device or face mask, **give 2 breaths (1 sec each)** with sufficient volume to cause the chest to rise. Do not over-

- ventilate. Note: If the chest does not rise, reposition the head, chin, and jaw, and attempt 2 more breaths. **If the chest still does not rise, follow instructions for unconscious adult with an obstructed airway (p 250)**
7. **Assess the carotid pulse and look for other signs of circulation (no more than 10 sec). If signs of circulation are present but the person is still not breathing, give rescue breaths at the rate of 10–12 breaths per minute (1 breath every 5–6 sec).**
 8. If a pulse and signs of circulation are not present, **begin compressions.** Place the heel of one hand over the center of the chest between the nipples; place the heel of your other hand over the first. **Firmly compress the chest 1.5–2.0 in. Give 30 compressions. Compress at a rate of 100 per minute.**
 9. **Continue to give 2 breaths followed by 30 compressions.** After the fifth cycle of 30:2 (2 min), **recheck the pulse** and look for other signs of circulation (no more than 10 sec). If circulation is not present, use the AED. Follow the instructions on how to use an AED on page 266. If an AED is unavailable, continue to give 2 breaths followed by 30 compressions. After each fifth cycle of 30:2 (2 min), recheck the pulse and look for other signs of circulation (no more than 10 sec).
 10. If circulation resumes but breathing does not resume or is inadequate, continue rescue breathing at 10–12 breaths per minute.
 11. If adequate breathing and circulation resume, place the person in the recovery position and monitor until help arrives.
2. If no response send a second rescuer, if available, for help.
 3. If you are alone, begin the steps for CPR.
 4. **Position the child supine** on a hard, flat surface.
 5. **Open the airway** by the head tilt–chin lift method or, if spinal injury is suspected, use the jaw thrust method, if possible.
 6. **Look, listen, and feel for adequate breathing (no more than 10 sec).**
 7. If the child is not breathing, **begin rescue breaths.** Using a bag-valve-mask device or face mask, **give 2 breaths (1 sec each) with sufficient volume to make the chest rise.** Do not over-ventilate. Note: If the chest does not rise, reposition the head, chin, and jaw and attempt 2 more breaths. **If the chest still does not rise, follow instructions for unconscious child with an obstructed airway (p 251).**
 8. **Assess the carotid or femoral pulse and look for other signs of circulation (no more than 10 sec). If signs of circulation are present but the child is still not breathing, give rescue breaths at the rate of 12–20 breaths per minute (1 breath every 3–5 sec).**
 9. **If the pulse and signs of circulation are not present or the heart rate is less than 60 bpm with signs of poor perfusion, begin compressions.** Place the heel of one or two hands over the lower half of the sternum on the nipple line. **Firmly compress chest at 1/3 to 1/2 depth of chest. Give 30 compressions. Compress at a rate of 100 per minute.**
 10. **Continue to give 2 breaths followed by 30 compressions.** After the fifth cycle of 30:2 (2 min), **recheck the pulse** and look for other signs of circulation (no more than 10 sec). If you are still alone and no signs of circulation are present, **summon help, call a code, or phone 911 and get an AED, if available.**
 11. Return to the child. If circulation is still not present, continue CPR until the AED is available. Follow the instructions on how to use an AED on page 266. If an AED is unavailable, continue to give 2 breaths followed by 30 compressions. After each fifth cycle of 30:2 (2 min), recheck the pulse and look for other signs of circulation (no more than 10 sec).
 12. If circulation resumes but breathing does not resume or is inadequate, continue rescue breathing at 12–20 breaths per minute.
 13. If adequate breathing and circulation resume, place the child in the recovery position and monitor until help arrives.



Clinical Tip:

Victims of asphyxial arrest (e.g., drowning, drug overdose, respiratory failure) should receive five cycles (2 min) of CPR before the lone rescuer calls for help (activates the EMS system).



Clinical Tip:

When two rescuers are available, give cycles of 30 compressions and 2 breaths for adult CPR.

CPR: Child (1 yr to Adolescent [12–14 yr])

1. Ensure that the scene is safe. **Check for unresponsiveness.** Gently tap child's shoulder. Ask, "Are you okay?"

**Clinical Tip:**

If you are alone and know a child has had a **sudden collapse** due to heart failure, request immediate help including an AED. Do not delay defibrillation.

**Clinical Tip:**

When two rescuers are available, give cycles of 15 compressions and 2 breaths for child CPR.

CPR: Infant (Younger than 1 yr)

1. Ensure that the scene is safe. **Check for unresponsiveness.** Gently rub the infant's back or chest.
2. If no response send a second rescuer, if available, for help.
3. If you are alone, begin the steps for CPR.
4. **Position the infant supine** on a hard, flat surface.
5. **Open the airway** by the head tilt–chin lift method. If spinal injury is suspected, use the jaw thrust method, if possible.
6. **Look, listen, and feel for adequate breathing (no more than 10 sec).**
7. If the infant is not breathing, **begin rescue breaths.** Using a bag-valve-mask device or face mask, **give 2 breaths (1 sec each) with sufficient volume to make the chest rise.** Do not over-ventilate. Note If the chest does not rise, reposition the head, chin, and jaw and give 2 more breaths. **If the chest still does not rise, follow instructions for unconscious infant with an obstructed airway (p 251).**
8. **Assess the brachial or femoral pulse and look for other signs of circulation (no more than 10 sec).** If signs of circulation are present but infant is still not breathing, **continue rescue breaths at the rate of 12–20 breaths per minute (1 breath every 3–5 sec).**
9. **If the pulse and signs of circulation are not present or the heart rate is less than 60 bpm with signs of poor perfusion, begin compressions.** Place two fingers of one hand on the sternum just below the nipple line. **Firmly compress the chest by 1/3 to 1/2 its depth. Give 30 compressions. Compress at a rate of 100 per minute.**
10. **Continue to give 2 breaths followed by 30 compressions.** After the fifth cycle of 30:2 (2 min), **recheck the pulse** and look for other

signs of circulation (no more than 10 sec). If you are still alone and no signs of circulation are present, **summon help, call a code, or phone 911.**

11. If circulation is still not present, continue CPR. After each fifth cycle of 30:2 (2 min), recheck the pulse and look for other signs of circulation (no more than 10 sec).
12. If circulation resumes but breathing does not resume or is inadequate, continue rescue breathing at 12–20 breaths per minute.
13. If adequate breathing and circulation resume, place the infant in the recovery position and monitor until help arrives.

**Clinical Tip:**

If you are alone and know an infant has had a **sudden collapse** due to heart failure, request immediate help.

**Clinical Tip:**

When two rescuers are available, give cycles of 15 compressions and 2 breaths for infant CPR; using the two thumb-encircling hands technique for compression.

**Clinical Tip:**

There is no evidence to support the use of an AED in infants at this time.

OBSTRUCTED AIRWAY: Conscious Adult or Child (1 yr or older)**Clinical Presentation**

- Grabbing at the throat with one or both hands
 - Inability to speak; high-pitched crowing sounds
 - Wheezing, gagging, ineffective coughing
1. **Determine that the airway is obstructed.** Ask, “Are you choking? Can you speak?”
 2. Let the person know you are going to help.
 3. **Stand behind the choking person and wrap your arms around the person's waist.** For someone who is obese or pregnant, wrap your arms around the chest.
 4. **Make a fist. Place the thumb side of your fist in middle of the person's abdomen, just above the navel.** Locate the middle of the sternum for obese or pregnant persons.



Fig. A.1 ■ Heimlich maneuver for the adult or child.

5. Grasp your fist with your other hand.
6. **Press your fist abruptly into the person's abdomen using an upward, inward thrust** (Fig A.1). Use a straight thrust back for someone who is obese or pregnant.
7. Continue thrusts until the object is dislodged or the person loses consciousness.
8. If the person loses consciousness, treat as an unconscious adult or a child with an obstructed airway (pp 250, 251).

OBSTRUCTED AIRWAY: Conscious Infant (younger than 1 yr)

Clinical Presentation

- Inability to breathe or cry
 - High-pitched crowing sounds
 - Sudden wheezing or noisy breathing
1. **Determine that the airway is obstructed. Notice if air exchange is poor or does not occur.**
 2. Lay the infant down on your forearm, with the chest in your hand and the jaw between your thumb and index finger.
 3. Using your thigh or lap for support, keep the infant's head lower than the body (Fig A.2).
 4. **Give five quick, forceful blows between the shoulder blades** with the heel of your hand.
 5. Turn the infant over to be face up on your other arm. Using your thigh or lap for support,




Fig. A.2 ■ Heimlich maneuver for the infant.

- keep the infant's head lower than the body (see Fig A.2).
6. Place two fingers on the sternum just below the nipple line.
 7. **Give five quick thrusts downward, depressing the chest by 1/3 to 1/2 its depth** each time (see Fig A.2).
 8. **Continue the sequence of five back blows and five chest thrusts until the object is dislodged or the infant loses consciousness.** If the infant loses consciousness, treat as an unconscious infant with an obstructed airway (p 251).

OBSTRUCTED AIRWAY: Unconscious Adult (adolescent [12–14 yr] or older)

Clinical Presentation

- Failure to breathe, cyanosis
 - Inability to move air into lungs with rescue breaths
1. **Establish unresponsiveness.** Gently tap person's shoulder. Ask, "Are you OK?"
 2. **If there is no response and you are alone, summon help, call a code, or phone 911 and get an AED, if available.** Send a second rescuer, if available, for help.
 3. **Position the person supine** on a hard, flat surface.
 4. **Open the airway** by the head tilt–chin lift method or, if spinal injury is suspected, use the jaw thrust method, if possible.
 5. **Look, listen, and feel for adequate breathing (no more than 10 sec).**
 6. If the person is not breathing, **begin rescue breaths.** Using a bag-valve-mask device or face mask, **give 2 breaths (1 sec each).** Check to see if the chest rises.
 7. **If the chest does not rise, reposition the head, chin, and jaw, and give 2 more breaths.** If breaths cannot be delivered, perform Adult CPR (p 247), beginning with compressions

and using a 30:2 ratio at a rate of 100/min. Each time the airway is opened, look for an object in the person's mouth. **Only use a finger sweep to remove solid material you can see obstructing the airway. Never perform a blind finger sweep if you do not see a foreign body in the airway.**

8. Continue CPR until breathing and circulation resume or until advanced life support measures are initiated.
9. If circulation resumes but breathing does not resume or is inadequate, continue rescue breathing at 10–12 breaths per minute.
10. If adequate breathing and circulation resume, place the person in the recovery position and monitor until help arrives.



Clinical Tip:

An airway obstruction is successfully relieved if you see and remove the object or feel air movement and see the chest rise when you give breaths.

object in the child's mouth. **Only use a finger sweep to remove material you see obstructing the airway. Never perform a blind finger sweep if you do not see a foreign body in the airway.**

9. Continue CPR until breathing and circulation resume or until advanced life support measures are initiated.
10. After the fifth cycle, if you are still alone, **summon help, call a code, or phone 911 and get an AED, if available.**
11. If circulation resumes but breathing does not resume or is inadequate, continue rescue breathing at 12–20 breaths per minute.
12. If adequate breathing and circulation resume, place the child in the recovery position and monitor until help arrives.



Clinical Tip:

Never perform a blind finger sweep.

OBSTRUCTED AIRWAY: Unconscious Child (1 yr to Adolescent [12–14 yr])

Clinical Presentation

- Failure to breathe, cyanosis
 - Inability to move air into lungs with rescue breaths
1. **Establish unresponsiveness.** Gently tap the child's shoulder. Ask, "Are you OK?"
 2. If there is no response, send a second rescuer, if available, for help.
 3. If you are alone, begin the steps for CPR.
 4. **Position the child supine** on a hard, flat surface.
 5. **Open the airway** by the head tilt–chin lift method or, if spinal injury is suspected, use the jaw thrust method, if possible.
 6. **Look, listen, and feel for adequate breathing (no more than 10 sec).**
 7. If the child is not breathing, **begin rescue breaths.** Using a bag-valve-mask device or face mask, **give 2 breaths (1 sec each) with sufficient volume to make the chest rise. Do not over-ventilate.**
 8. **If the chest does not rise, reposition the head, chin, and jaw, and attempt 2 more breaths.** If breaths cannot be delivered, perform Child CPR (p 248) beginning with chest compression using a 30:2 ratio at a rate of 100/min. Each time the airway is opened, look for an

OBSTRUCTED AIRWAY: Unconscious Infant (younger than 1 yr)

Clinical Presentation

- Inability to breathe, high-pitched noises
 - Inability to move air into lungs with rescue breaths
 - Cyanosis
1. **Establish unresponsiveness.** Gently rub the infant's back or chest.
 2. If no response send a second rescuer, if available, for help.
 3. If you are alone, begin the steps for CPR.
 4. **Position the infant supine** on a hard, flat surface.
 5. **Open the airway** by the head tilt–chin lift method or, if spinal injury is suspected, use the jaw thrust method, if possible.
 6. **Look, listen, and feel for adequate breathing (no more than 10 sec).**
 7. If the infant is not breathing, **begin rescue breaths.** Using a bag-valve-mask device or face mask, **give 2 breaths (1 sec each) using sufficient volume to make the chest rise. Do not over-ventilate.**
 8. **If the chest does not rise, reposition the head, chin, and jaw, and give 2 more breaths.** If breaths cannot be delivered, perform Infant CPR (p 249) beginning with chest compression using a 30:2 ratio at a rate of 100/min. Each time the airway is opened, look for an

object in the infant's mouth. **Only use a finger sweep to remove solid material you see obstructing the airway. Never perform a finger sweep if you do not see a foreign body in the airway.**

9. Continue CPR until breathing and circulation resume or until advanced life support measures are initiated.
10. After the fifth cycle, if you are still alone, summon help, call a code, or phone 911.
11. If circulation resumes but breathing does not resume or is inadequate, continue rescue breathing at 12–20 breaths per minute.
12. If adequate breathing and circulation resume, place the infant in the recovery position and monitor until help arrives.



Clinical Tip:

When you open an infant's airway by the head tilt–chin lift method, do not overextend the head or the airway will become obstructed.

CPR AND OBSTRUCTED AIRWAY POSITIONS




Head tilt-chin lift (adult or child).



Jaw thrust maneuver.



Bag-valve-mask.



Head tilt-chin lift (infant).



Universal choking sign.



Abdominal thrusts.

Advanced Cardiac Life Support Protocols

VENTRICULAR FIBRILLATION (VF) OR PULSELESS VENTRICULAR TACHYCARDIA (VT)

Clinical Presentation

- Unresponsive state
- No respiration, pulse, or blood pressure (BP)
 1. Establish unresponsiveness with no respiration or pulse. Call for help.
 2. Begin CPR, provide oxygen and attach AED or monitor-defibrillator when available without interrupting CPR.
 3. When device is attached, stop CPR and assess rhythm. If shock is advised when using an AED, defibrillate following AED prompts. If using a manual monitor-defibrillator and rhythm is ventricular fibrillation or pulseless ventricular tachycardia, defibrillate at 120–200 J if using a biphasic defibrillator following manufacturer's device-specific energy levels if known, or 200 J if unknown, or defibrillate at 360 J if using a monophasic defibrillator.
 4. Immediately resume CPR, beginning with compressions. Provide five cycles (2 min) of uninterrupted CPR. During CPR, establish IV or IO access. Prepare vasopressor dose (epinephrine or vasopressin).
 5. Assess rhythm. If the rhythm is shockable, follow AED prompts or defibrillate at same or higher energy for biphasic manual defibrillator or at 360 J for a monophasic manual defibrillator.
 6. Immediately resume CPR beginning with compressions and using five cycles of 30 compressions and 2 breaths.
 7. Consider insertion of an advanced airway (ET tube, LMA, or Combitube) if basic airway management is inadequate. Once an advanced airway is in place, compressions should be uninterrupted at 100/min and ventilations should be 8 to 10 breaths/min (1 breath every 6–8 sec).
 8. Administer epinephrine 1 mg (10 mL of 1:10,000) by the intravenous or intraosseous (IV/IO) method; follow with 20 mL IV flush. Repeat every 3–5 minutes; give 2.0–2.5 mg diluted in 10 mL normal saline if administering by endotracheal (ET) tube; or administer a single dose of vasopressin 40 U IV/IO to replace first or second dose of epinephrine.
 9. Continue CPR; check the rhythm every 2 minutes.
 10. If the rhythm is still shockable, defibrillate as in step 5.
 11. Immediately resume CPR, check the rhythm every 2 minutes.

Consider antiarrhythmics for shock-refractory VF or pulseless VT:

12. Administer amiodarone 300 mg IV/IO (diluted in 20–30 mL D5W); or lidocaine 1.0–1.5 mg/kg IV/IO.
13. Repeat antiarrhythmic therapy for shock-refractory VF or VT: amiodarone 150 mg IV/IO in 3–5 minutes; or lidocaine 0.5–0.75 mg/kg IV/IO. Repeat lidocaine every 5–10 minutes, maximum dose 3 mg/kg.
14. Consider magnesium sulfate 1–2 g (2–4 mL of a 50% solution) diluted in 10 mL of D5W IV/IO, given over 5–20 minutes, if arrhythmia is torsade de pointes.



Clinical Tip:

Do not delay defibrillation for a witnessed arrest. For an unwitnessed arrest with a down time greater than 4–5 minutes, perform two minutes of CPR prior to defibrillation.

**Clinical Tip:**

When giving drugs via peripheral IV, elevate extremity for 10–20 seconds to enhance drug delivery to core circulation.

**Clinical Tip:**

Airway must be secured and placement verified with observation of chest rise and auscultation of breath sounds plus a confirmatory device (exhaled CO₂ detector or esophageal detector device). Monitor tube for displacement during transport or whenever patient is moved.

PULSELESS ELECTRICAL ACTIVITY (PEA)**Clinical Presentation**

- Unresponsive state
 - No respiration, pulse, or BP
 - Identifiable electrical rhythm on monitor but no pulse
1. Establish unresponsiveness with no respiration or pulse. Call for help.
 2. Begin CPR, provide oxygen and attach manual monitor-defibrillator when available without interrupting CPR.
 3. When device is attached, stop CPR to assess rhythm. If identifiable rhythm noted on monitor, immediately resume CPR beginning with compressions. Establish IV/IO access.
 4. During CPR, consider and treat possible causes:

Trauma	Hypokalemia/hyperkalemia
Tension pneumothorax	Hypovolemia
Thrombosis (pulmonary or coronary)	Hypoxia
Tamponade, cardiac	Hypoglycemia
Toxins	Hypothermia
	Hydrogen ion (acidosis)
 5. Continue CPR using five cycles of 30 compressions and 2 breaths; check the rhythm every 2 minutes.
 6. Consider insertion of an advanced airway (ET tube, LMA, or Combitube) if basic airway management is inadequate. Once an advanced airway is in place, compressions should be uninterrupted at 100/min and ventilations should be 8 to 10 breaths/min (1 breath every 6–8 sec).
 7. If PEA persists, administer epinephrine 1 mg IV/IO (10 mL of 1:10,000) and follow with 20 mL IV flush; repeat every 3–5 minutes; give 2.0–2.5 mg diluted in 10 mL normal saline if administering via ET tube; or administer a single dose of vasopressin 40 U IV/IO to replace the first or second dose of epinephrine.

8. Consider atropine 1 mg IV/IO if the heart rate on the ECG is less than 60 bpm. Repeat every 3–5 minutes as needed up to a total of three doses (3 mg).
9. Continue CPR; check the rhythm every 2 minutes.
10. If the rhythm is shockable with no pulse, follow VF/VT protocol (p 253).
11. If the rhythm is not shockable with no pulse, resume CPR and repeat steps 4–8.
12. If a stable ECG rhythm returns with adequate breathing and circulation, monitor and reevaluate the patient.

**Clinical Tip:**

PEA is often caused by reversible conditions and can be treated if those conditions are identified and corrected.

ASYSTOLE**Clinical Presentation**

- Unresponsive state, no respiration, pulse, or BP
 - ECG shows flat line; no electrical activity
1. Establish unresponsiveness with no respiration or pulse. Call for help.
 2. Begin CPR, provide oxygen and attach manual monitor-defibrillator when available without interrupting CPR.
 3. When device is attached, stop CPR to assess rhythm. If no electrical activity (flat line or asystole) is noted on monitor, immediately resume CPR beginning with compressions. Establish IV/IO access.
 4. During CPR consider and treat possible causes:

Trauma	Hypokalemia/hyperkalemia
Tension pneumothorax	Hypovolemia
Thrombosis (pulmonary or coronary)	Hypoxia
Tamponade, cardiac	Hypoglycemia
Toxins	Hypothermia
	Hydrogen ion (acidosis)
 5. Continue CPR beginning with compressions and using five cycles of 30 compressions and 2 breaths; check the rhythm every 2 minutes.
 6. Consider insertion of an advanced airway (ET tube, LMA, or Combitube) if basic airway management is inadequate. Once an advanced airway is in place, compressions should be uninterrupted at 100/min and ventilations should be 8 to 10 breaths/min (1 breath every 6–8 sec).
 7. If asystole persists, administer epinephrine 1 mg IV/IO (10 mL of 1:10,000) and follow with 20 mL IV flush; repeat every 3–5 minutes; give 2.0–2.5 mg diluted in 10 mL normal saline if

- administering via ET tube; or administer a single dose of vasopressin 40 U IV/IO to replace the first or second dose of epinephrine.
8. Consider atropine 1 mg IV/IO if the ECG still shows asystole. Repeat every 3–5 minutes as needed up to a total of three doses (3 mg).
 9. Continue CPR; check the rhythm every 2 minutes.
 10. If the rhythm is shockable with no pulse, follow VF/VT protocol (p 253).
 11. If the rhythm is not shockable with no pulse, resume CPR and repeat steps 4–8.
 12. If asystole persists, consider whether proper resuscitation protocols were followed and reversible causes identified. If procedures were performed correctly, follow local criteria for terminating resuscitation efforts.
6. Obtain a 12 lead ECG.
 7. If the 12-lead ECG shows ST segment elevation, notify the attending physician to begin the checklist for fibrinolytic therapy. If possible, pre-hospital providers should transport the patient to the closest facility with rapid coronary intervention capabilities.
 8. Administer nitroglycerin by the sublingual route 0.3–0.4 mg (1 tablet), repeated for a total of three doses over 15-minute intervals; or administer aerosol spray for 0.5–1.0 seconds at 5-minute intervals (provides 0.4 mg per dose) not to exceed three sprays in 15 minutes. Nitroglycerin administration requires a systolic BP greater than 90 mm Hg.
 9. Repeat nitroglycerin (see step 8) until chest pain is relieved, systolic BP falls below 90 mm Hg, or signs of ischemia or infarction are resolved.
 10. If chest pain is not relieved by nitroglycerin, administer morphine 2–4 mg IV (over 1–5 min). If symptoms are not resolved, administer 2–8 mg every 5–15 min if hemodynamically stable. Do not administer morphine if systolic BP is less than 90 mm Hg.

**Clinical Tip:**

Always confirm asystole by checking the ECG in two different leads. Also, search to identify underlying VF.

**Clinical Tip:**

Transcutaneous pacing is not recommended for asystolic cardiac arrest.

**Clinical Tip:**

Study local policy to learn established criteria for stopping resuscitation efforts.

**Clinical Tip:**

Patients should not be given nitroglycerin if they have taken sildenafil (**Viagra**), or vardenafil (**Levitra**) in the last 24 hours or tadalafil (**Cialis**), within 48 hours. The use of nitroglycerin with these medication may cause irreversible hypotension.

**Clinical Tip:**

Nitroglycerin should be used with caution in patients with an inferior MI with possible right ventricular involvement. It is contraindicated with right ventricular MI.

**Clinical Tip:**

Diabetic patients, the elderly, and women frequently present with atypical symptoms (e.g., weakness, fatigue, complaints of indigestion).

ACUTE CORONARY SYNDROME (ACS)

Clinical Presentation

- History of acute MI or angina
- Chest pain or discomfort
- Pain spreading to neck, shoulders, arms, or jaw
- Sudden unexplained shortness of breath, weakness, fatigue with or without chest pain/discomfort
- Associated nausea, diaphoresis, lightheadedness, fainting.
 1. Establish responsiveness.
 2. Perform primary ABCD survey.
 3. Measure vital signs, including oxygen saturation.
 4. Administer oxygen at 4 L/min, start an IV, and attach cardiac monitor.
 5. Administer aspirin 160–325 mg orally (PO) if no history of aspirin allergy. Chewing the tablet is preferable; use non–enteric-coated tablets for antiplatelet effect. Give within minutes of onset.

BRADYCARDIA

Clinical Presentation

- Pulse rate less than 60 bpm
- Sinus bradycardia, junctional escape rhythm, or AV Block

- Symptoms of chest discomfort/pain, lightheadedness, dizziness
- Signs of hypotension, diaphoresis, altered mental status, congestive heart failure (CHF), shock
 1. Establish responsiveness.
 2. Perform primary ABCD survey.
 3. Measure vital signs, including oxygen saturation.
 4. Supply oxygen, start an IV and attach cardiac monitor to identify rhythm.
 5. Obtain a 12-lead ECG.
 6. If the patient is stable and asymptomatic with a heart rate less than 60 bpm, monitor and observe for any changes.
 7. If the patient is symptomatic with signs of poor perfusion, initiate treatment.
 8. In second-degree (Mobitz type II) or third-degree AV block, prepare for transcutaneous pacing.
 9. While awaiting the pacemaker, consider atropine 0.5 mg IV every 3–5 minutes, maximum total dose 3 mg.
 10. If the patient fails to respond to atropine, sedate and begin transcutaneous pacing.
 11. If the patient is still hypotensive with severe bradycardia, or if pacing is unavailable or ineffective, consider epinephrine infusion, 2–10 mcg/min IV (add 1 mg of 1:1000–500 mL normal saline and infuse at 1–5 mL/min) or dopamine with continuous infusions (titrate to patient response) of 2–10 mcg/kg/min. Mix 400 mg/250 mL in normal saline, lactated Ringer's solution, or D5W (1600 mcg/mL).



Clinical Tip:

If the patient is symptomatic, do not delay transcutaneous pacing while waiting for atropine to take effect or for IV access.



Clinical Tip:

Use atropine with caution in a suspected acute MI; atropine may lead to rate-induced ischemia.

TACHYCARDIA—UNSTABLE

Clinical Presentation

- Altered level of consciousness (LOC)
 - Symptoms: Shortness of breath, diaphoresis, weakness, fatigue, syncope or presyncope, chest discomfort or pain, palpitations
 - Signs: Hypotension, shock, congestive heart failure, ischemic ECG changes, poor peripheral perfusion
1. Establish responsiveness.
 2. Perform primary ABCD survey.
 3. Measure vital signs, including oxygen saturation.
 4. Supply oxygen, start an IV, and begin cardiac monitoring.
 5. Consider and treat possible causes:

Trauma	Hypokalemia/hyperkalemia
Tension pneumothorax	Hypovolemia
Thrombosis (pulmonary or coronary)	Hypoxia
Tamponade, cardiac	Hypoglycemia
Toxins	Hypothermia
	Hydrogen ion (acidosis)
 6. Establish that serious signs and symptoms are related to the tachycardia.
 7. If the patient is **unstable and symptomatic** with a heart rate >150 bpm, prepare for immediate synchronized cardioversion (patients with a healthy heart are unlikely to be unstable if the ventricular rate is less than 150 bpm; however patients with cardiac disease may be unstable with heart rates less than 150 bpm). It is important to look at the patient's stability in addition to monitoring heart rate as criteria for cardioversion.
 8. Premedicate with a sedative plus an analgesic whenever possible.
 9. Place the defibrillator in synchronized (sync) mode.
 10. Administer synchronized cardioversion at a monophasic energy level of 100 J (or, if using a biphasic manual defibrillator, use the manufacturer's device-specific energy levels for synchronized cardioversion, usually 120–200 J).
 11. If there is no change, administer synchronized cardioversion at a monophasic energy level of 200 J (or the equivalent energy level for a biphasic manual defibrillator).
 12. If there is no change, administer synchronized cardioversion at a monophasic energy level of 300 J (or the equivalent energy level for a biphasic manual defibrillator).
 13. If there is still no change, administer synchronized cardioversion at a monophasic energy level of 360 J (or the equivalent energy level for a biphasic manual defibrillator).
 14. If pulseless arrest develops, identify arrhythmia and follow algorithm for VF/VT (p 253), PEA (p 254), or asystole (p 254).



Clinical Tip:

Reactivate the "sync" mode before each attempted cardioversion.

**Clinical Tip:**

The “sync” mode delivers energy just after the R wave to avoid stimulation during the refractory, or vulnerable, period of the cardiac cycle when a shock could potentially produce VF.

**Clinical Tip:**

If the rhythm is very fast or irregular and it is not possible to synchronize a shock, deliver unsynchronized shocks (defibrillation) at higher energy levels.

NARROW-COMPLEX TACHYCARDIA— STABLE REGULAR RHYTHM

Clinical Presentation

- No *serious* signs or symptoms related to the tachycardia
- Regular ECG rhythm
- QRS narrow (<0.12 sec)
 1. Establish responsiveness.
 2. Perform primary ABCD survey.
 3. Measure vital signs, including oxygen saturation.
 4. Supply oxygen, start an IV, and attach cardiac monitor to identify rhythm. Obtain a 12-lead ECG.
 5. Consider and treat possible causes:

Trauma	Hypokalemia/hyperkalemia
Tension pneumothorax	Hypovolemia
Thrombosis (pulmonary or coronary)	Hypoxia
	Hypoglycemia
Tamponade, cardiac	Hypothermia
Toxins	Hydrogen ion (acidosis)
 6. If the arrhythmia is PSVT (stable reentry SVT), attempt vagal maneuvers (carotid sinus massage or Valsalva maneuver).
 7. If rhythm has not converted to sinus rhythm, administer adenosine 6 mg IV in the antecubital or other large vein given rapidly over 1–3 seconds followed by a 20-mL bolus of normal saline and elevate arm immediately.
 8. If the rhythm has not converted in 1–2 minutes, repeat adenosine at 12 mg IV. If the rhythm still does not convert, a third dose of 12 mg IV may be given after another 1–2 minutes, maximum 30 mg.
 9. If the rhythm still does not convert, it may be atrial flutter, atrial tachycardia, or junction-

al tachycardia. Consider rate control using diltiazem or beta blockers. Obtain expert consultation.

10. If the rhythm converts, the arrhythmia was possibly PSVT (reentry SVT). Observe the patient and treat any recurrence with adenosine, diltiazem, or beta blockers. Obtain expert consultation.

**Clinical Tip:**

If the patient’s condition becomes unstable during the tachycardia, perform immediate synchronized cardioversion.

**Clinical Tip:**

Use beta blockers with caution in patients with pulmonary disease or congestive heart failure. Avoid use in patients with bronchospastic disease.

NARROW-COMPLEX TACHYCARDIA— STABLE IRREGULAR RHYTHM

Clinical Presentation

- No *serious* signs or symptoms related to the tachycardia
- Irregular ECG rhythm
- QRS narrow (<0.12 sec)
 1. Establish responsiveness.
 2. Perform primary ABCD survey.
 3. Measure vital signs, including oxygen saturation.
 4. Supply oxygen, start an IV, and attach cardiac monitor to identify rhythm. Obtain 12-lead ECG.
 5. Consider and treat possible causes:

Trauma	Hypokalemia/hyperkalemia
Tension pneumothorax	Hypovolemia
Thrombosis (pulmonary or coronary)	Hypoxia
	Hypoglycemia
Tamponade, cardiac	Hypothermia
Toxins	Hydrogen ion (acidosis)
 6. Irregular rhythms (QRS <0.12 sec) are probably atrial fibrillation, atrial flutter, or multifocal atrial tachycardia (MAT). Consider using diltiazem or beta blockers to control the rate. Consider expert consultation.
 7. If pulseless arrest develops, identify arrhythmia and follow algorithm for VF/VT (p 253), PEA (p 254), or asystole (p 254).

**Clinical Tip:**

If the patient's condition becomes unstable during the tachycardia, perform immediate synchronized cardioversion.

**Clinical Tip:**

Beta blockers should be used with caution in patients with pulmonary disease or CHF. Avoid in patients with bronchospastic disease.

WIDE-COMPLEX TACHYCARDIA— STABLE REGULAR RHYTHM

Clinical Presentation

- No *serious* signs and symptoms related to the tachycardia
- Regular ECG rhythm
- QRS wide (>0.12 sec)
 1. Establish responsiveness.
 2. Perform primary ABCD survey.
 3. Measure vital signs, including oxygen saturation.
 4. Supply oxygen, start an IV, and attach cardiac monitor to identify rhythm. Obtain a 12-lead ECG.
 5. Consider and treat possible causes:

Trauma	Hypokalemia/hyperkalemia
Tension pneumothorax	Hypovolemia
Thrombosis (pulmonary or coronary)	Hypoxia
Tamponade, cardiac	Hypoglycemia
Toxins	Hypothermia
	Hydrogen ion (acidosis)
 6. If the arrhythmia is monomorphic VT or an uncertain rhythm, consider expert consultation. Administer amiodarone 150 mg IV/IO over 10 minutes. Repeat as needed to a maximum dose of 2.2 g in 24 hours.
 7. If the rhythm still does not convert, prepare for synchronized cardioversion.
 8. Premedicate with a sedative plus an analgesic.
 9. Administer synchronized cardioversion incrementally at monophasic energy levels of 100 J, 200 J, 300 J, and 360 J. If you are using biphasic equipment, the optimal initial dose is 100–120 J, with escalation as needed.
 10. If the rhythm still does not convert and the arrhythmia is SVT with aberrant conduction, administer adenosine 6 mg IV in the antecubital or other large vein given rapidly over 1–3 seconds followed by a 20-mL bolus of normal saline and elevate arm immediately.
 11. If the rhythm does not convert and arrhythmia is still SVT with aberrant conduction, wait 1–2 minutes, then give a second rapid dose of adenosine 12 mg IV. A third rapid dose of 12 mg IV may be given after another 1–2 minutes, maximum dose of 30 mg.

12. If pulseless arrest develops, identify arrhythmia and follow algorithm for VF/VT.

WIDE-COMPLEX TACHYCARDIA— STABLE IRREGULAR RHYTHM

Clinical Presentation

- No *serious* signs or symptoms related to the tachycardia
- Irregular rhythm
- QRS wide (>0.12 sec)
 1. Establish responsiveness.
 2. Perform primary ABCD survey.
 3. Measure vital signs, including oxygen saturation.
 4. Supply oxygen, start an IV, and attach cardiac monitor to identify rhythm. Obtain a 12-lead ECG.
 5. Consider and treat possible causes:

Trauma	Hypokalemia/hyperkalemia
Tension pneumothorax	Hypovolemia
Thrombosis (pulmonary or coronary)	Hypoxia
Tamponade, cardiac	Hypoglycemia
Toxins	Hypothermia
	Hydrogen ion (acidosis)
 6. If the arrhythmia is atrial fibrillation with aberrant conduction, consider expert consultation. Consider controlling the rate using diltiazem or beta blockers.
 7. If the arrhythmia is pre-excited A-fib (A-fib with Wolff-Parkinson-White Syndrome), seek expert consultation. Consider amiodarone, 150 mg IV/IO over 10 minutes. Avoid AV nodal blocking agents (e.g., adenosine, digoxin, diltiazem, verapamil).
 8. If the arrhythmia is recurrent polymorphic VT, seek expert consultation.
 9. If the arrhythmia is torsade de pointes, administer magnesium sulfate 1–2 g diluted in 50–100 mL of D5W IV/IO, given over 5–60 minutes, followed by an infusion at 0.5–1 g/hr titrated to control torsades.
 10. If pulseless arrest develops, identify the arrhythmia and follow the algorithm for VF/VT (p 253), PEA (p 254), or asystole (p 254).

**Clinical Tip:**

Monomorphic and polymorphic VT may rapidly deteriorate to VF.

Emergency Medications

This list is a reference list only. It is not meant to be exhaustive in clinical content.

- Always consult an authoritative, current reference about dose, dilution, route and rate of administration, and interactions before administering medications, especially IV medications. Have a second licensed person independently check dose calculations, preparation, original orders, and infusion pump programming.

❖ ACE INHIBITORS (Angiotensin-Converting Enzyme Inhibitors)

Class: Antihypertensive

Common Agents: Captopril, Enalapril, Lisinopril, Ramipril.

Indications: Myocardial infarction, hypertension (HTN), congestive heart failure (CHF), heart failure without hypotension, ST segment elevation, left ventricular dysfunction after MI.

Dose: See individual order and drug for route and dosage. Usually not started in the Emergency Department, but within 24 hours after fibrinolytic therapy has been completed and BP has stabilized.

Contraindications: Lactation, pregnancy, angioedema, hypersensitivity to ACE inhibitors, serum potassium more than 5 mEq/L.

Side Effects: Cough, dizziness, headache, fatigue, hypotension, hyperkalemia.

Precautions: Reduce dose in renal failure.

❖ ADENOSINE (Adenocard, Adenoscan)

Class: Antiarrhythmic

Indications: Regular narrow-complex tachycardias and PSVT.

Dose: 6 mg IV in the antecubital or other large vein given rapidly over 1–3 seconds followed by a 20-mL bolus of normal saline. Immediately elevate arm. If the rhythm does not convert, give 12 mg by IV in 1–2 min-

utes if needed. A third dose of 12 mg IV may be given in another 1–2 minutes, maximum total dose 30 mg.

Contraindications: Hypersensitivity, sick sinus syndrome, second- or third-degree AV block (unless a functional artificial pacemaker is present), drug- or poison-induced tachycardia.

Side Effects: Flushing, dizziness, headache, dyspnea, bronchospasm, chest pain or tightness, discomfort in neck, throat, or jaw, bradycardia, AV block, asystole, ventricular ectopic beats, VF.

Precautions: Ineffective in treating A-fib, A-flutter, or VT. Avoid in patients receiving dipyridamole, theophylline, or caffeine.

❖ AMIODARONE (Cordarone, Pacerone)

Class: Antiarrhythmic

Indications: Management of life-threatening recurrent VF or refractory hemodynamically-unstable VT. Conversion of atrial fibrillation, SVT. Control of rapid ventricular rate in pre-excited atrial arrhythmias. Control of hemodynamically stable VT, polymorphic VT with normal QT interval, or wide-complex tachycardia of uncertain origin.

Dose: *Cardiac arrest* 300 mg IV/IO (diluted in 20–30 mL D5W); consider additional 150 mg IV/IO in 3–5 minutes. *Wide- and narrow-complex tachycardia (stable)* 150 mg IV over first 10 minutes (15 mg/min)—may repeat infusion of 150 mg IV every 10 minutes as needed; slow infusion of 360 mg IV over next 6 hours (1 mg/min); maintenance infusion of 540 mg over next 18 hours (0.5 mg/min). Maximum cumulative dose 2.2 g IV in 24 hours.

Contraindications: Hypersensitivity, cardiogenic shock, symptomatic bradycardia or second- or third-degree AV block without functioning artificial pacemaker.

Side Effects: Vasodilation, hypotension, visual impairment, hepatotoxicity, pulmonary toxicity,

CHF; may prolong QT interval, producing torsade de pointes.

Precautions: Avoid concurrent use with procainamide. Correct hypokalemia and hypomagnesemia, if possible, before use. Draw up amiodarone through a large-gauge needle to reduce foaming. For slow or maintenance IV infusion, mix the medication only in a glass bottle containing D5W or NS and administer through an in-line filter.

❖ **ASPIRIN (Acetylsalicylic Acid)**

Class: Antiplatelet

Indications: Acute coronary syndrome, symptoms suggestive of cardiac ischemia.

Dose: 160–325 mg PO. Chewing the tablet is preferable; use non-enteric coated tablets for antiplatelet effect. Give within minutes of onset of ischemic symptoms.

Contraindications: Known allergy to aspirin, pregnancy.

Side Effects: Anorexia, nausea, epigastric pain, anaphylaxis.

Precautions: Active ulcers and asthma, bleeding disorders or thrombocytopenia.

❖ **ATROPINE SULFATE**

Class: Antiarrhythmic, Anticholinergic

Indications: Symptomatic sinus bradycardia, junctional escape rhythm, or second-degree type I block, asystole, bradycardic PEA (<60 bpm). Not likely to be effective in second-degree type II or third-degree AV block with wide QRS complex.

Dose: *Cardiac arrest* 1 mg IV/IO every 3–5 minutes (may give through an ET tube at 2.0–3.0 mg diluted in 10 mL normal saline). May repeat to a maximum of three doses (3 mg). *Bradycardia* 0.5mg IV given every 3–5 minutes as needed, maximum total dose 3 mg (0.04 mg/kg).

Contraindications: Atrial fibrillation, A-flutter, glaucoma, asthma, obstructive uropathy.

Side Effects: Tachycardia, headache, dry mouth, dilated pupils, flushing, hypotension.

Precautions: Use caution in myocardial ischemia and hypoxia. Avoid in hypothermic bradycardia and in second-degree (Mobitz type II) and third-degree AV block.

❖ **BETA BLOCKERS**

Class: Antihypertensive, Antiarrhythmic, Antianginal

Common Agents: Atenolol, Esmolol, Labetalol, Metoprolol tartrate, Propranolol.

Indications: Myocardial infarction, unstable angina, PSVT, A-fib, A-flutter, HTN.

Dose: See individual order and drug for route and dosage.

Contraindications: Heart rate less than 60 bpm, systolic BP less than 100 mm Hg, second- or third-degree AV block, severe left ventricular failure.

Side Effects: Hypotension, dizziness, bradycardia, headache, nausea and vomiting.

Precautions: Concurrent use with calcium channel blockers, such as verapamil or diltiazem, can cause hypotension. Use caution in patients with a history of bronchospasm or cardiac failure. Use caution in patients with peripheral arterial disease and with diabetic patients (monitor blood glucose levels frequently).

❖ **CALCIUM CHLORIDE**

Class: Minerals, Electrolytes, Calcium Salt

Indications: Hyperkalemia, hypocalcemia, hypomagnesemia; antidote to calcium channel blockers and beta blockers; given prophylactically with calcium channel blockers to prevent hypotension.

Dose: *Hyperkalemia and antidote to calcium channel blocker*

8–16 mg/kg (usually 5–10 mL of a 10% solution) IV; may be repeated as needed. *Prophylaxis prior to IV calcium channel blockers* 2–4 mg/kg (usually 2 mL of a 10% solution) IV.

Contraindications: Hypercalcemia, VF, digoxin toxicity, renal calculi.

Side effects: Bradycardia, hypotension, hypomagnesemia, VF, nausea and vomiting.

Precautions: Incompatible with sodium bicarbonate (precipitates).

❖ **DIGOXIN (Lanoxin)**

Class: Inotropic, Antiarrhythmic

Indications: To slow ventricular response in A-fib or A-flutter, as a positive inotrope in CHF or pulmonary edema. May be used as an alternative drug for PSVT (reentry SVT).

Dose: Loading dose of 10–15 µg/kg, administered IV over 5 minutes. Maintenance dose determined by body size and renal function.

Contraindications: Hypersensitivity, uncontrolled ventricular arrhythmias, AV block without functioning artificial pacemaker, idiopathic hypertrophic subaortic stenosis (IHSS), constrictive pericarditis, atrial fibrillation with Wolff-Parkinson-White syndrome.

Side Effects: Accelerated junctional rhythm, atrial tachycardia with block, AV block, asystole, VT, VF; dizziness, weakness, fatigue; nausea and vomiting; blurred or yellow vision; headache; hypersensitivity; hypokalemia.

Precautions: Avoid electrical cardioversion of stable patients. If the patient's condition is unstable, use lower current settings such as 10–20 J. Use cautiously in elderly patients. Correct electrolyte abnormalities, monitor digoxin levels, monitor for clinical signs of toxicity. Hypokalemia may precipitate digitalis toxicity.

❖ DIGOXIN IMMUNE FAB (Fragment Antigen Binding) (Digibind)

Class: Antidote to digoxin and digitoxin

Indications: Symptomatic digoxin toxicity or acute ingestion of unknown amount of digoxin.

Dose: Depends on serum digoxin levels. One 40-mg vial binds to approximately 0.6 mg of digoxin. Dose is typically administered over 30 minutes.

Contraindications: Allergy only, otherwise none known.

Side Effects: Worsening of CHF, rapid ventricular response in patients with A-fib, hypokalemia; increased serum digoxin levels due to bound complexes (clinically misleading since bound complex cannot interact with receptors).

Precautions: Allergy to sheep proteins or other sheep products.

❖ DILTIAZEM (Cardizem)

Class: Calcium Channel Blocker

Indications: To control ventricular rate in A-fib and A-flutter, PSVT (reentry SVT) refractory to adenosine with narrow QRS complex and adequate BP.

Dose: 15–20 mg (0.25 mg/kg) IV given over 2 minutes. May repeat in 15 minutes at 20–25 mg (0.35 mg/kg) IV given over 2 minutes. Start maintenance drip at 5–15 mg/hr and titrate to HR.

Contraindications: Drug- or poison-induced tachycardia, wide-complex tachycardia of uncertain origin, rapid A-fib and A-flutter with Wolff-Parkinson-White syndrome, sick sinus syndrome, second- or third-degree AV block (unless a functional artificial pacemaker is present), hypotension with systolic BP less than 90 mm Hg.

Side Effects: Hypotension, bradycardia (including AV block), chest pain, ventricular arrhythmias peripheral edema, flushing.

Precautions: Severe hypotension in patients receiving beta blockers, hepatic injury, renal disease.

❖ DOPAMINE (Intropin)

Class: Vasopressor, Inotropic, Adrenergic Agonist

Indications: Symptomatic bradycardia and hypotension, cardiogenic shock.

Dose: Continuous infusions (titrate to patient response): Low dose 1–5 µg/kg/min; moderate dose 5–10 µg/kg/min (cardiac doses); high dose 10–20 µg/kg/min (vasopressor doses). Mix 400 mg/250 mL in normal saline, lactated Ringer's solution, or D5W (1600 µg/mL).

Contraindications: Hypersensitivity to sulfites, pheochromocytoma, VF.

Side Effects: Tachyarrhythmias, angina, hypotension, palpitations, vasoconstriction, dyspnea, nausea and vomiting.

Precautions: Hypovolemia, MI. Adjust dosage in elderly patients and in those with occlusive vascular disease. Ensure adequate hydration prior to infusion. Taper slowly. Do not mix with sodium bicarbonate. Use care with peripheral administration; infiltration can cause tissue necrosis. A central line is preferred.

❖ EPINEPHRINE (Adrenalin)

Class: Adrenergic Agonist

Indications: Cardiac arrest: PEA, asystole, pulseless VT, VF; hypotension with severe bradycardia.

Dose: *Cardiac arrest* 1 mg IV/IO (10 mL of 1:10,000 solution) given every 3–5 minutes as needed; follow each dose with 20 mL IV flush. Give 2.0–2.5 mg diluted in 10 mL normal saline if administering by ET tube. *Profound bradycardia or hypotension* 2–10 µg/min IV; add 1 mg (1 mL of a 1:1000 solution) to 500 mL normal saline or D5W.

Contraindications: Hypersensitivity to adrenergic amines, hypovolemic shock, coronary insufficiency.

Side Effects: Angina, HTN, tachycardia, VT, VF, nervousness, restlessness, tremors, weakness, headache, nausea.

Precautions: Use caution in HTN and increasing heart rate (may cause increased myocardial oxygen demand). Higher doses can contribute to post-arrest cardiac impairment, but they may be required to treat poison- or drug-induced shock. Avoid mixing with alkaline solutions.

❖ FIBRINOLYTIC AGENTS

Class: Thrombolytic, Fibrinolytic

Common Agents: Alteplase (**Activase**, **t-PA**), Anistreplase (**Eminase**), Reteplase (**Retavase**), Streptokinase (**Streptase**), Tenecteplase (**TNKase**).

Indications: Acute MI symptoms within the last 12 hours. Alteplase is the only fibrinolytic agent approved for acute ischemic stroke and must be started less than 3 hours from the onset of symptoms.

Dose: See individual order and drug for route and dosage.

Contraindications: Active internal bleeding within 21 days (except menses), neurovascular event within 3 months, major surgery or trauma within 2 weeks, aortic dissection, severe (uncontrolled) HTN, bleeding disorders, prolonged CPR, lumbar puncture within 1 week.

Side Effects: Hypotension, reperfusion, arrhythmias, heart failure, headache, increased bleeding time, deep or superficial hemorrhage, flushing, urticaria, anaphylaxis.

Precautions: Use cautiously in patients with severe renal or hepatic disease. Initiate bleeding precautions. Monitor patient for bleeding complications.

❖ FUROSEMIDE (Lasix)

Class: Diuretic, Loop Diuretics

Indications: Congestive heart failure with acute pulmonary edema, hypertensive crisis, post-arrest cerebral edema, hepatic or renal disease.

Dose: 0.5–1.0 mg/kg IV given over 1–2 minutes; may repeat at 2 mg/kg IV given over 1–2 minutes.

Contraindications: Hypersensitivity (cross-sensitivity with thiazides and sulfonamides may occur), uncontrolled electrolyte imbalance, hepatic coma, anuria, hypovolemia.

Side Effects: Severe dehydration, hypovolemia, hypotension, hypokalemia, hyponatremia, hypochloremia, hyperglycemia, dizziness, ototoxicity.

Precautions: Use cautiously in severe liver disease accompanied by cirrhosis or ascites, electrolyte depletion, diabetes mellitus, pregnancy, lactation. Risk for ototoxicity with increased dose or rapid injection. Monitor electrolytes closely.

❖ IBUTILIDE (Corvert)

Class: Antiarrhythmic

Indications: Supraventricular tachycardia, including A-fib and A-flutter; most effective for conversion of A-fib or A-flutter of short duration (≤ 48 hrs).

Dose: *Patients weighing 60 kg or more* 1 mg IV given over 10 minutes; may repeat the same dose in 10 minutes if arrhythmia does not terminate. *Patients weighing less than 60 kg* 0.01 mg/kg IV given over 10 minutes; may repeat the same dose in 10 minutes if arrhythmia does not terminate.

Contraindications: Known hypersensitivity, history of polymorphic VT, QT_c greater than 440 msec.

Side Effects: Nonsustained or sustained monomorphic or polymorphic VT, torsade de pointes, AV block, CHF, HTN, headache, hypotension, nausea and vomiting.

Precautions: Monitor ECG for 4–6 hours after administration, with a defibrillator nearby. Correct electrolyte abnormalities prior to use. If A-fib has lasted longer than 48 hours, anticoagulation is required before cardioversion with ibutilide.

❖ ISOPROTERENOL (Isuprel)

Class: Sympathomimetic, Beta-Adrenergic Agonist

Indications: Medically-refractory symptomatic bradycardia when transcutaneous or transvenous pacing is not available, refractory torsade de pointes unresponsive to magnesium, bradycardia in heart transplant patients, beta blocker poisoning.

Dose: IV infusion: mix 1 mg/250 mL in normal saline, lactated Ringer's solution, or D5W, run at 2–10 $\mu\text{g}/\text{min}$, and titrate to patient response. In torsade de pointes, titrate to increase heart rate until VT is suppressed.

Contraindications: Hypersensitivity to drug or sulfites, digitalis intoxication, angina, concurrent use with epinephrine (can cause VF or VT).

Side Effects: Arrhythmias, cardiac arrest, hypotension, angina, anxiety, tachycardia, palpitations, skin flushing.

Precautions: May increase myocardial ischemia, tachycardia, restlessness. High doses are harmful except in beta blocker overdose.

❖ LIDOCAINE (Xylocaine)

Class: Antiarrhythmic, Local Anesthetic

Indications: Ventricular fibrillation or pulseless VT, stable VT, wide-complex tachycardia of uncertain origin.

Dose: *Cardiac arrest from VF or VT* 1.0–1.5 mg/kg IV/IO (or 2–4 mg/kg via ET tube); may repeat 0.5–0.75 mg/kg IV/IO every 5–10 minutes, maximum dose 3 mg/kg. *Stable VT, wide-complex tachycardia of uncertain origin* use 0.50–0.75 mg/kg and up to 1.0–1.5 mg/kg; may repeat 0.50–0.75 mg/kg every 5–10 minutes, maximum total dose 3.0 mg/kg. If conversion is successful, start an IV infusion of 1–4 mg/min (30–50 $\mu\text{g}/\text{kg}/\text{min}$) in normal saline or D5W.

Contraindications: Prophylactic use in acute MI, advanced AV block without functioning artificial pacemaker, hypotension, Wolff-Parkinson-White syndrome, hypersensitivity to amide local anesthetics.

Side Effects: Confusion, agitation, anxiety, tinnitus, tremors, hallucinations, seizures, hypotension, bradycardia, cardiovascular collapse, respiratory arrest.

Precautions: Congestive heart failure, respiratory depression, shock. Reduce maintenance dose (not loading dose) in presence of impaired liver function or left ventricular dysfunction or in the elderly. Stop infusion if signs of CNS toxicity develop.

❖ MAGNESIUM SULFATE

Class: Electrolyte, Antiarrhythmic

Indications: Torsade de pointes.

Dose: *Torsade de pointes (cardiac arrest)* 1–2 g IV (2–4 mL of a 50% solution) diluted in 10 mL of D5W over 5–20 minutes. *Torsade de pointes (non-cardiac arrest)* load with 1–2 g mixed in 50–100 mL of D5W infused over 5–60 minutes IV, then infuse 0.5–1.0 g/hr IV (titrate to control torsade).

Contraindications: Hypermagnesemia, hypocalcemia, renal disease, AV block, toxemia of pregnancy 2 hours prior to delivery.

Side Effects: Hypotension, bradycardia, cardiac arrest, respiratory depression, altered LOC, flushed skin, diaphoresis, hypocalcemia, hyperkalemia, hypophosphatemia.

Precautions: Renal insufficiency, occasional fall in BP with rapid administration. Monitor serum magnesium levels.

❖ MORPHINE SULFATE

Class: Opiate Narcotic Analgesic

Indications: Chest pain unrelieved by nitroglycerin, CHF and dyspnea associated with pulmonary edema.

Dose: 2–4 mg IV (given over 1–5 min), administer every 5–30 minutes if hemodynamically stable; may repeat dose of 2–8 mg at 5- to 15-minute intervals.

Contraindications: Hypersensitivity, heart failure due to chronic lung disease, respiratory depression, hypotension. Avoid in patients with RV infarction.

Side Effects: Respiratory depression, hypotension, nausea and vomiting, bradycardia, altered LOC, seizures.

Precautions: Administer slowly and titrate to effect. Reverse with naloxone (0.4–2.0 mg IV), if necessary. Use caution in cerebral edema and pulmonary edema with compromised respiration. Use caution with hypovolemic patients; be prepared to administer volume.

❖ NITROGLYCERIN (Nitrostat, Nitrolingual [Pump spray])

Class: Antianginal, Nitrate, Vasodilator

Indications: Acute coronary syndrome, angina, CHF associated with acute MI, hypertensive urgency with ACS.

Dose: Sublingual route, 0.3–0.4 mg (1 tablet); repeat every 3–5 minutes, maximum 3 doses/15 min. Aerosol, spray for 0.5–1.0 sec at 3- to 5-minute intervals (provides 0.4 mg/dose), maximum 3 sprays/15 min. Intravenous administration at 12.5–25.0 µg (if no sublingual or spray used). Intravenous infusion: mix 25 mg/250 mL (100 µg/mL) in D5W, run at 5–20 µg/min, and titrate to desired response.

Contraindications: Hypersensitivity, systolic BP less than 90 mm Hg; severe bradycardia or severe tachycardia associated with hypotension; sildenafil (**Viagra**), tadalafil (**Cialis**), vardenafil (**Levitra**) within 24 hours; right ventricular infarction.

Side Effects: Hypotension with reflex tachycardia, syncope, headache, flushed skin.

Precautions: Do not mix with other medications; titrate IV to maintain systolic BP above 90 mm Hg. Mix only in glass IV bottles and infuse only through tubing provided by manufacturer; standard polyvinyl chloride tubing can bind up to 80% of the medication, making it necessary to infuse higher doses. Do not shake aerosol spray (affects metered dose).

❖ OXYGEN

Class: Gas

Indications: Cardiopulmonary emergencies with shortness of breath and chest pain, cardiac or respiratory arrest, hypoxemia.

Dose: Nasal cannula 1–6 L/min (24%–44% oxygen), Venturi mask 4–8 L/min (24%–40% oxygen), simple mask 5–8 L/min (40%–60% oxygen), partial rebreathing mask 6–15 L/min (35%–60% oxygen), non-rebreathing mask 6–15 L/min (60%–90% oxygen), bag-valve-mask 15 L/min (up to 100% oxygen).

Contraindications: None reported.

Side Effects: Drying of respiratory mucosa, possible bronchospasm if oxygen is extremely cold and dry. Oxygen supports combustion and can fuel a fire.

Precautions: Respiratory arrest in patients with hypoxic respiratory drive. The patient needs an airway and adequate ventilation before oxygen is effective.

❖ PROCAINAMIDE (Pronestyl)

Class: Antiarrhythmic

Indications: Recurrent VT or VF, PSVT refractory to adenosine and vagal stimulation, rapid A-fib with Wolff-Parkinson-White syndrome, stable wide-complex tachycardia of uncertain origin, maintenance after conversion.

Dose: 20 mg/min IV infusion or up to 50 mg/min under urgent conditions, maximum 17 mg/kg loading dose. Maintenance IV infusion: mix 1 g/250 mL (4 mg/mL) in normal saline or D5W, run at 1–4 mg/min.

Contraindications: Second- and third-degree AV block (unless a functioning artificial pacemaker is in place), prolonged QT interval, torsade de pointes, hypersensitivity.

Side Effects: Hypotension, widening QRS, headache, nausea and vomiting, flushed skin, seizures, ventricular arrhythmias, AV block, cardiovascular collapse, arrest.

Precautions: Monitor BP every 2–3 minutes while administering procainamide. If QRS width increases by 50% or more, or if systolic BP decreases to less than 90 mm Hg, stop the drug. Monitor for prolonged PR interval and heart block. Monitor for QT prolongation. May precipitate or exacerbate CHF. Reduce the total dose to 12 mg/kg and maintenance infusion to 1–2 mg/min if cardiac or renal dysfunction is present. Use cautiously in myasthenia gravis, in hepatic or renal disease, and with drugs that prolong the QT interval (e.g., amiodarone, sotalolol).

❖ SODIUM BICARBONATE

Class: Alkalinizing Agent, Buffer

Indications: Known preexisting hyperkalemia, bicarbonate-responsive acidosis, prolonged resuscitation with effective ventilation.

Dose: 1 mEq/kg given rapidly; may repeat 0.5 mEq/kg every 10 minutes.

Contraindications: Metabolic and respiratory alkalosis, hypocalcemia, hypokalemia, hypercarbic acidosis.

Side Effects: Hypokalemia, hypocalcemia, hypernatremia, metabolic alkalosis, edema, seizures, tetany, exacerbation of CHF.

Precautions: Congestive heart failure, renal disease, cirrhosis, toxemia, concurrent corticosteroid therapy. Not recommended for routine use in cardiac arrest patients because adequate ventilation and CPR are the major “buffer agents” in cardiac arrest. Incompatible

with many drugs; flush the line before and after administration.

❖ **VASOPRESSIN (Pitressin)**

Class: Vasopressor, Hormone

Indication: Cardiac arrest: an alternative to epinephrine in shock-refractory VF and pulseless VT, PEA, and asystole.

Dose: *Cardiac arrest* 40 units IV single dose to replace first or second dose of epinephrine as an alternative.

Contraindications: Hypersensitivity.

Side Effects: Bradycardia, HTN, angina, MI, arrhythmias, dizziness, headache, nausea and vomiting, abdominal cramps, diaphoresis, bronchoconstriction, anaphylaxis.

Precautions: Coronary artery disease (may precipitate angina or MI), renal impairment; patients with seizure disorders, asthma, vascular disease.

❖ **VERAPAMIL (Calan, Isoptin)**

Class: Calcium Channel Blocker, Antiarrhythmic, Antihypertensive

Indications: Paroxysmal supraventricular tachycardia (with narrow QRS and adequate BP) refractory to adenosine, rapid ventricular rates in A-fib, A-flutter, or MAT.

Dose: 2.5–5.0 mg IV over 2 minutes; may give second dose, if needed, of 5–10 mg IV in 15–30 minutes, maximum dose 20 mg. An alternative second dose is 5 mg IV every 15 minutes, maximum dose 30 mg.

Contraindications: Atrial fibrillation with Wolff-Parkinson-White syndrome, wide-complex tachycardia of uncertain origin, second- or third-degree AV block (unless a functioning artificial pacemaker is in place), sick sinus syndrome, hypotension, severe CHF, cardiogenic shock.

Side Effects: Hypotension, exacerbation of CHF with left ventricular dysfunction, bradycardia, AV block, constipation, peripheral edema.

Precautions: Concurrent oral beta blockers, CHF, impaired hepatic or renal function; may decrease myocardial contractility. In geriatric patients administer slowly over 3 minutes.

Emergency Medical Skills

DEFIBRILLATION

Indications: Ventricular fibrillation or pulseless VT.

Energy Levels: Adult monophasic energy levels: deliver the first shock at 360 J. (If using a biphasic manual defibrillator, use the manufacturer's device-specific energy levels, usually 120–200 J.) Continue at a monophasic energy level of 360 J for further shocks. (With a biphasic manual defibrillator, use the device-specific energy levels, usually 120–200 J for further shocks.)

Application: Use handheld paddles or remote adhesive pads (Fig. D-1). Always use a conducting gel with paddles and apply firm pressure to the chest to ensure good skin contact. Dry moisture off skin; shave excessive hair.

Methods: Manual or automated.

Precautions: Place paddles and pads several inches away from an implanted pacemaker.




Fig. D.1 ■ Defibrillation.



Clinical Tip:

Defibrillation may be used on children (1 yr to adolescent [12–14 yr]). Always use pediatric paddles or pads and follow pediatric protocols. Data are insufficient to allow a recommendation for or against the use of AEDs for infants younger than 1 year. Refer to your local protocols for the use of an AED on an infant.

Manual Defibrillation

A manual defibrillator is used to restore a normal heart rhythm. For a patient experiencing sudden cardiac arrest, first assess for responsiveness, respiration, and pulse. If the ECG tracing is available, use it to verify that the rhythm is either VF or pulseless VT, and then manually deliver an electric shock to the heart.

Procedure

1. Verify that the patient is in cardiac arrest, with no pulse or respiration. Have someone provide CPR while the defibrillator is obtained and placed next to the patient.
2. Turn on the defibrillator; verify that all cables are connected.
3. Turn “lead select” to “paddles” or “defibrillator.”
4. Select the initial energy level for an adult to a monophasic energy level of 360 J. (With a biphasic manual defibrillator, use the manufacturer's device-specific energy levels, usually 120–200 J.)
5. Paddles: Use conducting gel and place on the apex (lower left chest, midaxillary) and sternum (right of sternum, midclavicular).
Pads: Place in locations specified for paddles.
6. Verify rhythm as VF or pulseless VT.
7. Say, “Charging defibrillator, stand clear!”
8. Charge the defibrillator.

9. Say, “I’m going to shock on three. One, I’m clear; two, you’re clear; three, everybody’s clear.” Perform a visual sweep to ensure all rescue personnel are clear of the patient, bed, and equipment.
10. Discharge the defibrillator, reassess the rhythm, and refer to appropriate advanced cardiac life support protocol.

Automated External Defibrillator

An automated external defibrillator (AED) is a small, lightweight device used by both professionals and laypersons to assess heart rhythm by computer analysis. Using voice and visual prompts, it administers an electric shock, if necessary, to restore a normal rhythm in patients with sudden cardiac arrest. A shock is administered only if the rhythm detected is VF or pulseless VT.

Automated external defibrillators are available from medical device manufacturers and local pharmacies. Although the AEDs all operate in basically the same way, they may vary from model to model. Be sure to follow the manufacturer’s recommendations.

Indications: Ventricular fibrillation or pulseless VT in adults and children. Insufficient data are available to allow a recommendation for or against the use of AEDs for infants younger than 1 year of age. Refer to your local protocols for the use of an AED on an infant.

Dose: The AED will automatically select the energy dose for each defibrillation. Some devices are equipped with pediatric systems that include a pad–cable system or a key to reduce the delivered energy to a suitable dose for children.

Procedure

1. Verify that the patient is in cardiac arrest, with no pulse or respiration. Have someone provide CPR while the AED is obtained and placed next to the patient.
2. Power on the AED. Follow the voice prompts and visual messages.
3. Open the package of adhesive electrode pads and attach pads to the patient’s bare chest.
4. Use adult pads for an adult and child pads for a child. If there are no child pads available, you may use adult pads on a child, but be sure the pads do not touch.
5. Attach one pad to the right sternal border (superior–anterior right chest) and place the second pad over the left apex (inferior–lateral left chest). Alternatively, follow the diagrams on each of the AED electrodes.
6. Attach the pads to the patient cables.
7. Clear the patient and stop CPR.
8. The AED may automatically analyze the patient’s rhythm or may be equipped with an “analyze” button.

9. If a shock is advised, say, “I’m going to shock on three. One, I’m clear; two, you’re clear; three, everybody’s clear.” Perform a visual sweep to ensure rescue personnel are not touching the patient or equipment. Press the shock button.
10. Once the shock is delivered, continue CPR beginning with chest compression.
11. After about 2 minutes of CPR the AED will prompt you with further verbal and visual cues.



Clinical Tip:

A fully automated AED analyzes the rhythm and delivers a shock, if one is indicated, without operator intervention.



Clinical Tip:

A semiautomated AED analyzes the rhythm and tells the operator that a shock is indicated. If it is, the operator initiates the shock.

CARDIOVERSION (Synchronized)

Indications: Unstable tachycardias with a perfusing rhythm. The patient may present with an altered LOC, dizziness, chest pain, or hypotension.

Energy Levels: With monophasic equipment, use energy levels of 100 J, 200 J, 300 J, and 360 J. Biphasic equipment supports an initial dose of 100–120 J with escalation as needed.

Application: Use handheld paddles or remote adhesive pads. Always use a conducting gel with paddles. For conscious patients explain the procedure and use a medication for sedation. Consider 2.5–5.0 mg of midazolam (**Versed**) or 5.0 mg diazepam (**Valium**).

Methods: Place defibrillator in synchronized (sync) mode. Charge to appropriate level. Say, “I’m going to shock on three. One, I’m clear; two, you’re clear; three, everybody’s clear.” Perform a visual sweep and press the shock button. Reassess the patient and treat according to the appropriate advanced cardiac life support protocol.

Precautions: Reactivate the “sync” mode after each attempted cardioversion; defibrillators default to the unsynchronized mode. Place paddles and pads several inches away from an implanted pacemaker.



Clinical Tip:

The “sync” mode delivers energy just after the R wave to avoid stimulation during the refractory, or vulnerable, period of the cardiac cycle when a shock could potentially produce VF.

TRANSCUTANEOUS PACING

Indications: Symptomatic bradycardia (with a pulse) unresponsive to atropine, bradycardia with ventricular escape rhythms, symptomatic second-degree AV block Type II, or third-degree AV block.

Pacing Modes: *Demand-mode (synchronous)* pacemakers sense the patient's heart rate and pace only when the heart rate falls below the level set by the clinician. *Fixed-mode (asynchronous)* pacemakers cannot sense the heart rate and always operate at the rate set by the clinician. Rate selections vary between 30 and 180 bpm. Output is adjustable between 0 and 200 mA. Pulse duration varies from 20–40 ms.

Application: Pacemaker pads work most effectively if placed in an anterior-posterior position on the patient. For correct pad placement, see Figure D-2.

Contraindications: Not effective in VF, pulseless VT, or asystole.

Side Effects: Chest muscle contraction, burns, and chest discomfort.

Precautions: Make sure pads have good skin contact to achieve capture and avoid burns.

CAROTID SINUS MASSAGE (Vagal Maneuver)

Indications: Can increase vagal nerve stimulation and slow SVT, or even convert SVT to NSR, without severe hemodynamic compromise.




Fig. D.2 ■ Placement of anterior-posterior pacemaker pads.

Method: Place the patient in a supine position, head tilted to either side with the neck hyperextended. Place your index and middle fingers over the carotid artery just below the angle of the jaw, as high on the neck as possible. Massage the artery for 5–10 sec by pressing it firmly against the vertebral column and rubbing (Fig. D-3).

Contraindications: Unequal carotid pulses, carotid bruits, cervical spine injury, or history of cerebrovascular accident or carotid atherosclerosis.

Side Effects: Slow HR or AV block, PVCs, VT, VF, syncope, seizure, hypotension, nausea or vomiting, stroke.

Precautions: Be sure the patient is receiving oxygen and an IV is in place. Never massage both arteries simultaneously.



Clinical Tip:

Each carotid artery should be palpated and auscultated before the procedure to maintain safety measures.



Clinical Tip:

Alternate vagal maneuvers include encouraging the patient to cough, bear down, or hold his or her breath.




Fig. D.3 ■ Carotid sinus massage.



Abbreviations Used in Text

AED	Automated external defibrillator	IV	Intravenous
A-fib	Atrial fibrillation	LOC	Level of consciousness
A-flutter	Atrial flutter	MAT	Multifocal atrial tachycardia
AV	Atrioventricular	MI	Myocardial infarction
BBB	Bundle branch block	NSR	Normal sinus rhythm
BP	Blood pressure	PAC	Premature atrial contraction
bpm	Beats per minute	PAT	Paroxysmal atrial tachycardia
BUN	Blood urea nitrogen	PEA	Pulseless electrical activity
CHF	Congestive heart failure	PJC	Premature junctional contraction
CO	Cardiac output	PO	Per os (by mouth)
COPD	Chronic obstructive pulmonary disease	PSVT	Paroxysmal supraventricular tachycardia
CPR	Cardiopulmonary resuscitation	PVC	Premature ventricular contraction
EMD	Electromechanical dissociation	SA	Sinoatrial
ET	Endotracheal	SC	Subcutaneous
HTN	Hypertension	SVT	Supraventricular tachycardia
IHSS	Idiopathic hypertrophic subaortic stenosis	VF	Ventricular fibrillation
IM	Intramuscular	VT	Ventricular tachycardia
IO	Intraoral	WAP	Wandering atrial pacemaker
		WPW	Wolff-Parkinson-White (Syndrome)



Selected References

1. American Heart Association: Basic Life Support for Healthcare Providers (Student Manual), Dallas, TX, 2006.
2. American Heart Association: Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care (supplement to *Circulation* 112: 24, December 13, 2005).
3. Deglin, JH, Vallerand, AH: *Davis's Drug Guide for Nurses*, ed 9. F.A. Davis, Philadelphia, 2006.
4. Deglin, JH, Vallerand, AH: *Med Notes*. F.A. Davis, Philadelphia, 2004.
5. Jones, SA: *ECG Notes*. F.A. Davis, Philadelphia, 2005.
6. Jones, SA: Author's personal collection.
7. Myers, E: *RNotes*. F.A. Davis, Philadelphia, 2003.
8. Myers, E, Hopkins, T: *MedSurg Notes*. F.A. Davis, Philadelphia, 2004.
9. *Physicians' Desk Reference*, ed 60. Thomson Healthcare, Montvale, NJ, 2006.
10. Scanlon, VC, Sanders, T: *Essentials of Anatomy and Physiology*, ed 4. F.A. Davis, Philadelphia, 2003.
11. *Taber's Cyclopedic Medical Dictionary*, ed 20. F.A. Davis, Philadelphia, 2005.

Illustration Credits

The ECG strips in Chapters 3–9 and 11–16 are from Jones: Author's personal collection, with permission. The unnumbered figure in Appendix A is from Jones: *ECG Notes*. Philadelphia: F.A. Davis, 2005, p 115, with permission. The cover and, unit page photographs are from Jones: Author's personal collection, with permission.

Fig. 1.1 ■ From Jones: *ECG Notes*. Philadelphia: F.A. Davis, 2005; p 1, with permission.

Fig. 1.2 ■ From Scanlon and Sanders: *Essentials of Anatomy and Physiology*, ed 4. Philadelphia: F.A. Davis, 2005; p 260, with permission.

Fig. 1.3 ■ From Scanlon and Sanders: *Essentials of Anatomy and Physiology*, ed 4. Philadelphia: F.A. Davis, 2005; p 263, with permission.

Fig. 1.4 ■ From Scanlon and Sanders: *Essentials of Anatomy and Physiology*, ed 4. Philadelphia: F.A. Davis, 2005; p 262, with permission.

Fig. 1.5 ■ From Jones: *ECG Notes*. Philadelphia: F.A. Davis, 2005; p 5, with permission (adapted from Scanlon and Sanders: *Essentials of Anatomy and Physiology*, ed 4. Philadelphia: F.A. Davis, 2005; p 263.

Fig. 1.6 ■ From Scanlon and Sanders: *Essentials of Anatomy and Physiology*, ed 4. Philadelphia: F.A. Davis, 2005; p 279, with permission.

Fig. 1.7 ■ From Scanlon and Sanders: *Essentials of Anatomy and Physiology*, ed 4. Philadelphia: F.A. Davis, 2005; p 283, with permission.

Fig. 1.8 ■ From Scanlon and Sanders: *Essentials of Anatomy and Physiology*, ed 4. Philadelphia: F.A. Davis, 2005, p 284, with permission.

Fig. 1.10 ■ From Jones: *ECG Notes*. Philadelphia: F.A. Davis, 2005; p 10, with permission (adapted from Myers and Hopkins: *MedSurg Notes*. Philadelphia: F.A. Davis, 2004, p 184.

Fig. 2.1 ■ From Myers and Hopkins: *MedSurg Notes*. Philadelphia, F.A. Davis, 2004, p 186, with permission.

Fig. 2.2 ■ From Jones: *ECG Notes*. Philadelphia, F.A. Davis, 2005; p 13, with permission.

Fig. 2.3 ■ From Jones: *ECG Notes*. Philadelphia, F.A. Davis, 2005; p 14, with permission.

Fig. 2.4 ■ From Jones: *ECG Notes*. Philadelphia, F.A. Davis, 2005; p 15, with permission.

Fig. 2.5 ■ From Myers and Hopkins: *MedSurg Notes*. Philadelphia, F.A. Davis, 2004; p 185, with permission.

Fig. 2.6 ■ From Jones: *ECG Notes*. Philadelphia, F.A. Davis, 2005; p 17, with permission.

Fig. 2.7 ■ From Jones: *ECG Notes*. Philadelphia, F.A. Davis, 2005; p 17, with permission.

Fig. 2.8 ■ From Jones: *ECG Notes*. Philadelphia, F.A. Davis, 2005; p 18, with permission.

Fig. 2.9 ■ From Jones: *ECG Notes*. Philadelphia, F.A. Davis, 2005; p 18, with permission.

Fig. 2.10 ■ From Jones: *ECG Notes*. Philadelphia, F.A. Davis, 2005; p 19, with permission (adapted from Myers and Hopkins: *MedSurg Notes*. Philadelphia, F.A. Davis, 2004; p 185).

Fig. 2.11 ■ From Jones: *ECG Notes*. Philadelphia, F.A. Davis, 2005; p 20, with permission.

Fig. 2.12 ■ From Jones: *ECG Notes*. Philadelphia, F.A. Davis, 2005; p 21, with permission.

Fig. 2.13 ■ From Jones: *ECG Notes*. Philadelphia, F.A. Davis, 2005; p 21, with permission.

Fig. 2.14 ■ From Jones: *ECG Notes*. Philadelphia, F.A. Davis, 2005; p 23, with permission.

Fig. 2.15 ■ From Jones: Author's personal collection, with permission.

Fig. 10.1 ■ From Myers and Hopkins: *MedSurg Notes*. Philadelphia: F.A. Davis, 2004; p 185, with permission.

Fig. 10.2 ■ From Jones: *ECG Notes*. Philadelphia, F.A. Davis, 2005; p 74, with permission.

Fig. 10.3 ■ From Jones: *ECG Notes*. Philadelphia, F.A. Davis, 2005; p 75, with permission.

Fig. 10.4 ■ From Jones: *ECG Notes*. Philadelphia, F.A. Davis, 2005; p 75, with permission.

Fig. 10.5 ■ From Jones: *ECG Notes*. Philadelphia, F.A. Davis, 2005; p 75, with permission.

Fig. 10.6 ■ From Jones: *ECG Notes*. Philadelphia, F.A. Davis, 2005; p 76, with permission.

Fig. 10.7 ■ From Jones: *ECG Notes*. Philadelphia, F.A. Davis, 2005, p 77, with permission.

Fig. 10.8 ■ From Jones: *ECG Notes*. Philadelphia, F.A. Davis, 2005, p 77, with permission.

Fig. 10.9 ■ From Jones: *ECG Notes*. Philadelphia, F.A. Davis, 2005; p 77, with permission.

Fig. 10.10 ■ From Jones: *ECG Notes*. Philadelphia, F.A. Davis, 2005; p 78, with permission.

Fig. 10.11 ■ From Jones: *ECG Notes*. Philadelphia, F.A. Davis, 2005; p 79, with permission.

Fig. 10.12 ■ From Jones: *ECG Notes*. Philadelphia, F.A. Davis, 2005; p 80, with permission.

Fig. 10.13 ■ From Jones: *ECG Notes*. Philadelphia, F.A. Davis, 2005; p 81, with permission.

Fig. 10.14 ■ From Jones: *ECG Notes*. Philadelphia, F.A. Davis, 2005; p 82, with permission.

Fig. 10.15 ■ From Jones: *ECG Notes*. Philadelphia, F.A. Davis, 2005; p 83, with permission.

Fig. 10.16 ■ From Jones: *ECG Notes*. Philadelphia, F.A. Davis, 2005; p 84, with permission.

Fig. 10.17 ■ From Jones: *ECG Notes*. Philadelphia, F.A. Davis, 2005; p 85, with permission.

Fig. A.1 ■ From Jones: *ECG Notes*. Philadelphia: F.A. Davis, 2005, p 110, with permission.

Fig. A.2 ■ From Jones: *ECG Notes*. Philadelphia: F.A. Davis, 2005; p 111, with permission (adapted from Myers and Hopkins: *MedSurg Notes*. Philadelphia: F.A. Davis, 2004, p 11).

Fig. D.1 ■ From Myers and Hopkins: *MedSurg Notes*. Philadelphia: F.A. Davis, 2004, p 15, with permission.

Fig. D.2 ■ From Jones: *ECG Notes*. Philadelphia: F.A. Davis, 2005, p 103, with permission.

Fig. D.3 ■ From Jones: *ECG Notes*. Philadelphia: F.A. Davis, 2005, p 105, with permission.



Index

Note: Page numbers in **boldface** refer to definition of rhythms; page numbers followed by f refer to figures and rhythm strips; page numbers followed by t refer to tables.

A

- A-fib. *See* Atrial fibrillation (A-fib)
- A-flutter. *See* Atrial flutter (A-flutter)
- Abbreviations, 268
- Accelerated idioventricular rhythm, **45**, **45f**, 52f, 90f, 115, 130f, 142f, 149, 152, 170f, 183
 - bigeminal premature ventricular contraction with, 145f, 153
 - sinus rhythm conversion to, 170f, 183
- Accelerated junctional rhythm, **38**, **38f**, 39f, 41f, 93f, 116
 - case study of, 226–228, 226f, 227f, 228f, 231
 - inverted T waves with, 131f, 149, 199f, 214
 - paroxysmal supraventricular tachycardia with, 143f, 152
 - ST segment depression with, 131f, 149, 199f, 214
 - ST segment elevation with, 99f, 117
- Accelerated ventricular rhythm
 - atrial flutter with, 141f, 152
 - unknown underlying rhythm with, 141f, 152
- ACE (angiotensin-converting enzyme) inhibitors, 259
- Acetylsalicylic acid, 260
- Acute coronary syndrome, advanced cardiac life support protocol for, 255
- Adenocard, 259
- Adenoscan, 259
- Adenosine, 259
- Adrenalin, 261
- Adrenergic agonists, 261
- Advanced cardiac life support protocols, 253–258
 - for acute coronary syndrome, 255
 - for asystole, 254–255
 - for bradycardia, 255–256
 - for narrow-complex tachycardia, 257
 - for pulseless electrical activity, 254
 - for pulseless ventricular tachycardia, 253–254
 - for unstable tachycardia, 256–257
 - for ventricular fibrillation, 253–254
 - for wide-complex tachycardia, 257–258
- AED (automated external defibrillator), 266
- Afterload, 6
- Agonal rhythm, **44**, 192f, 211f, 212, 217
- Airway obstruction. *See also* Cardiopulmonary resuscitation (CPR)
 - case study of, 232, 242–243
 - in conscious adult or child, 249–250, 250f
 - in conscious infant, 250, 250f
 - in unconscious adult, 250–251
 - in unconscious child, 251
 - in unconscious infant, 251–252
- Alkalinizing agents, 263
- Alteplase, 261
- Amiodarone, 258–259
- Analgesics, 262–263
- Anesthetics, 262
- Angiotensin-converting enzyme (ACE) inhibitors, 259
- Anistreplase, 261
- Anterior myocardial infarction, 78f, 79f, **80**, **80f**
- Antianginals, 263
- Antiarrhythmics, 259–260, 262, 263
- Anticholinergics, 260
- Antihypertensives, 259
- Antiplatelets, 260
- Aorta, 3, 4f
- Arrhythmia(s). *See also specific arrhythmias*
 - atrial, 28–36, 28f–35f
 - junctional, 37–43, 37f–42f
 - sinoatrial node, 20–27, 20f–26f
 - ventricular, 44–55, 44f–54f
 - Arteries, anatomy of, 4, 4f, 5f
- Artifact(s), 71–76
 - loose connection, **71**, **71f**, 74f, 155f, 179
 - muscle, **72**, **72f**
 - atrial fibrillation with, 75f
 - junctional rhythm with, 189f, 212
 - sinus bradycardia with, 74f
 - sinus rhythm with, 73f, 97f, 109f, 117, 120, 171f, 176f, 183, 184, 188f, 211
 - torsade de pointes with, 75f
 - respiration, **71**, **71f**
 - sinus arrhythmia with, 89f, 115
 - sinus bradycardia with, 97f, 117
 - sinus pause with, 135f, 150
 - ventricular pacemaker with, 135f, 150
 - sinus tachycardia with, 109f, 120
 - sixty-cycle interference, **72**, **72f**
 - atrial fibrillation with, 73f
 - sinus rhythm with, 138f, 151
 - torsade de pointes with, 74f
 - vs. ventricular tachycardia, 191f, 212
- Artificial cardiac pacemaker, 63–70
 - atrial, **65f**, 67f, 87f, 113f, 115, 121, 164f, 181
 - failure to capture of, 96f, 117, 135f, 150
 - first-degree atrioventricular block with, 174f, 184
 - inverted T waves with, 123f, 147
 - nonpaced P wave with, 68f
 - premature ventricular contraction with, 176f, 184
 - ST segment depression with, 123f, 147
 - atrial and ventricular, **65f**, 67f, 104f, 114f, 119, 121, 193f, 213
 - absent atrial conduction with, 69f
 - bundle branch block with, 68f
 - failure to capture of, 107f, 119, 163f, 181

- Artificial cardiac pacemaker (*Continued*)
- premature junctional contraction with, 206f, 216
 - premature ventricular contraction with, 203f, 215
 - ST segment depression with, 203f, 215
 - in capture, 65
 - codes for, 64, 64t
 - demand (synchronous), 64
 - dual-chamber, 64, 65f
 - failure to capture of, **66f, 66t**, 67f, 96f, 107f, 117, 119, 135f, 150, 163f, 181
 - asystole with, 186f, 211
 - failure to pace of, 66t, 68f
 - failure to sense of, **66f, 66t**
 - fixed rate (asynchronous), 64
 - inverted T wave with, 203f, 215
 - loss of capture of, 91f, 116
 - malfunctions of, 65–66, 66f, 66t
 - modes of, 64
 - oversensing of, **66f, 66t**
 - P-R interval with, 64
 - P wave with, 64
 - permanent, 63–64
 - premature ventricular contraction with, 67, 195f, 203f, 213, 215
 - QRS interval with, 64
 - rate of, 64
 - rhythm of, 64, 65f
 - single-chamber, 63, 65f
 - temporary, 63
 - transcutaneous, 267, 267f
 - ventricular, **65f**, 67f, 69f, 158f, 161f, 178f, 180, 181, 185
 - conducted sinus complex with, 109f, 120
 - fusion beat with, 210f, 217
 - loss of capture of, 91f, 116
 - monomorphic ventricular tachycardia with, 195f, 213
 - no pacing with, 68f
 - oversensing of, 197f, 214
 - polymorphic ventricular tachycardia with, 173f, 184
 - premature ventricular contraction with, 195f, 213
 - respiration artifact with, 135f, 150
 - sinus bradycardia with, 171f, 183
 - wide intrinsic complexes with, 210f, 217
- Aspirin, 260
- Asystole, **51, 51f**, 54f, 106f, 119, 166f, 171f, 182, 183
 - advanced cardiac life support protocol for, 254–255
 - case study of, 235–237, 235f, 236f, 237f, 243–244
 - pacemaker failure to capture with, 186f, 211
 - ventricular, 207f, 216
 - ventricular fibrillation conversion from, 156f, 179, 190f, 212
 - ventricular fibrillation conversion to, 126f, 129f, 148, 149
- Atenolol, 260
- Atria, 3, 3f
- Atrial arrhythmias, 28–36, 28f–35f. *See also specific arrhythmias*
- Atrial fibrillation (A-fib), **32, 32f**, 34f, 35f, 140f, 151, 166f, 182, 199f, 214
 - atrial flutter conversion from, 162f, 181
 - atrial flutter conversion to, 102f, 118
 - bundle branch block with, 96f, 112f, 117, 120, 127f, 148, 193f, 202f, 213, 215
 - controlled ventricular response with, 202f, 215
 - couplet premature ventricular contraction with, 158f, 180
 - inverted T waves with, 134f, 150
 - multiform ventricular tachycardia with, 136f, 150
 - muscle artifact with, 75f
 - premature ventricular contraction with, 129f, 149
 - rapid ventricular response with, 103f, 107f, 118, 119, 146f, 153, 154f, 155f, 179, 187f, 190f, 211, 212
 - sixty-cycle interference with, 73f
 - slow ventricular response with, 162f, 181, 193f, 213
 - ST segment depression with, 92f, 98f, 116, 117, 134f, 150, 155f, 160f, 179, 180, 190f, 212
 - ST segment elevation with, 158f, 180
- Atrial flutter (A-flutter), **31, 31f**, 33f, 35f
 - 2:1 block with, 165f, 182
 - 4:1 block with, 122f, 133f, 147, 150
 - 8:1 block with, 157f, 180
 - 3:1 conduction with, 204f, 215
 - accelerated ventricular rhythm with, 141f, 152
 - atrial fibrillation conversion from, 102f, 118
 - atrial fibrillation conversion to, 162f, 181
 - bundle branch block with, 136f, 150
 - couplet premature ventricular contraction with, 127f, 148
 - slow ventricular response with, 198f, 214
 - variable block with, 100f, 118, 136f, 146f, 150, 153, 188f, 198f, 211, 214
- Atrial pacemaker, wandering, **28, 28f**, 107f, 119, 172f, 175f, 183, 184
- Atrial quadrigeminy, sinus tachycardia with, 173f, 184
- Atrial tachycardia, **30, 30f**, 33f, 138f, 151
 - multifocal, **29, 29f**
 - premature ventricular contraction with, 53f
- Atrioventricular (AV) block
 - first-degree, **56, 56f**, 59f, 61f, 93f, 116
 - atrial pacemaker with, 174f, 184
 - normal sinus rhythm with, 92f, 103f, 116, 118, 144f, 152
 - sinus bradycardia with, 156f, 163f, 172f, 179, 181, 183, 208f, 216
 - sinus rhythm with, 131f, 149
 - second-degree
 - type I (Mobitz I, Wenckebach), **57, 57f**, 59f, 60f, 90f, 93f, 115, 116, 177f, 185
 - type II (Mobitz II), **57, 57f**, 59f, 60f, 191f, 212
 - 3:1 block with, 143f, 152, 209f, 217
 - bundle branch block with, 209f, 217
 - third-degree, **58, 58f**, 60f, 61f, 95f, 100f, 117, 118, 159f, 165f, 168f, 180, 182, 193f, 207f, 208f, 213, 216
 - case study of, 241–242, 241f, 242f, 244–245
 - couplet premature ventricular contraction with, 110f, 120
 - multiform premature ventricular contraction with, 147f, 153
 - ventricular fibrillation conversion from, 139f, 151
 - ventricular fibrillation conversion to, 156f, 179
- Atrioventricular node, 7, 7f
 - intrinsic rate of, 63
- Atrioventricular valves, 3, 3f
- Atropine sulfate, 260
- Automated external defibrillator (AED), 266
- Automaticity, 6–7
- AV. *See at* Atrioventricular
- Axillary artery, 4, 5f
- Axillary vein, 5, 5f
- Axis deviation, 78, 78f
- B**
- BBB. *See* Bundle branch block (BBB)
- Beta-adrenergic agonists, 262
- Beta blockers, 260
- Bigeminy
 - atrial, sinus rhythm with, 179f, 185
 - junctional
 - junctional rhythm with, 160f, 180
 - sinus rhythm with, 125f, 148, 205f, 210f, 216, 217
 - ventricular, **46, 46f**
 - accelerated idioventricular rhythm with, 145f, 153
 - normal sinus rhythm with, 165f, 182, 189f, 202f, 212, 215
 - sinus rhythm with, 138f, 151
- Block
 - 2:1, atrial flutter with, 165f, 182

- 3:1, type II second-degree atrioventricular block with, 143f, 152, 209f, 217
- 4:1, atrial flutter with, 122f, 133f, 147, 150
- atrioventricular. *See* Atrioventricular (AV) block
- bundle branch. *See* Bundle branch block (BBB)
- sinoatrial, **23, 23f**, 26f
- sinus rhythm with, 111f, 120, 201f, 215
 - sinus tachycardia with, 192f, 212
- sinus, sinus rhythm with, 94f, 116
- variable, atrial flutter with, 100f, 118, 136f, 146f, 150, 153, 188f, 198f, 211, 214
- Blood flow, cardiac, 5–6, 6f
- Brachial artery, 4, 5f
- Bradycardia. *See* Sinus bradycardia
- Bundle branch
- left, 7, 7f
 - right, 7, 7f
- Bundle branch block (BBB), **58, 58f**, 61f
- atrial fibrillation with, 96f, 112f, 117, 120, 127f, 148, 193f, 202f, 213, 215
 - atrial flutter with, 136f, 150
 - junctional rhythm with, 89f, 115, 160f, 167f, 180, 182
 - junctional tachycardia with, 137f, 151
 - left, **82, 82f**
 - normal sinus rhythm with, 95f, 117, 128f, 140f, 148, 151, 158f, 160f, 167f, 177f, 180, 182, 185, 196f, 201f, 210f, 213, 215, 217
 - on pacemaker electrocardiogram, 68f
 - right, **82, 83f**
 - sinus bradycardia with, 172f, 183, 194f, 200f, 213, 214
 - type II second-degree atrioventricular block with, 209f, 217
- Bundle of His, 7, 7f
- C**
- Calan, 264
- Calcium channel blockers, 261, 264
- Calcium chloride, 260
- Captopril, 259
- Cardiac cycle, 5–6
- on electrocardiogram, 14, 14f
- Cardiac output, 6
- Cardiopulmonary resuscitation (CPR). *See also* Airway obstruction
- adult
- case study of
 - for accelerated junctional rhythm with pulseless electrical activity, 226–228, 226f, 227f, 228f, 231
 - for asystole, 235–237, 235f, 236f, 237f, 243–244
 - for sinus rhythm with ST segment elevation, 239–241, 239f, 240f, 241f, 244
 - for third-degree atrioventricular block, 241–242, 241f, 242f, 244–245
 - for ventricular tachycardia, 223–226, 224f, 225f, 226f, 230, 237–239, 237f, 238f, 239f, 244
 - guidelines for, 247–248
- child (1 year to 14 years)
- case study of, 222–223, 229–230
 - guidelines for, 247, 248–249, 251
- infant (younger than 1 year), guidelines for, 247, 249, 251–252
- positions for, 252
- Cardiovascular system, 4–5, 5f
- Cardioversion, 266
- Cardizem, 261
- Carotid artery, 4, 5f
- Carotid sinus massage, 267, 267f
- Case study
- adult CPR
 - for accelerated junctional rhythm with pulseless electrical activity, 226–228, 226f, 227f, 228f, 231
 - for asystole, 235–237, 235f, 236f, 237f, 243–244
 - for sinus rhythm with ST segment elevation, 239–241, 239f, 240f, 241f, 244
 - for third-degree atrioventricular block, 241–242, 241f, 242f, 244–245
 - for ventricular tachycardia, 223–226, 224f, 225f, 226f, 230, 237–239, 237f, 238f, 239f, 244
 - pediatric CPR, 222–223, 229–230
 - sinus bradycardia, 228–229, 228f, 229f, 231
 - ventricular fibrillation, 220–222, 220f, 221f, 222f, 229
- Children (1 year to 14 years)
- airway obstruction in, 249–250, 250f, 251
 - cardiopulmonary resuscitation for
 - case study of, 222–223, 229–230
 - guidelines for, 247, 248–249
 - Heimlich maneuver for, 249–250, 250f
- Choking
- case study of, 232, 242–243
 - Heimlich maneuver for
 - for adult or child, 249–250, 250f
 - for infant, 250, 250f
 - universal sign for, 252
- Complex, 14, 14f
- QRS, 14, 14f, 17
- Conductivity, 7, 7f, 8f
- Contractions. *See* Premature atrial contraction (PAC); Premature junctional contraction (PJC); Premature ventricular contraction (PVC)
- Cordarone, 258–259
- Coronary arteries, 3–4, 4f
- blockage of. *See* Myocardial infarction (MI)
- Coronary syndrome, acute, advanced cardiac life support protocol for, 255
- Coronary veins, 4, 4f
- Couplet premature ventricular contractions, **47, 47f**
- atrial fibrillation with, 158f, 180
 - atrial flutter with, 127f, 148
 - normal sinus rhythm with, 167f, 182
 - sinus tachycardia with, 92f, 116, 145f, 153
 - third-degree atrioventricular block with, 110f, 120
- Covert, 262
- CPR. *See* Cardiopulmonary resuscitation (CPR)
- D**
- Defibrillation, 103f, 118, 265–266, 265f
- Depolarization, 7, 8f
- on electrocardiogram, 9, 9f
- Diastole, 5–6, 6f
- Digibind, 261
- Digoxin, 260
- Digoxin immune fab, 261
- Diltiazem, 261
- Diuretics, 261–262
- Dopamine, 261
- Dropped beats, 16t, 23, 23f
- E**
- Electrical axis deviation, 78, 78f
- Electrocardiogram, 9–17
- 12-lead, 77, **80, 80f**
 - 15-lead, 13, 13f, 13t
 - analysis of, 16–17, 16t
 - arrhythmias on. *See* Arrhythmia(s) and specific patterns
 - artifacts on. *See* Artifact(s)
 - artificial pacemaker. *See* Artificial cardiac pacemaker

Electrocardiogram (*Continued*)

- augmented leads for, 10, 11f, 11t
- cable connections for, 11, 11t
- cardiac cycle on, 14, 14f
- chest leads for, 10, 11f, 11t, 12, 12f
- components of, 14, 14f
- definition of, 9–10
- depolarization on, 9, 9f
- electrode placement for, 11–12, 11f, 12f
- five-wire cable for, 12, 12f
- graph paper for, 13, 14f
- heart rate calculation with, 14–16, 15f, 15t
- leads for, 9–10, 10f, 11f
- limb leads for, 10, 10f
- modified chest leads for, 12, 12f
- monitoring lead of, 12
- myocardial infarction on. *See* Myocardial infarction (MI)
- R wave progression on, 78, 78f
- repolarization on, 9, 9f
- rhythm strip of, 13, 14f
- right-sided, 12–13, 13f, 13t
- six-second, 15, 15f
- standard leads for, 10, 10f, 10t
- three-wire cable for, 11–12, 11f
- troubleshooting for, 77–78

Electrodes

- loose, **71**, **71f**, 74f, 155f, 179
- placement of, 11–12, 11f, 12f

Electromechanical dissociation (pulseless electrical activity), **50–51**, **50f**

- advanced cardiac life support protocol for, 254
- case study of, 226–228, 226f, 227f, 228f, 231

Emergency medications, 259–264

- adrenergic agonist, 261
- alkalizing, 263
- analgesic, 262–263
- anesthetic, 262
- antianginal, 263
- antiarrhythmic, 259–260, 262, 263
- anticholinergic, 260
- antihypertensive, 259, 260
- antiplatelet, 260
- beta-adrenergic agonist, 262
- calcium channel blocker, 261, 264
- diuretic, 261–262
- fibrinolytic, 261
- inotropic, 260, 261
- sympathomimetic, 262
- thrombolytic, 261
- vasopressor, 261, 264

Enalapril, 259

Endocardium, 2, 3f

Epicardium, 2, 3f

Epinephrine, 261

Escape beat, junctional, **38**, **38f**

- normal sinus rhythm with, 174f, 184
- sinus rhythm with, 164f, 181

Esmolol, 260

Excitability, 7

FFailure to capture, **66f**, **66t**, 67f, 150

- asystole with, 186f, 211
- atrial and ventricular pacemaker, 107f, 119, 163f, 181
- atrial pacemaker, 96f, 117, 135f, 150

Failure to sense, **66f**, **66t**

Femoral artery, 4, 5f

Femoral vein, 5, 5f

Fibrillation. *See* Atrial fibrillation (A-fib); Ventricular fibrillation (VF)

Fibrinolytics, 261

Fifteen-lead electrocardiogram, 13, 13f, 13t

First-degree atrioventricular (AV) block, **56**, **56f**, 59f, 61f, 93f, 116

- atrial pacemaker with, 174f, 184
- sinus bradycardia with, 156f, 163f, 172f, 179, 181, 183, 208f, 216
- sinus rhythm with, 92f, 103f, 116, 118, 131f, 144f, 149, 152

Flutter. *See* Atrial flutter (A-flutter)

Furosemide, 261–262

Fusion beat, artificial pacemaker with, 210f, 217

G

Graph paper, 13, 14f

Great vessels, 3, 3f

H

Heart

- anatomy of, 2–4, 2f, 3f
- anterior wall of, 78f
- atria of, 3, 3f
- automaticity of, 6–7
- blood flow through, 5–6, 6f
- chambers of, 3, 3f
- conduction system of, 7, 7f, 8f
- contractile state of, 6
- electrical axis deviation of, 78, 78f
- electrophysiology of, 6–8, 7f, 8f
- excitability of, 7, 7f, 8f
- inferior wall of, 79f
- ischemia of, 78, 79f. *See also* Myocardial infarction (MI)
- lateral wall of, 79f
- layers of, 2, 3f
- mechanical physiology of, 5–6, 6f
- physiology of, 5–8, 6f, 8f
- septal wall of, 79f
- shape of, 2, 2f
- valves of, 2–3, 3f
- ventricles of, 3, 3f

Heart failure, 63

Heart rate, 63

- calculation of, 14–16, 15f, 15t, 17–18
- cardiac output and, 6

Heimlich maneuver, 232, 242

- for adult or child, 249–250, 250f
- for infant, 250, 250f

I

Ibutilide, 262

Idioventricular rhythm, **44**, **44f**, 54f, 163f, 177f, 181, 185, 192f, 211f, 212, 217

- accelerated, **45**, **45f**, 52f, 90f, 115, 130f, 142f, 149, 152, 170f, 183

- bigeminal premature ventricular contraction with, 145f, 153
- normal sinus rhythm conversion to, 170f, 183

Iliac artery, 4, 5f

Iliac vein, 5, 5f

Infant

- airway obstruction in, 250, 250f, 251–252
- cardiopulmonary resuscitation for, 247, 249
- Heimlich maneuver for, 250, 250f

Infarction. *See* Myocardial infarction (MI)Inferior myocardial infarction, 79f, **80–81**, **81f**

Inferior vena cava, 5, 5f

Injury, 78, 79f. *See also* Myocardial infarction (MI)

Inotropic agents, 260, 261

Interatrial septum, 3

Internodal pathways, 7, 7f

Interpolated premature ventricular contraction, **48**, **48f**

- normal sinus rhythm with, 142f, 144f, 152
- sinus bradycardia with, 97f, 117, 169f, 183, 205f, 216

- Interval, 14, 14f
 PR, 14, 14f, 16t
 artificial pacemaker with, 64
 QRS, 16t
 artificial pacemaker with, 64
 QT, 14, 14f, 16t
 R-R, 16
 measurement of, 17
- Interventricular septum, 3, 3f
- Intropin, 261
- Ischemia, 78, 79f. *See also* Myocardial infarction (MI)
- Isoelectric line, 14, 14f
- Isoproterenol, 262
- Isoptin, 264
- Isuprel, 262
- J**
- Jugular vein, 5, 5f
- Junctional arrhythmias, 37–43, 37f–42f. *See also specific arrhythmias*
- Junctional contraction, premature. *See* Premature junctional contraction (PJC)
- Junctional escape beat, **38, 38f**
 sinus rhythm with, 164f, 174f, 181, 184
- Junctional rhythm, **37, 37f**, 40f, 41f, 159f, 180, 204f, 215
 accelerated, **38, 38f**, 39f, 41f, 93f, 116
 case study of, 226–228, 226f, 227f, 228f, 231
 inverted T waves with, 131f, 149, 199f, 214
 paroxysmal supraventricular tachycardia with, 143f, 152
 ST segment depression with, 131f, 149, 199f, 214
 ST segment elevation with, 99f, 117
 bigeminal premature junctional contraction with, 160f, 180
 bundle branch block with, 89f, 115, 160f, 167f, 180, 182
 interpolated premature junctional contraction with, 167f, 182
 muscle artifact with, 189f, 212
 premature junctional contraction with, 142f, 152
 premature ventricular contraction with, 189f, 212
 ST segment depression with, 134f, 142f, 150, 152, 207f, 216
 supraventricular tachycardia conversion with, 125f, 148
- Junctional tachycardia, **38, 38f**, 40f, 41f, 42f, 168f, 182
 bundle branch block with, 137f, 151
 couplet premature ventricular contraction with, 187f, 211
 inverted T waves with, 128f, 148
- L**
- Labetalol, 260
- Lanoxin, 260
- Lasix, 261–262
- Lateral myocardial infarction, 79f, **81, 81f**
- Lead(s), 9–10, 10f, 11f
 augmented, 10, 11f, 11t
 chest, 10, 11f, 11t, 12, 12f
 limb, 10, 10f
 monitoring, 12
 standard, 10, 10f, 10t
- Left bundle branch block, **82, 82f**. *See also* Bundle branch block (BBB)
- Lidocaine, 262
- Lisinopril, 259
- Loop diuretics, 261–262
- Loose connection artifact, **71, 71f**, 74f, 155f, 179
- M**
- Magnesium sulfate, 262
- MAT (multifocal atrial tachycardia), **29, 29f**
- Metoprolol tartrate, 260
- MI. *See* Myocardial infarction (MI)
- Mitral valve, 3, 3f
- Mobitz I atrioventricular (AV) block, **57, 57f**, 59f, 60f, 90f, 93f, 115, 116, 177f, 185
- Mobitz II atrioventricular (AV) block, **57, 57f**, 59f, 60f, 191f, 212
 3:1 block with, 143f, 152, 209f, 217
 bundle branch block with, 209f, 217
- Monomorphic ventricular tachycardia, **48, 48f**, 51f, 52f, 106f, 119, 144f, 152, 198f, 214
 sinus bradycardia conversion to, 145f, 153
- Morphine sulfate, 262–263
- Multifocal atrial tachycardia (MAT), **29, 29f**
- Multiform premature ventricular contraction, **46f**
 normal sinus rhythm with, 88f, 94f, 105f, 115, 116, 119
 sinus rhythm with, 52f, 168f, 182
 third-degree atrioventricular block with, 147f, 153
- Muscle artifact, **72, 72f**
 junctional rhythm with, 189f, 212
 normal sinus rhythm with, 97f, 109f, 117, 120, 176f, 184, 188f, 211
 sinus bradycardia with, 74f
 sinus rhythm with, 72f, 73f, 171f, 183
 torsade de pointes with, 75f
- Myocardial infarction (MI), 78–83
 advanced cardiac life support protocol for, 255
 anterior, 78f, 79f, **80, 80f**
 inferior, 79f, **80–81, 81f**
 lateral, 79f, **81, 81f**
 location of, 78–79, 78f, 79f
 posterior, 79f, **82, 82f**
 progression of, 79, 79f
 septal, 79f, **81–82, 81f**
 ST segment elevation in, 79–80, 80f
- Myocardium, 2, 3f
 infarction of. *See* Myocardial infarction (MI)
- N**
- Nitroglycerin, 263
- Nitrostat, 263
- Normal sinus rhythm, **20, 20f**, 26f, 27, 91f, 116, 146f, 153, 178f, 185, 199f, 214
 accelerated idioventricular rhythm conversion from, 170f, 183
 bigeminal multiform pre-ventricular contraction with, 194f, 213
 bigeminal premature atrial contraction with, 179f, 185
 bigeminal premature junctional contraction with, 125f, 148, 205f, 210f, 216, 217
 bigeminal premature ventricular contraction with, 128f, 148, 165f, 182, 202f, 215
 bundle branch block with, 95f, 117, 128f, 140f, 148, 151, 158f, 160f, 167f, 177f, 180, 182, 185, 196f, 201f, 210f, 213, 215, 217
 couplet premature ventricular contraction with, 167f, 182
 first-degree atrioventricular block with, 92f, 103f, 116, 118, 144f, 152
 interpolated premature ventricular contraction with, 142f, 144f, 152
 inverted T waves with, 99f, 118, 144f, 152, 201f, 210f, 215, 217
 junctional escape beat with, 174f, 184
 multiform premature ventricular contraction with, 88f, 94f, 105f, 115, 116, 119
 muscle artifact with, 97f, 109f, 117, 120, 176f, 184, 188f, 211
 paroxysmal supraventricular tachycardia conversion from, 155f, 179
 paroxysmal supraventricular tachycardia conversion to, 104f, 119, 235, 235f
 peaked T waves with, 99f, 118
 premature atrial contraction with, 98f, 105f, 117, 119, 140f, 151, 179f, 185
 premature junctional contraction with, 195f, 213
 premature ventricular contraction with, 93f, 94f, 100f, 116, 118, 132f, 149, 164f, 170f, 181, 183, 195f, 202f, 213, 215
 sinoatrial block with, 201f, 215
 sinus block with, 94f, 116

Normal sinus rhythm (*Continued*)

- sinus bradycardia with, 200f, 214
- sinus pause with, 138f, 151, 162f, 174f, 181, 184
- ST segment depression with, 132f, 139f, 151, 169f, 183, 196f, 201f, 202f, 210f, 213, 215, 217
- ST segment elevation with, 99f, 118, 134f, 150, 186f, 192f, 206f, 211, 212, 216
- supraventricular tachycardia conversion from, 203f, 215
- supraventricular tachycardia conversion to, 133f, 150
- triplet premature ventricular contraction with, 123f, 134f, 147, 150, 166f, 176f, 182, 184, 201f, 215
- U wave with, 24f, 26, 100f, 118, 161f, 181
- ventricular bigeminy with, 165f, 182, 189f, 202f, 212, 215
- ventricular fibrillation conversion to, 195f, 213
- ventricular quadrigeminy with, 197f, 214
- ventricular tachycardia conversion to, 137f, 151, 225–226, 226f
- ventricular trigeminy with, 158f, 161f, 175f, 180, 181, 184

O

Oxygen therapy, 263

P

P wave, 14, 14f, 16t, 17

with artificial cardiac pacemaker, 64

Pacemaker

artificial. *See* Artificial cardiac pacemaker

atrial, wandering, **28, 28f**, 107f, 119, 172f, 175f, 183, 184

sinoatrial, 7, 7f, 63

Pacerone, 258–259

Paroxysmal supraventricular tachycardia (PSVT), **31, 31f**, 34f, 35f

accelerated junctional rhythm with, 143f, 152

case study of, 233–235, 233f, 234f, 235f, 243

normal sinus rhythm conversion from, 104f, 119, 235, 235f

normal sinus rhythm conversion to, 155f, 179

sinus tachycardia conversion to, 169f, 183

Pause, 16t

sinus (sinus arrest), **22, 22f**, 25f, 102f, 118, 123f, 147

respiration artifact with, 135f, 150

sinus bradycardia with, 126f, 148

sinus rhythm with, 138f, 151, 162f, 174f, 181, 184

ventricular tachycardia with, 206f, 216

PEA. *See* Pulseless electrical activity (PEA)

Pericardial cavity, 2, 3f

Pericardium, 2, 3f

Pitressin, 264

PJC. *See* Premature junctional contraction (PJC)

Polymorphic ventricular tachycardia, **49, 49f**, 113f, 121, 136f, 150

sinus tachycardia with, 188f, 211

ventricular pacemaker with, 173f, 184

vs. artifact, 108f, 120

Posterior myocardial infarction, 79f, **82, 82f**

PR interval, 14, 14f, 16t

with artificial cardiac pacemaker, 64

Preload, 6

Premature atrial contraction (PAC), **29, 29f**, 34f

bigeminal, sinus rhythm with, 179f, 185

normal sinus rhythm with, 98f, 105f, 117, 119, 140f, 151, 179f, 185

sinus bradycardia with, 154f, 179

sinus rhythm with, 133f, 150

sinus tachycardia with, 196f, 209f, 213, 217

Premature junctional contraction (PJC), **39, 39f**, 40f

artificial pacemaker with, 206f, 216

bigeminal

junctional rhythm with, 160f, 180

normal sinus rhythm with, 125f, 148, 205f, 210f, 216, 217

in first-degree atrioventricular block, 59f

interpolated, junctional rhythm with, 167f, 182

junctional rhythm with, 142f, 152

normal sinus rhythm with, 195f, 213

sinus bradycardia with, 157f, 180

sinus rhythm with, 114f, 121

Premature ventricular contraction (PVC), **45, 45f**

accelerated junctional rhythm with, 41, 41f

artificial pacemaker with, 67, 195f, 203f, 213, 215

atrial and ventricular pacemaker with, 203f, 215

atrial fibrillation with, 129f, 149

atrial pacemaker with, 176f, 184

atrial tachycardia with, 53f

bigeminal, **46f**

accelerated idioventricular rhythm with, 145f, 153

normal sinus rhythm with, 128f, 148, 165f, 182, 202f, 215

sinus rhythm with, 138f, 151

couplet, **47, 47f**

atrial fibrillation with, 158f, 180

atrial flutter with, 127f, 148

normal sinus rhythm with, 167f, 182

sinus tachycardia with, 92f, 116, 145f, 153

third-degree atrioventricular block with, 110f, 120

first-degree atrioventricular block with, 61f

idioventricular rhythm with, 54f

interpolated, **48, 48f**

normal sinus rhythm with, 142f, 144f, 152

sinus bradycardia with, 97f, 117, 169f, 183, 205f, 216

junctional rhythm with, 189f, 212

junctional tachycardia with, 40, 40f, 42, 42f

multiform, **46f**, 52f

normal sinus rhythm with, 88f, 94f, 105f, 115, 116, 119

sinus rhythm with, 52f, 168f, 182

third-degree atrioventricular block with, 147f, 153

normal sinus rhythm with, 93f, 100f, 116, 118, 132f, 149, 164f,

166f, 176f, 181, 182, 184, 195f, 202f, 213, 215

quadrigeminal, **47, 47f**

sinus rhythm with, 197f, 214

R on T phenomenon with, **47, 47f**

sinus bradycardia with, 97f, 117, 169f, 183, 187f, 211

sinus rhythm with, 52f, 72f, 126f, 148

sinus tachycardia with, 140f, 151, 159f, 180, 188f, 190f, 211, 212

trigeminal, **46f**

sinus rhythm with, 111f, 120, 158f, 161f, 175f, 180, 181, 184

triplet, normal sinus rhythm with, 123f, 134f, 147, 150, 166f,

176f, 182, 184, 201f, 215

uniform, **46f**

sinus rhythm with, 170f, 183

Procainamide, 263

Pronestyl, 263

Propranolol, 260

PSVT. *See* Paroxysmal supraventricular tachycardia (PSVT)

Pulmonary artery, 3, 3f

Pulmonary vein, 3, 3f

Pulseless electrical activity (PEA), **50–51, 50f**

advanced cardiac life support protocol for, 254

case study of, 226–228, 226f, 227f, 228f, 231

Pulseless ventricular tachycardia, advanced cardiac life support

protocol for, 253–254

Purkinje system, 7, 7f

intrinsic rate of, 63

PVC. *See* Premature ventricular contraction (PVC)

Q

Q wave, 14, 14f

QRS complex, 14, 14f, 17

QRS interval, 16t

with artificial cardiac pacemaker, 64

QT interval, 14, 14f, 16t

Quadrigeminy

- atrial, sinus tachycardia with, 173f, 184
ventricular, **47, 47f**
normal sinus rhythm with, 197f, 214
- R**
- R on T phenomenon, 41f, 42f, **47, 47f**, 165f, 182
- R-R interval, 16
measurement of, 17
- R wave, 14, 14f
progression of, 78, 78f
- Radial artery, 4, 5f
- Ramipril, 259
- Rapid ventricular response, atrial fibrillation with, 103f, 107f, 118, 119, 146f, 153, 154f, 155f, 179, 187f, 190f, 211, 212
- Rate, 16t
- Regularity, 16t
- Repolarization, 7, 8f
on electrocardiogram, 9, 9f
- Respiration artifact, **71, 71f**
sinus arrhythmia with, 89f, 115
sinus bradycardia with, 97f, 117
sinus pause with, 135f, 150
ventricular pacemaker with, 135f, 150
- Retenol, 261
- Rhythm
agonal, **44**, 163f, 181, 192f, 211f, 212, 217
idioventricular. *See* Idioventricular rhythm
indeterminate, 188f, 211
junctional. *See* Junctional rhythm
sinus. *See* Sinus rhythm
ventricular, accelerated
atrial flutter with, 141f, 152
unknown underlying rhythm with, 141f, 152
- Rhythm strip, 13, 14f
analysis of, 16–17, 16t
- Right bundle branch block, **82, 83f**. *See also* Bundle branch block (BBB)
- Right-sided electrocardiogram, 12–13, 13f, 13t
- S**
- S wave, 14, 14f
- SA. *See at* Sinoatrial
- Second-degree atrioventricular (AV) block
type I (Mobitz I, Wenckebach), **57, 57f**, 59f, 60f, 90f, 93f, 115, 116, 177f, 185
type II (Mobitz II), **57, 57f**, 59f, 60f, 191f, 212
3:1 block with, 143f, 152, 209f, 217
bundle branch block with, 209f, 217
- Segment, 14, 14f
ST. *See* ST segment
- Semilunar valves, 3, 3f
- Septal myocardial infarction, 79f, **81–82, 81f**
- Sinoatrial block, **23, 23f**, 26f
normal sinus rhythm with, 201f, 215
sinus rhythm with, 111f, 120
sinus tachycardia with, 192f, 212
- Sinoatrial node, 7, 7f, 63
- Sinoatrial node arrhythmias, 20–27, 20f–26f. *See also specific arrhythmias*
- Sinus arrhythmia, **22, 22f**, 25f, 122f, 147, 172f, 183, 202f, 215
respiration artifact with, 89f, 115
sinus bradycardia with, 127f, 141f, 148, 152, 198f, 205f, 214, 216
supraventricular tachycardia conversion from, 130f, 149
- Sinus block
normal sinus rhythm with, 94f, 116
sinus tachycardia with, 129f, 149
- Sinus bradycardia, **21, 21f**, 23f, 34f, 86f, 101f, 115, 118, 132f, 149, 175f, 184, 189f, 212
- advanced cardiac life support protocol for, 255–256
bundle branch block with, 172f, 183, 194f, 200f, 213, 214
case study of, 228–229, 228f, 229f, 231
first-degree atrioventricular block with, 156f, 163f, 172f, 178f, 179, 181, 183, 185, 208f, 216
interpolated premature ventricular contraction with, 97f, 117, 156f, 169f, 179, 183, 205f, 216
inverted T waves with, 87f, 115, 154f, 156f, 179, 198f, 205f, 214, 216
monomorphic ventricular tachycardia conversion from, 145f, 153
multiform premature ventricular contraction with, 156f, 179
muscle artifact with, 74f
normal sinus rhythm with, 200f, 214
premature atrial contraction with, 154f, 179
premature junctional contraction with, 157f, 180
premature ventricular contraction with, 97f, 117, 169f, 183, 187f, 211
respiration artifact with, 97f, 117
sinus arrhythmia with, 127f, 141f, 148, 152, 198f, 205f, 214, 216
sinus pause with, 126f, 148
sinus tachycardia with, 191f, 212
ST segment depression with, 87f, 115, 154f, 179
ST segment elevation with, 205f, 216
ventricular pacemaker conversion from, 171f, 183
- Sinus pause (sinus arrest), **22, 22f**, 25f, 102f, 118, 123f, 147
normal sinus rhythm with, 138f, 151, 162f, 174f, 181, 184
respiration artifact with, 135f, 150
sinus bradycardia with, 126f, 148
sinus rhythm with, 123f, 147
ventricular tachycardia with, 206f, 216
- Sinus rhythm, 204f, 215. *See also* Normal sinus rhythm
bigeminal premature ventricular contraction with, 138f, 151
bundle branch block with, 61f
first-degree atrioventricular block with, 131f, 149
inverted T waves with, 164f, 181
junctional escape beat with, 164f, 181
multiform premature ventricular contraction with, 52f, 168f, 182
muscle artifact with, 72f, 73f, 171f, 183
paroxysmal supraventricular tachycardia conversion to, 155f, 179
premature atrial contraction with, 133f, 150
premature junctional contraction with, 114f, 121
premature ventricular contraction with, 72f, 126f, 148
sinoatrial block with, 111f, 120
sinus pause with, 123f, 147
sixty-cycle interference with, 138f, 151
ST segment depression with, 24f, 26, 61f, 131f, 133f, 149, 150, 164f, 181
ST segment elevation with, 91f, 116
case study of, 239–241, 239f, 240f, 241f, 244
third-degree atrioventricular block conversion to, 242, 242f
ventricular bigeminy with, 138f, 151
ventricular fibrillation conversion to, 222, 222f
ventricular tachycardia conversion from, 104f, 119
ventricular tachycardia conversion to, 239, 239f
ventricular trigeminy with, 111f, 120
- Sinus tachycardia (ST), **21, 21f**, 24f, 25f, 87f, 88f, 110f, 115, 120, 128f, 148, 157f, 180
artifact with, 109f, 120
atrial quadrigeminy with, 173f, 184
couplet premature ventricular contraction with, 92f, 116, 145f, 153
inverted T waves with, 140f, 151, 159f, 180, 209f, 217
paroxysmal supraventricular tachycardia conversion from, 169f, 183
peaked T waves with, 110f, 120, 200f, 214
polymorphic ventricular tachycardia with, 188f, 211
premature atrial contraction with, 196f, 209f, 213, 217

- Sinus tachycardia (ST) (*Continued*)
 premature ventricular contraction with, 140f, 151, 159f, 180, 188f, 190f, 211, 212
 sinoatrial block with, 192f, 212
 sinus block with, 129f, 149
 sinus bradycardia with, 191f, 212
 ST segment depression with, 92f, 116, 130f, 140f, 149, 151, 176f, 184, 188f, 190f, 209f, 211, 212, 217
 supraventricular tachycardia with, 209f, 217
- Sixty-cycle interference artifact, **72, 72f**
 atrial fibrillation with, 73f
 sinus rhythm with, 138f, 151
 torsade de pointes with, 74f
- Slow ventricular response
 atrial fibrillation with, 162f, 181, 193f, 213
 atrial flutter with, 198f, 214
- Sodium bicarbonate, 263
- ST segment, 14, 14f. *See also* ST segment depression; ST segment elevation
 baseline, 79, 79f
- ST segment depression, 80, 80f
 accelerated junctional rhythm with, 199f, 214
 artificial pacemaker with, 67f, 123f, 147, 203f, 215
 atrial and ventricular pacemaker with, 203f, 215
 atrial fibrillation with, 92f, 98f, 116, 117, 134f, 150, 155f, 160f, 179, 180, 190f, 212
 atrial flutter with, 35f
 atrial pacemaker with, 123f, 147
 atrial tachycardia with, 33f
 bundle branch block with, 61f
 junctional rhythm with, 134f, 142f, 150, 152, 207f, 216
 junctional tachycardia with, 42f
 normal sinus rhythm with, 132f, 139f, 151, 169f, 183, 196f, 201f, 202f, 210f, 213, 215, 217
 on pacemaker electrocardiogram, 69f
 sinus bradycardia with, 87f, 115, 154f, 179
 sinus rhythm with, 24f, 26, 61f, 131f, 133f, 149, 150, 164f, 181
 sinus tachycardia with, 92f, 116, 130f, 140f, 149, 151, 176f, 184, 188f, 190f, 209f, 211, 212, 217
 third-degree atrioventricular block with, 60f
- ST segment elevation, 79–80, 80f
 accelerated junctional rhythm with, 99f, 117
 atrial fibrillation with, 158f, 180
 junctional rhythm with, 41f
 in myocardial infarction, 79–80, 80f
 normal sinus rhythm with, 99f, 118, 134f, 150, 186f, 192f, 206f, 211, 212, 216
 sinus bradycardia with, 205f, 216
 sinus rhythm with, 91f, 116
 case study of, 239–241, 239f, 240f, 241f, 244
- Starling's law, 6
- Streptokinase, 261
- Stroke volume, 6
- Subclavian artery, 4, 5f
- Subclavian vein, 5, 5f
- Superior vena cava, 5, 5f
- Supraventricular tachycardia (SVT), **30, 30f**, 33f, 112f, 120, 124f, 139f, 147, 151
 junctional rhythm with, 125f, 148
 normal sinus rhythm conversion from, 133f, 150
 normal sinus rhythm conversion to, 203f, 215
 paroxysmal, **31, 31f**, 34f, 35f
 accelerated junctional rhythm with, 143f, 152
 case study of, 233–235, 233f, 234f, 235f, 243
 sinus rhythm conversion from, 104f, 119
 sinus rhythm conversion to, 155f, 179
 sinus tachycardia conversion to, 169f, 183
 sinus arrhythmia with, 130f, 149
 sinus tachycardia with, 209f, 217
- SVT. *See* Supraventricular tachycardia (SVT)
- Sympathomimetics, 262
- Systole, 5, 6f
- T**
- T wave(s), 14, 14f
 inverted
 accelerated junctional rhythm with, 199f, 214
 artificial pacemaker with, 67f, 69f, 203f, 215
 atrial fibrillation with, 134f, 150
 atrial pacemaker with, 123f, 147
 junctional tachycardia with, 128f, 148
 normal sinus rhythm with, 99f, 118, 144f, 152, 201f, 210f, 215, 217
 on pacemaker electrocardiogram, 67f, 69f
 second-degree atrioventricular block with, 60f
 sinus bradycardia with, 87f, 115, 154f, 179, 198f, 205f, 214, 216
 sinus rhythm with, 164f, 181
 sinus tachycardia with, 140f, 151, 159f, 180, 209f, 217
- peaked
 normal sinus rhythm with, 99f, 118
 sinus tachycardia with, 110f, 120, 200f, 214
- Tachycardia
 atrial, **30, 30f**, 33f, 138f, 151
 multifocal, **29, 29f**
 junctional, **38, 38f**, 40f, 41f, 42f, 168f, 182
 bundle branch block with, 137f, 151
 couplet premature ventricular contraction with, 187f, 211
 inverted T waves with, 128f, 148
 narrow-complex, advanced cardiac life support protocol for, 257
 sinus. *See* Sinus tachycardia (ST)
 supraventricular. *See* Supraventricular tachycardia (SVT)
 unstable, advanced cardiac life support protocol for, 256–257
 ventricular. *See* Ventricular tachycardia (VT)
 wide-complex, advanced cardiac life support protocol for, 257–258
- Tenecteplase, 261
- Third-degree atrioventricular (AV) block, **58, 58f**, 60f, 61f, 95f, 100f, 117, 118, 159f, 165f, 168f, 180, 182, 193f, 207f, 208f, 213, 216
 case study of, 241–242, 241f, 242f, 244–245
 couplet premature ventricular contraction with, 110f, 120
 multiform premature ventricular contraction with, 147f, 153
 ventricular fibrillation conversion from, 139f, 151
 ventricular fibrillation conversion to, 156f, 179
- Thrombolytics, 261
- Torsade de pointes, **49–50, 49f**, 53f, 108f, 120, 173f, 184
 muscle artifact with, 75f
 sixty-cycle interference with, 74f
- Transcutaneous pacing, 267, 267f
- Tricuspid valve, 3, 3f
- Trigeminy, ventricular, **46f**
 normal sinus rhythm with, 158f, 161f, 175f, 180, 181, 184
 sinus rhythm with, 111f, 120
- Tunica externa, 4, 4f
- Tunica intima, 4, 4f
- Tunica media, 4, 4f
- Twelve-lead electrocardiogram, 77, **80, 80f**
- U**
- U wave, 14, 14f
 normal sinus rhythm with, 24f, 26, 100f, 118, 161f, 181
- Uniform premature ventricular contraction, **46f**
 sinus rhythm with, 170f, 183
- V**
- Vagal maneuver, 267, 267f
- Variable block, atrial flutter with, 100f, 118, 136f, 146f, 150, 153, 188f, 198f, 211, 214

- Vasopressin, 264
 Vasopressors, 261, 264
 Veins, 4, 5, 5f
 Ventricles, 3, 3f
 Ventricular arrhythmias, 44–55, 44f–54f. *See also specific arrhythmias*
 Ventricular asystole, 207f, 216
 Ventricular bigeminy, **46, 46f**
 accelerated idioventricular rhythm with, 145f, 153
 normal sinus rhythm with, 165f, 182, 189f, 202f, 212, 215
 sinus rhythm with, 138f, 151
 Ventricular contraction, premature. *See* Premature ventricular contraction (PVC)
 Ventricular couplet
 sinus tachycardia with, 176f, 184
 unknown underlying rhythm with, 141f, 152
 Ventricular fibrillation (VF), **50, 50f**, 53f, 94f, 101f, 116, 118, 137f, 151, 197f, 208f, 209f, 214, 216, 217
 advanced cardiac life support protocol for, 253–254
 asystole conversion from, 126f, 129f, 148, 149
 asystole conversion to, 156f, 179, 190f, 212
 atrial fibrillation conversion with, 95f, 117
 case study of, 220–222, 220f, 221f, 222f, 229
 defibrillation with, 124f, 147
 normal sinus rhythm conversion from, 195f, 213
 sinus rhythm conversion from, 222, 222f
 third-degree atrioventricular block conversion from, 156f, 179
 third-degree atrioventricular block conversion to, 139f, 151
 ventricular tachycardia conversion from, 132f, 149
 ventricular tachycardia conversion to, 105f, 119, 124f, 147, 174f, 184
 wandering atrial pacemaker conversion to, 175f, 184
 Ventricular quadrigeminy, **47, 47f**
 normal sinus rhythm with, 197f, 214
 Ventricular rhythm, accelerated
 atrial flutter with, 141f, 152
 unknown underlying rhythm with, 141f, 152
 Ventricular tachycardia (VT)
 artificial pacemaker with, 195f, 213
 case study of, 223–226, 224f, 225f, 226f, 230, 237–239, 237f, 238f, 239f, 244
 monomorphic, **48, 48f**, 51f, 52f, 106f, 119, 144f, 152, 198f, 214
 sinus bradycardia conversion to, 145f, 153
 multiform, atrial fibrillation with, 136f, 150
 normal sinus rhythm conversion from, 137f, 151, 225–226, 226f
 normal sinus rhythm with, 123f, 134f, 147, 150, 166f, 176f, 182, 184, 201f, 215
 polymorphic, **49, 49f**, 113f, 121, 136f, 150
 sinus tachycardia with, 188f, 211
 ventricular pacemaker with, 173f, 184
 vs. artifact, 108f, 120
 pulseless, advanced cardiac life support protocol for, 253–254
 sinus pause with, 206f, 216
 sinus rhythm conversion from, 239, 239f
 sinus rhythm conversion to, 104f, 119
 ventricular fibrillation conversion from, 105f, 119, 124f, 147, 174f, 184
 ventricular fibrillation conversion to, 132f, 149
 vs. artifact, 191f, 212
 Ventricular trigeminy, **46, 46f**
 normal sinus rhythm with, 158f, 161f, 175f, 180, 181, 184
 sinus rhythm with, 111f, 120
 Verapamil, 264
 VF. *See* Ventricular fibrillation (VF)
 VT. *See* Ventricular tachycardia (VT)
- W**
- Wandering atrial pacemaker, **28, 28f**, 107f, 119, 172f, 175f, 183, 184
 Wave, 14, 14f
 P, 14, 14f, 16t, 17
 artificial pacemaker with, 64
 Q, 14, 14f
 R, 14, 14f
 progression of, 78, 78f
 S, 14, 14f
 T, 14, 14f. *See also* T wave(s)
 U, 14, 14f
 sinus rhythm with, 24f, 100f, 118, 161f, 181
 Wenckebach atrioventricular (AV) block, **57, 57f**, 59f, 60f, 90f, 93f, 115, 116, 177f, 185
 Wide intrinsic complexes, artificial pacemaker with, 210f, 217
 Wolff-Parkinson-White (WPW) syndrome, **32, 32f**
- X**
- Xylocaine, 262