

Vectors:

## Vector:


A vector is a matrix that has only one row – then we call the matrix a **row vector** – or only one column – then we call it a **column vector**.

A **row vector** is of the form:  $a = [a_1 \ a_2 \ \dots \ a_n]$


A **column vector** is of the form:

$$b = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix}$$

A quantity such as force, displacement, or velocity is called a vector and is represented by a directed line segment



A **vector** in the plane is directed line segment. The directed line segment  $\overrightarrow{AB}$  has **initial point** A and **terminal point** B; its **length** is denoted by  $|\overrightarrow{AB}|$ . Two vectors are **equal** if they have the same length and direction.



### Component form


If  $v$  is a **two dimensional** vector in the plane equal to the vector with initial point at the origin and terminal point  $(v_1, v_2)$ , then the **Component form** of  $v$  is:

$$v = (v_1, v_2)$$

If  $v$  is a **three dimensional** vector in the plane equal to the vector with initial point at the origin and terminal point  $(v_1, v_2, v_3)$ , then the **Component form** of  $v$  is:

Vectors:

$$v = (v_1, v_2, v_3)$$



The numbers  $v_1, v_2$  and  $v_3$  are called the components of  $v$ .

Given the points  $P(x_1, y_1, z_1)$  and  $Q(x_2, y_2, z_2)$ , the standard position vector

$v = (v_1, v_2, v_3)$  equal to  $\overrightarrow{PQ}$  is

$$v = (x_2 - x_1, y_2 - y_1, z_2 - z_1)$$

The **magnitude** or **length** of the vector  $v = \overrightarrow{PQ}$  is the nonnegative number

$$|v| = \sqrt{v_1^2 + v_2^2 + v_3^2} = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$$

The only vector with length **0** is the zero vector  $0 = (0,0)$  or  $0 = (0,0,0)$ . This vector is also the only vector with no specific direction.

**Ex.:** Find **a)** component form and **b)** length of the vector with initial point  $P(-3,4,1)$  and terminal point  $Q(-5,2,2)$

**Solution:**

**a)**  $v = (-5+3, 2-4, 2-1)$

The component form of  $\overrightarrow{PQ}$  is  $v = (-2, -2, 1)$

**b)** The length or magnitude of  $v = \overrightarrow{PQ}$  is  $|v| = \sqrt{(-2)^2 + (-2)^2 + (1)^2} = \sqrt{9} = 3$

**Vector Addition and Multiplication of a vector by a scalar**

Let  $u = (u_1, u_2, u_3)$  and  $v = (v_1, v_2, v_3)$  be vectors with  $k$  a scalar.


**Addition:**

$$u + v = (u_1 + v_1, u_2 + v_2, u_3 + v_3)$$

Vectors:



**Scalar multiplication:**  $ku = (ku_1, ku_2, ku_3)$

If the length of  $ku$  is the absolute value of the scalar  $k$  times the length of  $u$ .  
The vector  $(-1)u = -u$  has the same length as  $u$  but points in the opposite direction.



If  $u = (u_1, u_2, u_3)$  and  $v = (v_1, v_2, v_3)$ ,  $u - v = (u_1 - v_1, u_2 - v_2, u_3 - v_3)$

Note that  $(u - v) + v = u$  and the difference  $u - v$  as the sum  $u + (-v)$



**Ex.:**

Let  $u = (-1, 3, 1)$  and  $v = (4, 7, 0)$ , find

- a)  $2u + 3v$       b)  $u - v$       c)  $\left| \frac{1}{2}u \right|$

**Solution:**

a)  $2u + 3v = (-2, 6, 2) + (12, 21, 0) = (10, 27, 2)$

b)  $u - v = (-5, -4, 1)$

c)  $\left| \frac{1}{2}u \right| = \left| \left( \frac{-1}{2}, \frac{3}{2}, \frac{1}{2} \right) \right| = \frac{1}{2}\sqrt{11}$

**Properties of vector operations:**

Let  $u$ ,  $v$  and  $w$  be vectors and  $a$  and  $b$  be scalars.

1)  $u + v = v + u$

2)  $(u + v) + w = u + (v + w)$

Vectors:

- |                         |                         |
|-------------------------|-------------------------|
| 3) $u + 0 = u$          | 4) $u + (-u) = 0$       |
| 5) $0u = 0$             | 6) $1u = u$             |
| 7) $a(bu) = (ab)u$      | 8) $a(u + v) = au + av$ |
| 9) $(a + b)u = au + bu$ |                         |

**Unit vectors**

A vector  $v$  of length 1 is called **unit vector**. The standard unit vectors are:

$$i = (1,0,0) \quad , \quad j = (0,1,0) \quad , \quad k = (0,0,1)$$

$$\begin{aligned} v = (v_1, v_2, v_3) &= (v_1, 0, 0) + (0, v_2, 0) + (0, 0, v_3) \\ &= v_1(1, 0, 0) + v_2(0, 1, 0) + v_3(0, 0, 1) \\ &= v_1i + v_2j + v_3k \end{aligned}$$

We call the scalar (or number)  $v_1$  the ***i*-component** of the vector  $v$  ,  $v_2$  the ***j*-component** of the vector  $v$  , and  $v_3$  the ***k*-component**. In component form,


$P_1(x_1, y_1, z_1)$  and  $P_2(x_2, y_2, z_2)$  is

$$\overrightarrow{P_1P_2} = (x_2 - x_1)i + (y_2 - y_1)j + (z_2 - z_1)k$$

If  $v \neq 0$ , then

$$u = \frac{v}{|v|} \text{ is a unit vector in the direction of } v, \text{ called } \textit{the direction} \text{ of the}$$

nonzero vector  $v$ .



**Ex.:**

Find a unit vector  $u$  in the direction of the vector  $P_1(1,0,1)$  and  $P_2(3,2,0)$ .

**Solution**

$$\begin{aligned} \overrightarrow{P_1P_2} &= (3-1)i + (2-0)j + (0-1)k = 2i + 2j - k \\ |\overrightarrow{P_1P_2}| &= \sqrt{(2)^2 + (2)^2 + (-1)^2} = \sqrt{9} = 3 \end{aligned}$$

Vectors:


$$u = \frac{\overrightarrow{P_1P_2}}{|\overrightarrow{P_1P_2}|} = \frac{2i + 2j - k}{3} = \frac{2}{3}i + \frac{2}{3}j - \frac{1}{3}k$$

The unit vector  $u$  is the **direction** of  $\overrightarrow{P_1P_2}$ .

### Midpoint of a line segment

The Midpoint  $M$  of a line segment joining points  $P_1(x_1, y_1, z_1)$  and  $P_2(x_2, y_2, z_2)$  is the point

$$\left( \frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}, \frac{z_1 + z_2}{2} \right)$$



**Ex.:**

The midpoint of the segment joining  $P_1(3, -2, 0)$  and  $P_2(7, 4, 4)$  is

$$\left( \frac{3+7}{2}, \frac{-2+4}{2}, \frac{0+4}{2} \right) = (5, 1, 2)$$

### Product of vectors

$u$  &  $v$  are vectors,

There are two kinds of multiplication of two vectors:

- 1- The scalar product (dot product)  $u \cdot v$ . The result is a **scalar**.
- 2- The vector product (cross product)  $u \times v$ . The result is a **vector**.

#### 1) The dot product

In this section, we show how to calculate easily the angle between two vectors directly from their components. The dot product is also called **inner** or **scalar** products because the product results in scalar, not a vector.

Vectors:

**Def.:** The dot product  $u \cdot v$  of vectors  $u = (u_1, u_2, u_3)$  and  $v = (v_1, v_2, v_3)$  is:

$$u \cdot v = u_1v_1 + u_2v_2 + u_3v_3$$

**Note:**

$$\left. \begin{array}{l} i \cdot i \\ j \cdot j \\ k \cdot k \end{array} \right] = 1.1 = 1 \quad , \quad \left. \begin{array}{l} i \cdot j \\ j \cdot k \\ k \cdot j \end{array} \right] = 0$$

**Ex.:**

a)

$$(3,5) \cdot (-1,2) = 3(-1) + 5(2) = 7 \quad \text{scalar}$$

$$(3i + 5j) \cdot (-i + 2j) = 7$$

b)

$$(1,-3,4) \cdot (1,5,2) = 1 - 15 + 8 = -6 \quad \text{scalar}$$

$$(i - 3j + 4k) \cdot (i + 5j + 2k) = -6$$

**Angle between two vectors**

The angle  $\theta$  between two nonzero vectors  $u = (u_1, u_2, u_3)$  and  $v = (v_1, v_2, v_3)$  is given by

$$\vec{u} \cdot \vec{v} = |\vec{u}| \cdot |\vec{v}| \cdot \cos\theta$$

$$\theta = \cos^{-1} \left( \frac{\vec{u} \cdot \vec{v}}{|\vec{u}| \cdot |\vec{v}|} \right) \quad \text{where } \theta \quad (0 \leq \theta \leq \pi)$$

**Ex.:** Find the angle between two vectors in space

$$\vec{u} = 2\vec{i} - \vec{j} + 2\vec{k} \quad , \quad \vec{v} = \vec{i} - 2\vec{j} + 2\vec{k}$$


$$\cos\theta = \frac{\vec{u} \cdot \vec{v}}{|\vec{u}| \cdot |\vec{v}|} = \frac{2 + 2 + 4}{\sqrt{4+1+4} \cdot \sqrt{1+4+4}}$$

$$\cos\theta = \frac{8}{9} \Rightarrow \theta = \cos^{-1} \frac{8}{9}$$

Vectors:

**Ex.:**Find the angle  $\theta$  in the triangle ACB determined by the vertices

$$A = (0,0) , B(3,5) \text{ and } C(5,2)$$



$$\vec{CA} = (-5, -2) \quad \text{and} \quad \vec{CB} = (-2, 3)$$

$$\vec{CA} \cdot \vec{CB} = (-5)(-2) + (-2)(3) = 4$$

$$|\vec{CA}| = \sqrt{(-5)^2 + (-2)^2} = \sqrt{29}$$

$$|\vec{CB}| = \sqrt{(-2)^2 + (3)^2} = \sqrt{13}$$

$$\theta = \cos^{-1}\left(\frac{4}{\sqrt{29} \cdot \sqrt{13}}\right)$$

### ***Orthogonal vectors***

Vectors  $u = (u_1, u_2, u_3)$  and  $v = (v_1, v_2, v_3)$  are ***orthogonal*** (or ***perpendicular***)

if and only if  $u \cdot v = 0$

**Ex.:**

a)  $u = (3, -2)$  and  $v = (4, 6)$  are orthogonal because  $u \cdot v = 0$

b)  $u = 3i - 2j + k$  and  $v = 2j + 4k$  are orthogonal because  $u \cdot v = 0$

c)  $\mathbf{0}$  is orthogonal to every vector  $\mathbf{u}$  since

$$\begin{aligned} \mathbf{0} \cdot u &= (0, 0, 0) \cdot (u_1, u_2, u_3) \\ &= 0 \end{aligned}$$

### ***Properties of the Dot product***

If  $u$ ,  $v$  and  $w$  are any vectors and  $c$  is a scalar, then

1)  $u \cdot v = v \cdot u$

2)  $(cu) \cdot v = u \cdot (cv) = c(u \cdot v)$

Vectors:

3)  $u \cdot (v + w) = u \cdot v + u \cdot w$

4)  $u \cdot u = |u|^2$

5)  $0 \cdot u = 0$

**Vector projection**Vector projection of  $u$  onto  $v$ 

$$proj_v u = \left( \frac{u \cdot v}{|v|^2} \right) v \quad \dots\dots (1)$$

 $proj_v u$  ("The vector projection of  $u$  onto  $v$ ")**Ex.:**

Find the vector projection of  $u = 6i + 3j + 2k$  onto  $v = i - 2j - 2k$  and the scalar component of  $u$  in the direction of  $v$ .

**Solution:**We find  $proj_v u$  from eq.(1):

$$proj_v u = \left( \frac{u \cdot v}{|v|^2} \right) v = \frac{u \cdot v}{v \cdot v} v = \frac{6 - 6 - 4}{1 + 4 + 4} (i - 2j - 2k) = \frac{-4}{9} (i - 2j - 2k) = \frac{-4}{9} i + \frac{8}{9} j + \frac{8}{9} k$$

We find the scalar component of  $u$  in the direction of  $v$  from eq.(2):**Problems:**

1) Let  $u = (3, -2)$  and  $v = (-2, 5)$ . Find the **a)** component form and **b)** magnitude (length) of the vector.

1.  $-2u + 5v$

2.  $\frac{3}{5}u + \frac{4}{5}v$

2) Find the component form of the vector:

a. The vector  $\overrightarrow{PQ}$  where  $P = (1, 3)$  and  $Q = (2, -1)$ .b. The vector  $\overrightarrow{OP}$  where  $O$  is the origin and  $P$  is the midpoint of segment  $RS$ , where  $R = (2, -1)$  and  $S = (-4, 3)$ .c. The vector from the point  $A = (2, 3)$  to the origin.




Vectors:

d. The sum of  $\overrightarrow{AB}$  and  $\overrightarrow{CD}$ , where

$$A = (1, -1), B = (2, 0), C = (-1, 3) \text{ and } D = (-2, 2)$$

3) Let  $v$ ,  $u$  and  $w$  as in the figure: find a)  $u + v$ , b)  $u + v + w$ , c)  $u - v$  and d)  $u - w$



4) Find the vectors whose lengths and directions are given. Try to do the calculation without writing:

|    | <u>Length</u> | <u>Direction</u>                             |
|----|---------------|--|
| a. | 2             | $i$  |
| b. | $\sqrt{3}$    | $-k$   |
| c. | $\frac{1}{2}$ | $\frac{3}{5}j + \frac{4}{5}k$                |
| d. | 7             | $\frac{6}{7}i - \frac{2}{7}j + \frac{3}{7}k$ |

5) Find a) the direction of  $\overline{P_1P_2}$  and b) the midpoint of line segment  $P_1P_2$ .

- a.  $P_1(-1, 1, 5)$  and  $P_2(2, 5, 0)$
- b.  $P_1(0, 0, 0)$  and  $P_2(2, -2, -2)$

6) Find  $v \cdot u$ ,  $|v|$ ,  $|u|$ , the cosine of the angle between  $v$  and  $u$ , the scalar component of  $u$  in the direction of  $v$  and the vector  $proj_v u$ .

- a)  $v = 2i - 4j + \sqrt{5}k$ ,  $u = -2i + 4j - \sqrt{5}k$
- b)  $v = (\frac{3}{5})i + (\frac{4}{5})k$ ,  $u = 5i + 12j$
- c)  $v = -i + j$ ,  $u = \sqrt{2}i + \sqrt{3}j + 2k$
- d)  $v = 5i + j$ ,  $u = 2i + \sqrt{17}j$

Vectors:

$$\text{e) } v = \left( \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{3}} \right), \quad u = \left( \frac{1}{\sqrt{2}}, \frac{-1}{\sqrt{3}} \right)$$

7) Find the angles between the vectors:

$$\text{a) } u = 2i - 2j + k, \quad v = 3i + 4k$$

$$\text{b) } u = \sqrt{3}i - 7j, \quad v = \sqrt{3}i + j - 2k$$

$$\text{c) } u = i + \sqrt{2}j - \sqrt{2}k, \quad v = -i + j + k$$

8) Find the measures of the angles between the diagonals of the rectangle whose vertices are  $A = (1,0)$ ,  $B(0,3)$ ,  $C(3,4)$  and  $D(4,1)$ **References:**

- 1- Advanced Engineering Mathematics (Erwin Kreyszc)- 8<sup>th</sup> Edition.
- 2- Calculus (Howard Anton).
- 3- Advanced Mathematics for Engineering Studies (أ. رياض احمد عزت)